

USE OF SPOROTHECAE BY PHORETIC *TARSONEMUS* MITES TO
TRANSPORT ASCOSPORES OF CONIFEROUS BLUESTAIN FUNGI

BY JOHN C. MOSER

Southern Forest Experiment Station, USDA Pineville, LA 71360, U.S.A.

TRANS. BR. MYCOL. SOC.
84:750-753
June 1985

USE OF SPOROTHECAE BY PHORETIC *TARSONEMUS* MITES TO
TRANSPORT ASCOSPORES OF CONIFEROUS BLUESTAIN FUNGI

BY JOHN C. MOSER

Southern Forest Experiment Station, USDA Pineville, LA 71360, U.S.A.

Certain female *Tarsonemus* mites, like their host bark beetles, carry a specific fungus upon which the mites feed, inside a special morphological, spore-carrying structure called a sporotheca. The sporotheca is similar to the mycangium described for bark beetles, but differs in that no gland cells are present and the fungal spores do not multiply in the structure. So far, the phenomenon has been observed only in North America and China where the phoretic hosts of the mites are *Dendroctonus* and *Ips*.

The term sporotheca was first used by Suski (1973) (Peck) Wollenw., a grass pathogen with which the two *Siteroptes* were associated. A third species of *Siteroptes*, also associated with the fungal disease, did not possess the sporotheca. Two more types of sporothecae have apparently

evolved independently in females of the genera *Siteroptes* and *Trochometruidium* which belong to the closely related mite families Pygmephoridae and Pyemotidae, both of which belong to the subcohort Heterostigmae. *Trochometruidium tribulatum* Cross has sporotheca openings on coxal plates IV; Alfaro (1946) describes another sporotheca for a species of *Siteroptes* which carries one or two conidia of *Nigrospora oryzae* (Berk. & Br.) Petch, which causes lint rot of cotton. This type of sporotheca forms a pouch on the hysterosoma just behind the fourth pair of legs. Again, the mite and fungus are associated and the fungus is pathogenic to the plant. The *Siteroptes* feed on the mycelia of *N. oryzae*, which is required for the mite's normal growth and reproduction (Laemmlen & Hall, 1973).

Most bluestain diseases of conifers caused by *Ceratocystis* spp. are carried by scolytid beetles (Upadhyay, 1981). Although Leach, Orr & Christensen (1934), Hetrick (1949), and Moser, Wilkinson & Clark (1974) suspected that mites phoretic on these bark beetles might carry spores of *Ceratocystis* spp., Bridges & Moser (1983) were the first to provide good evidence. They reported that *Ceratocystis minor* (Hedgcock) Hunt associated with the

southern pine beetle, *Dendroctonus frontalis* Zimmerman, is hypervectored by two species of phoretic female mites of the family Tarsonemidae, subcohort Heterostigmae. Both mites (*Tarsonemus ips* Lindquist and *T. krantzi* Smiley & Moser) feed on *C. minor* and move from areas of fungus-colonized bark to cracks in the outer bark where they attach to emerging adult beetles (Roton, 1978). This report concerns mites carrying bluestain ascospores in a fourth type of sporotheca.

Several thousand mites have been microscopically examined for the presence of *C. minor* ascospores on the 15 mite species that are commonly phoretic on *Dendroctonus frontalis*, which attacks *Pinus* spp. in the southern United States.

Tarsonemus mites have also been scanned from adults of at least 50 other species of Scolytidae reared or trapped from various trees (mostly conifers) around the world.

The mites were mounted on slides and examined microscopically for *Ceratocystis* ascospores, and, if positive, whether they were concentrated on certain body surface areas of the mites.

Ascospores were abundant on body surfaces on individuals of all mite species removed from fruiting colonies of *C. minor* under bark (Bridges & Moser, 1983). However, *Tarsonemus ips* and *T. krantzi* were the only mites that carried ascospores in the phoretic state on *D. frontalis*.

The number of ascospores varied from 1 to over 100 on *Tarsonemus* mites, but they were usually located under one or both sides of tergite 1 (Fig. 1). The area under tergite 1 contains no unique morphological features to hold the spores and looks like the tergite 1 of other non-spore carrying species of *Tarsonemus*. But, functionally, the structure appears to fit the definition of a sporotheca as defined by Suski (1973) as 'paired external sacs used to transport fungal spores'. But this system is more primitive than those described for *Siteroptes* and *Trochometruidium*, because in *Tarsonemus* a unique morphological structure for carrying the spores never evolved.

The sporotheca is similar to the mycangium of scolytid beetles defined by Batra (1963) except that a sporotheca does not contain glands to nourish the fungi, and the fungi do not multiply within the sac.

Upadhyay (1981) splits the genus *Ceratocystis* into four sections: *Ips*, *Ceratocystis*, *Endoconidiophora*, and *Ophiostoma*. Because the ascospores of each have distinctive shapes, they can often be identified to section by viewing them microscopically. *C. minor* belongs to section *Ophiostoma*, the spores of which are often thin and crescent-shaped. Ascospores of *C. minor* and other species of *Ophiostoma* were almost always located in sporothecae of phoretic *Tarsonemus* mites; only rarely were these ascospores seen in a random fashion on




Fig. 1. Female of *Tarsonemus ips* showing the location of tergite 1 (arrows).



Fig. 2. Ventral surface of *Tarsonemus ips* female phoretic on *Dendroctonus frontalis* from *Pinus taeda*, Louisiana. Note the numerous ascospores of *Ceratocystis minor* in sporothecae (arrows) formed by tergite 1 on lateral sides of the mite.

body surfaces of *Tarsonemus* and other phoretic mites. Ascospores of all of the other sections except *Endocomidiophora* were seen on the body surfaces of various phoretic mites, but none occurred under tergite 1 on any *Tarsonemus* or in structures that might be construed as sporothecae in other mites (Moser, unpubl.).

Crescent-shaped *Ophiostoma* ascospores in *Tarsonemus* sporothecae have been seen from localities other than Louisiana (Fig. 2). All were morphologically similar or identical to those of *C. minor*, but all were seen on mites that had been preserved in alcohol. For this reason their identities could not be confirmed by culturing. Hence, the following *Ceratocystis* names are speculative, based on literature references following most records: (1) *Ceratocystis minor* on *Tarsonemus krantzi* phoretic on *Dendroctonus frontalis* from *Pinus leiophylla* Schiede, Puebla, Mexico (Bridges & Moser, 1983); (2) *Ceratocystis minor* on *Tarsonemus krantzi* on

Dendroctonus frontalis from *Pinus oocarpa* Schiede, Cedros, Honduras (Bridges & Moser, 1983); (3) *Ceratocystis minor* and/or *C. nigrocarpa* Davidson on *Tarsonemus endophloeus* Lindquist and *T. ips* phoretic on *Dendroctonus brevicomis* LeConte from *Pinus ponderosa* Douglas ex Lawson, Bass Lake, California (Whitney, 1982); (4) *Ceratocystis minor* and/or *C. nigrocarpa* on *Tarsonemus* prob. n.sp. nr *endophloeus* phoretic on *Dendroctonus pseudotsugae* Hopkins from *Pseudotsuga menziesii* (Mirbel) Franco, Moscow, Idaho (Upadhyay, 1981); (5) *Ceratocystis minor* and/or *C. nigrocarpa* on *Tarsonemus* prob. *endophloeus* phoretic on *Ips paraconfusus* Lanier from *Pinus coulteri* D. Don, Lake Arrowhead, California; (6) *Ceratocystis piceae* (Munch) Bakshi on *Tarsonemus* n.sp. nr *endophloeus* phoretic on *Ips nitidis* Eggers from *Picea asperata* Masters, Ginsu Provence, China (H. P. Upadhyay, pers. comm., agrees that *piceae* may be this species of *Ceratocystis*) (Fig. 3). Thus the sporothecae that

Fig. 3. Ventral surface of *Tarsonemus* n.sp. nr *endophloeus* female phoretic on *Ips nitidis* from *Picea asperata*, Ginsu Provence, China. Note the 61 ascospores of *Ceratocystis* sp. in sporothecae (arrow) formed by tergite 1 on lateral sides of the mite.




Fig. 3. For caption see opposite.

carry *Ceratocystis* ascospores on conifers are found on just a few species of *Tarsonemus*, but are associated with at least two genera of bark beetles on a rather broad range of host trees, and the phenomenon may be widespread geographically.

I wish to thank E. E. Lindquist and R. N. Sinha for helpful discussions, D. Barstow, J. H. Cain, J. F. Coyne, W. E. Rose, P. E. Tilden, and H.-f. Yin, for collection of bark beetles which contained the mites, and G. W. Krantz and H. P. Upadhyay for reviewing the manuscript.

This article was written and prepared by a U.S. Government employee on official time and it is therefore in the public domain.

REFERENCES

- ALFARO, A. (1946). El acaro *Pediculopsis graminum* Rent. y el hongo *Nigrospora oryzae* (Berk. et B.) Petch, en asociacion parasitaria sobre trigos aragonensis. *Boletín de Patología Vegetal Entomología Agrícola* **14**, 321-334.
- BATRA, I. R. (1963). Ecology of ambrosia fungi and their dissemination by beetles. *Transactions Kansas Academy of Science* **66**, 213-236.
- BRIDGES, J. R. & MOSER, J. C. (1983). Role of two phoretic mites in transmission of bluestain fungus, *Ceratocystis minor*. *Ecological Entomology* **8**, 9-12.
- HETRICK, L. A. (1949). Some overlooked relationships of southern pine beetle. *Journal of Economic Entomology* **42**, 466-469.
- LAEMMLEN, F. F. & HALL, D. H. (1973). Interdependence of a mite, *Siteroptes reniformis*, and a fungus, *Nigrospora oryzae*, in the nigrospora lint rot of cotton. *Phytopathology* **63**, 308-315.
- LEACH, J. G., ORR, L. W. & CHRISTENSEN, C. (1934). The interrelationships of bark beetles and blue-staining fungi in felled Norway pine timber. *Journal of Agricultural Research* **49**, 315-341.
- MOSER, J. C., WILKINSON, R. C. & CLARK, E. W. (1974). Mites associated with *Dendroctonus frontalis* Zimmerman (Scolytidae: Coleoptera) in Central America and Mexico. *Turrialba* **24**, 379-381.
- ROTON, L. M. (1978). Mites phoretic on the southern pine beetle: when and where they attach. *Canadian Entomologist* **110**, 557-558.
- SUSKI, Z. W. (1973). A revision of *Siteroptes cerealium* (Kirchner) complex (Acarina, Heterostigmata, Pycnotidae). *Annales Zoologici* **30**, 509-517.
- UPADHYAY, H. P. (1981). *A monograph of Ceratocystis and Ceratocystiopsis*. The University of Georgia Press: Athens.
- WHITNEY, H. S. (1982). Relationships between bark beetles and symbiotic organisms. In *Bark Beetles in North American Conifers* (ed. J. B. Mitton & K. B. Sturgeon). University of Texas Press: Austin.