

United States  
Department of  
Agriculture  
Forest Service




Southeastern Forest  
Experiment Station

General Technical  
Report SE-37

## A Loblolly Pine Management Guide

# Foresters' Primer in Nutrient Cycling

Jacques R. Jorgensen  
Carol G. Wells



February 1986

Southeastern Forest Experiment Station  
200 Weaver Boulevard  
P.O. Box 2680  
Asheville, North Carolina 28802

# **Foresters' Primer in Nutrient Cycling**

Jacques R. Jorgensen  
and  
Carol G. Wells

Soil Scientists  
Southeastern Forest Experiment Station  
Research Triangle Park, North Carolina

**A Loblolly Pine Management Guide**



## ABSTRACT

The nutrient cycle, which includes the input of nutrients to the site, their losses, and their movement from one soil or vegetation component to another, can be modified by site preparation, rotation length, harvest system, fertilization, and fire, and by using soil. We report how alternative procedures affect site nutrients, and we provide general principles that can be followed to enhance long-term productivity of loblolly pine.

**Keywords:** Pinus taeda, nitrogen, phosphorus, potassium, rotation length, nitrogen fixing, whole-tree harvesting.

---

One key to sustained yield in forestry is the maintenance or improvement of soil fertility through good soil management. Such management requires understanding of nutrient cycling, which is the circulation of nutrients in the forest ecosystem. In each ecosystem, circulation of nutrients follows specific pathways. Cycling patterns can only be described in broad terms because rates and directions of movement vary with stand conditions. Some general processes regulating nutrient flow within a forest system can be identified, however: (1) uptake by higher plants, (2) translocation and use within the plants, (3) return to the soil and forest floor, (4) mineralization, immobilization, and leaching of the returned nutrients, (5) inputs of nutrients from atmospheric, geologic, and biological sources, and (6) losses in streamflow, harvesting, and volatilization to the atmosphere.

Concerns about potential declines in forest productivity through loss of soil nutrients are not new. Over a century ago, experiments determined that excessive removal of nutrient-containing litter by Bavarian peasants was detrimental to tree growth. When this adverse effect was clearly established, litter gathering was discontinued.

Despite that experience, the prevailing view of silviculturists until recently was that mineral elements would "take care of themselves" or that the loss of nutrient elements through the exploitation of timber was not very serious if leaves and twigs were left behind. In 1955, an analysis of the requirements for forest growth on a poor site in England showed demands for nutrients exceeded the supply. Few noticed that this forest soil was similar to many forest soils in the South-eastern United States. Only recently have southern foresters recognized that conservation and replenishment of nutrients are as important in forest management as is determination of rotation length or thinning schedule.

Pines are adapted to infertile soils in part because they require smaller amounts of nutrients than do other plants. Recent work, however, has revealed some additional possible mechanisms: (1) conservation of available nutrients by minimizing losses to deeper soil layers or to the atmosphere (tight nutrient cycles), (2) minimizing losses from the cycle by temporarily immobilizing nutrients, primarily in the soil organic layers, (3) collection and retention of nutrients from inputs to the cycle (from the atmosphere and weathering), (4) reuse of nutrients within the tree, and (5) evolution of special mechanisms, such as mycorrhizae, that obtain nutrients from sources not readily available. Regardless of the mechanisms used, the pine ecosystem, like those of other plants, is sensitive to disturbances that might reduce its ability to obtain, retain, or cycle nutrients. Disturbances such as harvesting, fertilization, burning, and changes in atmospheric inputs all affect cycling. Understanding of these processes and their interrelationships, therefore, is necessary for maintenance of forest productivity.

## Ecosystem Components

The forest ecosystem can be divided into three major components--mineral soil, forest

floor, and vegetation. Each major component can be further divided into smaller components or pools (fig. 1). With data for each pool and information on the influence of management practices, flow models can be developed that show the rate of nutrient transfer from one pool to another. Predictions of nutrient availability, and thus production, can be made based upon changes in the pool quantities and on the transfer rates between pools.

A number of factors complicate such predictions at any given time, however. For example, relatively small additions by biological nitrogen




Figure 1.--The nutrient cycle in a loblolly pine stand.

(N) fixation, or nutrient losses by leaching, may influence growth for several rotations of trees. Therefore, extrapolation of the effects of extensive treatments, such as fertilization, over long time periods should be done cautiously and with some knowledge of the potential for error.

Quantitative nutrient cycling data are needed, however, to address some major forestry problems with harvesting systems, fertilization, and environmental protection. Currently, research in fertilization is primarily directed toward increased yield and toward delineation of sites where trees will respond to fertilization. In the long term, forest fertilization practices based upon a knowledge of nutrient cycling will sustain or improve yields.

Each nutrient or element has its own peculiar system of cycling, but the systems have similarities. N is largely in organic compounds and requires biological action for its release. Phosphorus (P) is in both organic and inorganic materials and its availability is strongly influenced by chemical reactions. Potassium (K) is almost totally inorganic, and is in equilibrium between the native, fixed, and exchangeable forms. Due to these and other variations, there is no single cycle: Rather, there are many cycles, one for each element under each separate condition.

From this general introduction, it is evident that cycling can be altered by modifications of the forest environment. We shall examine the components of the cycles and the influences of management practices thereon. Because it has been shown that loblolly pine (Pinus taeda L.) responds most readily to N and P fertilization, emphasis will be placed on these two elements.

## Nutrient Inputs to the Forest

In nature, nutrients are added to a loblolly pine system from: (1) the atmosphere as dry gases

and particulates or as materials dissolved in wet precipitation, (2) weathering of soil minerals, and (3) biological fixation of N.

### Atmospheric Inputs

Precipitation contributes significant amounts of nutrients for growth and maintenance of loblolly pine. Average annual inputs per acre (ha) throughout the tree's range are 5.4 lb (6 kg) of N, 0.4 lb (0.4 kg) of P, 1.5 lb (1.7 kg) of K, 6.4 lb (7.2 kg) of calcium (Ca), and 1.5 lb (1.7 kg) of magnesium (Mg). Variation between areas may be great; areas of high precipitation tend to have more nutrient inputs than those with low precipitation. Other factors may influence the chemical composition of precipitation. Areas near the ocean can receive marine aerosols that are high in sodium, magnesium, chloride, and sulphate. Sites near urban and industrial areas can receive increased quantities of sulphate sulfur and nitrate and ammonium N. Around smelters, quantities of sulfur and minor elements may be sufficient to adversely affect plant growth. At worst, all but the most resistant vegetation may be destroyed.

Gaseous and particulate inputs in the absence of precipitation are usually referred to as dry deposition. Unfortunately, there are few quantitative measurements of dry deposition. Mechanisms related to dry deposition include filtering of atmospheric particles by tree canopies, absorption of water-soluble gases onto moist surfaces, and the direct gaseous uptake via leaf stomata. Sulfur dioxide and gaseous ammonia may also be directly absorbed by the soil.

### Weathering Inputs

Weathering refers to nutrient release from primary and secondary soil minerals by geochemical and biogeochemical processes. In the South, release of nutrients depends on soil parent material and may vary greatly between sites. In general,

the older and more deeply weathered the soil, the less will be the potential for nutrient release. On highly weathered soils, atmospheric deposition, rather than weathering, is the major source of natural replenishment for the small leaching losses that may occur.

### Biological N Fixation

Biological fixation can be an important source of N. N is fixed by legumes, nodulated nonlegumes, and, in some instances, by lichens that include blue-green algae. Except under unusual circumstances, free-living organisms--bacteria and algae--contribute only insignificant amounts of N. In loblolly pine stands with closed canopies, N fixation from all sources probably does not exceed a few pounds per acre annually. The major sources of this fixation are nodulated nonlegumes such as southern bayberry (Myrica cerifera L.) and many species of legumes. The latter are especially important for brief periods after fires. Fixation by free-living bacteria and algae is limited by energy sources, low soil pH, and nutrients; it usually contributes less than 1 pound per acre of N annually. Mycorrhizae have not been shown to fix N. The apparent fixation sometimes associated with mycorrhizae is accomplished by free-living organisms in the soil areas influenced by the roots and their mycorrhizal associates.

### Nutrient Pools and Cycling

Nutrient pools change rapidly after ecosystem disturbances such as wildfire, harvesting, and site preparation but again reach an equilibrium as stands mature (figs. 2, 3, 4).

### Understory Vegetation

The nutrient content of herbaceous biomass peaks soon after plantation establishment. This




Figure 2.--Biomass accumulation in an unthinned loblolly pine plantation during the first 20 years. (Adapted from Switzer and Nelson 1972.)

vegetation, an important sink for nutrients, is usually eliminated after stand closure, 5 to 8 years after establishment. Along with the residues from harvesting, it later provides important quantities of nutrients to the developing loblolly pine stand. In one plantation, aboveground herbaceous vegetation contained 67 lb (75 kg) of N, 7 lb (8 kg) of P, 9 lb (10 kg) of K, 21 lb (23 kg) of Ca, and 7 lb (8 kg) of Mg per acre (ha).

### Overstory Vegetation

Rapidly growing young pines quickly accumulate nutrients and develop a forest floor at the expense of the mineral soil and the herbaceous components of the stand. Accumulation of nutrients in the stand proceeds at a fast pace during the first 20 years, slows through age 40, and reaches an equilibrium thereafter (fig. 3). Maximum nutrient accumulation in foliage of loblolly stands occurs sometime after stand closure, at about 15 years of age, but before there is extensive suppression mortality. As trees are suppressed, foliar nutrient accumulation declines. In the stand as a whole, stem and branch nutrients accumulate relatively uniformly through age 40, at which time a new equilibrium is reached. A different pattern of nutrient accumulation occurs in individual surviving trees (fig. 4). For the survivors, foliage continues to accumulate nutrients at a uniform rate throughout the rotation. Stems and branches of these trees gradually increase their rates of nutrient accumulation in contrast to an equilibrium for the stand as a whole.

A vigorously growing loblolly pine plantation can tie up a large portion of the site's nutrients (table 1). Of 2,124 lb of N/acre (2,400 kg/ha) on a site, 286 lb (321 kg) or 13 percent were in 16-year-old trees. An equal amount of N was in the forest floor. Thus, 26 percent of the site N was immobilized 16 years after planting loblolly pine. The relatively large proportion of site N




Figure 3.--Accumulation of macronutrients in whole trees in loblolly pine stands on good sites. (Adapted from Switzer and others 1968.)




Figure 4.--Accumulation of N in fractions of loblolly pine stands and trees on good sites. (Adapted from Switzer and others 1968.)

in the mineral soil and its low availability explains why N fertilization can often increase growth even though there is an abundance of N on the site. The proportions of P, K, and other elements that are extractable vary widely with site, and reliable measures of their nutritional availability have not been developed.

Needles have relatively little of a tree's biomass, but they have the highest concentrations of nutrients. Among tree parts, stemwood has the lowest concentrations of nutrients. Nutrient concentrations in root biomass could be more accurately described if roots were separated into two categories: permanent woody roots for anchoring the tree and providing channels for conduction, and nonwoody ephemeral roots for collecting nutrients and moisture required for growth. The latter contain relatively high nutrient concentrations compared with the former.

Table 1.--Distribution of tree biomass and nutrients in a 16-year-old loblolly pine plantation in the North Carolina Piedmont

Component	Biomass	N	P <sup>a</sup>	K <sup>a</sup>
	<u>Tons/acre</u>	- - -	<u>Pounds/acre (%)</u>	- - -
<b>Trees</b>				
Needles	3.6	73	9	43
Branches	10.3	54	5	25
Stemwood	48.9	70	10	58
Stembark	6.8	32	4	21
Roots	16.2	57	15	54
Total tree	85.8	286 (13)	43 (11)	201 (34)
Forest floor	--	274 (13)	27 (7)	25 (4)
Mineral soil, 0-28 in.	--	1,564 (74)	331 (82)	360 (62)
Site total	--	2,124	401	586

<sup>a</sup>Values represent total quantities in vegetation and forest floor and extractable amounts in mineral soil.

## Forest Floor

Nutrients and organic matter accumulate in the forest floor as a loblolly pine stand develops. Litter accumulates most rapidly between the time the stand closes and the time when stand foliage is at a maximum, about 15 years of age. In one plantation, the accumulation rate was nearly 2 tons/acre/year (4 t/ha) between ages 13 and 16. Accumulated litter reaches an equilibrium of 13 to 16 tons/acre (30 to 35 t/ha) somewhere between stand age 16 and 30 (fig. 5).

Nutrient contents of forest floors in loblolly pine stands vary greatly, probably because of differences in climate and soil. The forest floor of a 15-year-old plantation in Mississippi contained only about one-third as much N and P, half as much K, and two-thirds as much Ca and Mg as did that of a 16-year-old North Carolina plantation. In the forest floors of five 15-year-old




Figure 5.--Accumulations of organic matter, N, P, K, Ca, and Mg in the forest floor of three plantations at various ages.

plantations in Virginia, there was slightly less N, P, K, and Mg, but more Ca than in the North Carolina plantation. Although the forest floors of the three Piedmont North Carolina plantations shown in figure 5 had approximately reached a biomass equilibrium at age 16, nutrient contents continued to change. Differences between plantations were probably caused by the influences of stand density and soil on initial needle composition and rate of decomposition.

Decomposition and nutrient release rates of forest floor materials are highest in the first year after litterfall, but each element or component changes at its own rate. During the first year, about 25 percent of the organic material, 50 percent of the P, 70 percent of the K, 25 percent of the Ca, and 57 percent of the Mg, but only 10




Figure 6.--Percentages of initial organic matter, N, P, K, Ca, and Mg remaining in the forest floor after 8 years of decomposition.

percent of the N, are lost. After 8 years of decomposition, about two-thirds of the organic material, 60 percent of the P, 90 percent of the K, 67 percent of the Ca, and 79 percent of the original quantity of Mg are lost. Net decrease of N during this period is only 27 percent (fig. 6).

Forest floors of closed, growing loblolly pine plantations, regardless of age, may be similar in weight, nutrient content, and, except for

N, nutrient release (tables 2, 3). Differences may occur, however, in the distribution of nutrients within the forest floor layers and in the proportions of nutrients released from the layers. In a 19-year-old plantation, the majority of biomass and nutrients were in forest floor materials less than 8 years old (table 2). In a 40-year-old stand, forest floor materials younger and older than 8 years each held about half of the nutrients. Except for N, however, the majority of nutrients are released from forest floor layers less than 8 years old, regardless of stand age (table 3). Older forest floor layers in a young stand accumulate rather than release elements. In the 40-year-old stand, nutrients in the forest floor layers older than 8 years of age are released, but more slowly than from younger material. Thus, it appears that until forest floor biomass is in equilibrium with litterfall, older layers do not contribute important amounts of nutrients for tree growth. Before about age 20, the older layers act as nutrient sinks or reserves for cycling at a later time.

Many factors control the accumulation rate and the point at which forest floor biomass decomposition and nutrient release come into equilibrium with litterfall. These equilibria differ

Table 2.--Components of the forest floor of two loblolly pine plantations

Component	Forest floor layer age (years)					
	1 through 8		Over 8		All ages	
	Plantation age (years)					
	19	40	19	40	19	40
	----- Pounds/acre -----					
Organic matter	22,276	18,259	3,623	11,055	25,899	29,314
Nitrogen	283	212	71	159	354	371
Phosphorus	23.7	15.6	9.2	16.9	32.9	32.5
Potassium	14.3	12.3	3.8	13.5	18.1	25.8
Calcium	94.2	62.1	21.6	43.2	115.8	105.3
Magnesium	12.8	8.9	4.0	9.6	16.8	18.5

Table 3.--Annual release (accumulation) of nutrients from the forest floors of two loblolly pine plantations

Nutrient	Forest floor layer age (years)					
	1 to 8		Over 8		All ages	
	Plantation age (years)					
	19	40	19	40	19	40
	- - - - - Pounds/acre - - - - -					
Nitrogen	13.1	9.1	2.5	16.6	15.6	25.7
Phosphorus	4.8	3.3	(0.4)	1.8	4.4	5.1
Potassium	11.2	10.1	(0.2)	1.2	11.0	11.3
Calcium	15.6	11.8	0.4	3.7	16.0	15.5
Magnesium	4.4	3.4	0.2	0.7	4.5	4.6

among nutrients. Understanding the agents controlling decomposition and nutrient release will lead to improved tree growth and a reduction in the need for fertilization under intensive forest management.

### Mineral Soil

Mineral soil contains a majority of the ecosystem's nutrient reserves. In one Piedmont plantation, about three-quarters of all the site's N was in the mineral soil to a depth of 28 inches (70 cm) (table 1). Comparisons of the proportions of P, K, Ca, and Mg in biomass plus forest floor and in the mineral soil are meaningless because the bases for element measurements differ. In the forest floor, total nutrients are measured; in the soil, only extractable forms are measured. Nutrient analyses of mineral soils have been adapted from agriculture. While appropriate for annual agronomic crops, these tests are often inadequate as indices of long-term nutrient supply in forest ecosystems. The tests are not capable of estimating the size of the mineral soil nutrient reservoir. They indicate only the soil's ability to release nutrients under a particular set of conditions. Even where total nutrients are

measured, as with N, availability can only be roughly gauged. Nitrogen mineralization and availability are the result of numerous uncontrolled, unforeseen, and unknown factors that affect the presence and activity of the microorganisms responsible for N transformation. Thus, soil tests provide only rough approximations of potential total nutrient supply.

During the first few years after loblolly pine is established on sites without a forest floor, most of the nutrients required for tree growth and accumulation of the forest floor are supplied by the mineral soil. In a loblolly pine plantation established on an old field in the South Carolina Piedmont, N decreased markedly throughout the surface 24 inches (60 cm) of mineral soil during the first 15 years (fig. 7). During the next 5 years, there was little decline. Declines over the first 15 years were also observed for extractable K, Ca, and Mg. Extractable P decreases were noted only to a depth of 3 inches




Figure 7.--Changes in N content of mineral soil in an old-field loblolly pine plantation over a 20-year period.

(8 cm), possibly because below this depth the extractable P concentrations were extremely low. Nutrient changes in the mineral soil of loblolly pine stands between 20 and 40 years of age appear to be small. During this time, the forest floor can provide the majority of nutrients required for growth. Lack of further nutrient drawdown in the mineral soil may also be due to the depletion of readily available reserves.

Little information is available on the recharge of mineral soil depleted of nutrients during the first stages of a stand's growth. If recharge occurs, it would be after about 40 years of age (figs. 4, 5). At this time, uptake of nutrients by vegetation may be more than balanced by a loss in dying vegetation. The excess nutrients released by the decomposing roots and forest floor could aid in the preparation of the mineral soil to become a nutrient source when a succeeding stand develops.

Declines of nutrients in the mineral soil of old fields after pine planting illustrate the importance of the soil's nutrient supply during plantation establishment. What is not answered, however, is whether this old-field situation is representative of what occurs when natural stands or plantations with a forest floor are regenerated. Treatments that diminish the forest floor as a source of nutrients may prolong the stand's reliance on the mineral soil. On infertile or intensively harvested sites, declines related to nutrient deficiencies may take place unless fertilizer is applied.

### The Loblolly Pine Nutrient Cycle

Within the stand, nutrients move between and within the three major components: the living vegetation, the forest floor, and the mineral soil. Some factors that determine from which source nutrients are obtained include age and stage of stand development, and amount and availability of nutrients in the component or subcomponent.

Nutrients are transferred from one component to another by trees through litterfall, crown leaching (throughfall), root death and exudation, and translocation from one organ to another. Other than movement from roots to crown, the most common form of nutrient translocation is from foliage to stems before leaf abscission. Translocation may also occur in stemwood where nutrients in 1- to 2-year-old wood are conveyed to newer growth. There is no information available on translocation of nutrients out of branches and roots before their death, but it appears--at least in fine roots as in needles--that translocation is an effective nutrient conservation measure.

Although each nutrient has unique component pools, relative pool sizes, and transfer rates between pools, the basic concepts of nutrient transfers are similar. Figure 8 shows a simplified N cycle in a 16-year-old loblolly pine plantation. Input of N to the system is primarily atmospheric, with precipitation contributing 4.8 lb/acre (5.4 kg/ha) annually to the site. An unknown amount of N is added through dry fallout and by the direct absorption of ammonia by the foliage and the forest floor. Leaching from new and old needles by precipitation brings an additional 3.7 lb (4.1 kg) to the forest floor in throughfall, making a total input to the forest floor of 8.5 lb (9.5 kg) of dissolved N. At age 16, the forest floor contains 274 lb of N/acre (307 kg/ha) and has annual inputs from the atmosphere, leaves, branches, and fixation that total 43.5 lb (48.9 kg). Additional inputs are 43.4 lb (49 kg) from roots in both the forest floor and mineral soil. Total inputs to the forest floor and mineral soil are 86.9 lb (97.6 kg), but uptake by the trees amounts to 93.3 lb (104.6 kg). Thus, there is a net loss of N from soil and forest floor of 6.4 lb (7 kg). Approximately half of the N taken up by roots is translocated to the above-ground portions of the tree. Most of the other




Figure 8.--The N cycle in a 16-year-old loblolly pine plantation. Values in boxes indicate N storage in individual components. Values outside boxes indicate annual N transfer. All values are in pounds per acre.

portion is recycled one or more times into short-lived nonwoody feeder roots. Little is known about these roots concentrated in the lower forest floor and upper mineral soil, except that they appear to have several cycles of biomass increase and decline during a year. Even less information is available on whether or not nutrients are translocated from the ephemeral roots to more permanent structures to be used as reserves when a new growth cycle is begun.

Of the 48.5 lb (54.4 kg) of N translocated into the aboveground portions of the tree, 4.7 lb (5.4 kg) are retained in the stem biomass, and 43.8 lb (49.1 kg) are initially moved into branches to be augmented by 15.2 lb (17 kg) translocated from older needles just before their fall. Of the total branch input of 59.0 lb (66.1 kg), 4.0 (4.5) are lost through branch death, 5.9 (6.6) are retained for branch growth, and 49.1 lb (55 kg) are used for the production of new leaves. The 15.2 lb (17 kg) translocated from old needles before fall, if used for new needles, will provide about one-third of the N needs. During the next growing season, the cycle is renewed. New leaves and other components are grown, leached, aged, and lost, and the cycle begins again. As the stand ages, component sizes and their transfer rates change. Stem N will continue to gain until stand breakup, whereas foliage N will decrease slowly. As the stand ages, the annual nutrient needs decrease and the accumulated deficit in the mineral soil should be gradually restored.

## Silvicultural Practices Affecting the Nutrient Cycle

The most intensive forest management in North America is practiced in the Southeast, where loblolly pine is grown in 20- to 30-year rotations. Site preparation, thinning, fertilization, and various degrees of harvesting all affect the cycling of nutrients and the ability of the site

to maintain its productivity. Nutritional relationships should be considered when management alternatives such as rotation length, harvest intensity, prescribed fire, site preparation, and fertilization are selected to meet a particular objective.

### Effects of Harvesting on Nutrient Removal

Past forestry practices of long rotations and harvesting only the bole of the tree have led silviculturists to ignore nutrient removal and believe that nutrients take care of themselves. New practices are forcing silviculturists to think again. Intensive harvesting, in which more than the bole is removed and trees are grown in short rotations, is contemplated as a means to substantially increase biomass production for fiber, fuel, and chemicals. It is apparent that substantial quantities of so-called waste materials (branches and foliage) are available and could be utilized. More complete utilization does more than increase the biomass supply; it also can improve site esthetics, reduce site preparation and planting costs, and decrease the danger of wildfires or the need for prescribed burning. However, there are negative aspects that require examination. Increased nutrient removals along with the increased biomass removals may cause a decline in site productivity. Residual nutrients may also be lost more readily by leaching in intensively managed stands.

There is concern that complete-tree harvesting combined with short rotations can deplete the nutrient capital of a site and hence reduce future production. The quantities of nutrients removed are affected by the density of the particular stand, the age and stage of development, and the biomass components that are harvested. The N losses associated with aboveground biomass harvests and stem harvest in 16- and 32-year-old

plantations are shown in table 4. Annual nutrient removal is more affected by intensity of harvest than by rotation length. Whole-tree harvesting in both 16- and 32-year rotations results in high depletion rates, whereas stem harvest, regardless of rotation length, is approximately balanced by the N inputs from fixation and the atmosphere. In the intensive harvest system, annual nutrient drain is reduced by increasing rotation length. Nitrogen losses attributed to other management practices may be larger than those from harvest and are discussed in other sections of this report.

Most upland Piedmont soils are low in N (table 4). If all site N were completely available and productivity could be maintained as the

Table 4.--Changes in site nitrogen (N) with harvest intensity and rotation length of loblolly pine

Pool or process	Harvest age			
	16 years		32 years	
	Whole-tree harvest	Stem harvest	Whole-tree harvest	Stem harvest
	----- Pounds/acre -----			
Site reserves (immediately before harvest)				
Mineral soil, 0-28 inches	1,564	1,564	1,269	1,269
Forest floor	<u>274</u>	<u>274</u>	<u>321</u>	<u>321</u>
Total reserves	1,838	1,838	1,590	1,590
Harvest demand	230	103	382	208
Inputs (during rotation)				
N fixation	32	32	46	46
Atmospheric	<u>77</u>	<u>77</u>	<u>154</u>	<u>154</u>
Total input	109	109	200	200
Change due to harvest				
Per rotation	-121	+6	-182	-8
Per year (avg.)	-8	+1	-6	-1

quantities of soil N declined, soil N would be exhausted after 200 to 300 years of whole-tree harvesting. However, only a small proportion of mineral soil N becomes available for plant use annually. If the percentage of total N made available is constant, then available N will gradually decrease until a balance between growth demands, inputs, and available site N is reached. Depletion and the accompanying decline in tree growth may be prevented by applying fertilizer, by adjusting rotation length, or by limiting the removal of nutrient-rich biomass in harvests.

Although the nutrient drain by loblolly pine over a rotation appears large, the annual removal, compared with that of agricultural crops, is relatively small (table 5). Average annual production of pulpwood over a 16-year span is 3.2 tons of biomass/acre (7.2 t/ha). Averages of 5.8 lb (6.5 kg) of N, 0.8 lb (0.9 kg) of P, 4.5 lb (5.0 kg) of K, and 5.7 lb (6.4 kg) of Ca/acre (ha) are removed annually. Harvesting the entire tree, including roots, doubles or triples annual removal, but only

Table 5.--Comparison of the average annual yield and nutrient removal by a 16-year-old loblolly pine plantation with that of agricultural crops

Crop	Yield	N	P	K	Ca
	<u>Tons/ acre</u>	- - -	<u>Pounds/acre</u>	- - -	
Loblolly pine, whole tree	5.2	15.7	2.1	11.2	11.4
Loblolly pine, pulpwood	3.2	5.8	0.8	4.5	5.7
Corn (grain)	4.2	116	26.4	33.2	--
Soybeans (beans)	1.1	129	13.2	41.5	--
Alfalfa (forage)	3.6	189	20.7	165.8	67

about 60 percent more biomass (mostly leaves, branches, and roots of low commercial value) is produced.

Even with as severe a system as complete-tree harvest, nutrient removals are low compared with those of annual agronomic crops. Corn, soybeans, and alfalfa can deplete the soil at 10 times the rate of loblolly pine harvest. The major reason for the difference in nutrient removal of annual and tree crops is the type of biomass harvested. In agronomic crops, harvests are primarily seeds or leafy materials high in nutrients. Soybeans, for example, contain about 5 percent N. Loblolly pine wood has an N concentration of less than 0.1 percent, and needles, a reservoir of high nutrient concentrations, contain only about 1 percent. Yields of agronomic crops are maintained by fertilizing and good management practices. If nutrients to supplement those in the soil are not supplied, crop yields fall until availability, input, and output are in balance. Soil nutrient reserves and their availability control the rate of decline in yield of unfertilized agronomic crops. On poor sandy soils, the decline can be precipitous. On fertile prairie soils, where yields are often controlled by factors other than nutrients, decades may pass before yields fall below an acceptable level. Yield decline of agronomic crops may be analogous to what can happen in loblolly pine. On poor soils, short rotations and complete harvest can, over a few rotations, lead to unacceptably low production. On deep, fertile alluvial soils, many short biomass harvest rotations may pass before there are effects of nutrient depletion on productivity. Therefore, the harvest system used on any particular site should be one that, in the long term, will minimize the loss of nutrient capital.

## Influence of Harvesting on the Cycling of Residual Nutrients

In addition to the direct removal of nutrients in biomass, harvesting also influences cycling of the remaining site nutrients through chemical and physical site changes. Compared with preharvest conditions, the postharvest site has reduced evapotranspiration and the forest floor receives more solar radiation and precipitation. There is also a greater pool of nutrients from the harvest residues available for transfer to other components of the ecosystem than in the unharvested stand. Secondary effects of the harvest may include mixing organic residues with the mineral soil, transfer of nutrient-containing components from one area of the site to another, exposure of soil surfaces to erosion, and degradation of soil physical properties.

All harvesting, regardless of the intensity, will have some influence on cycling. After harvest, higher site temperature and moisture and greater availability of tree residues stimulate microbial activity. Organic residues decompose more rapidly, releasing their nutrients. The activities of nondecomposers, such as the microorganisms associated with N mineralization, are also stimulated. In one area of the North Carolina Piedmont, harvesting all trees increased N mineralization and nitrate production threefold and 36-fold, respectively, over an unharvested stand. Increased N losses due to leaching and denitrification may accompany increased nitrate production, whereas organic and ammonia N is subject to little direct loss. If the stand is thinned instead of clearcut, N mineralization will be greater than in uncut stands but still less than in clearcuts. In the thinned stand, the

greater portion of the mineralized N is likely to be taken up by the residual trees, rather than lost to leaching and denitrification. N mineralization and nitrate production may be less after complete-tree harvests than after stem harvests. The reason is that complete-tree harvests remove from the site the needles and other high N-content tree parts in addition to the low nutrient woody material.

## Fire

Two common types of prescribed fires used in southern forests--underburns and site-preparation burns--strongly influence nutrient cycling. Underburns are used in existing stands for fuel reduction, brush removal, range improvement, etc. These burns, which are relatively cool, remove only the most recently fallen litter and kill only the smaller hardwood stems. Site-preparation burns, which are relatively hot, are used for slash and forest floor reduction, and competition control in regeneration areas. The differences in the intensities of the burns and the amounts of materials consumed are major factors that produce the varying effects on nutrients and their cycling.

Burning can lead to nutrient losses by volatilization to the atmosphere, by increasing erosion, by ash removal from the site in air currents, and by leaching of elements that have been converted to more available or soluble form.

The most obvious effect of fire is a reduction in the forest floor. In the Southeastern Coastal Plain, 20 years of annual summer burning reduced the forest floor to 3.5 tons/acre (7.8 t/ha) compared with 13.4 tons/acre (30 t/ha) on unburned plots. Burning in the winter, after allowing a litter buildup for 4 or 5 years, consumed 3.2 tons (7.3 t) or 27 percent of the 12

tons/acre (26.9 t/ha) of litter present. The winter burn volatilized 100 lb (112 kg) of the 300 lb/acre (336 kg/ha) of N in the litter. Practically all of the organic N volatilized was converted to molecular N ( $N_2$ ), which cannot be used directly by plants. Burning may convert some organic N to ammonia, which can be adsorbed by the soil in its gaseous state. Absorption of ammonia on charred residues may help account for increases in soil N under some burning conditions. Low-intensity fires used in other studies have produced similar results. As fire intensity increases, a larger proportion of the forest floor will be consumed and the amount of N volatilized will be increased. Sulfur may also be lost by volatilization, but volatilization temperatures of other elements are high and their loss in this manner should be small.

Although much N can be lost in burning, burning may influence compensating processes and also influence N availability. Biological fixation by free-living micro-organisms can increase after fires, but fixation by these organisms probably amounts to only a few pounds per acre. It is severely limited by a lack of organic compounds to supply energy and by poor overall environmental conditions. A larger potential source of N may be fixation by higher plants. Following fire, N-fixing plants, both legumes and nonlegumes, are often prominent in the understory growth flush. Fixation by plants can range from a few to a hundred or more pounds per year, with the amount dependent on the plant density and the growth environment.

The nonvolatile components of the burned organic layers may either remain on the site, filtering into the organic and mineral soil layers, or be moved from the site by the wind or the convection currents set up by the fire. In one prescribed fire in South Carolina (Kodama and Van Lear 1980), nutrients lost from the litter (L)

layer (nearly 60 percent) were approximately proportional to the fraction of the L layer consumed by fire (table 6). Only small nutrient increases were measured in the F and H organic layers, and it was concluded that most of the nutrients had been lost from the site as ash in air currents and smoke created by burning.

Recurrent burning over long periods can transfer nutrients from the forest floor to the mineral soil. A severe treatment, annual burning for 20 years in the summer, reduced nutrient contents of the forest floor by 12, 18, 89, and 19 lb/acre (13, 20, 100, and 21 kg/ha) for P, K, Ca, and Mg, respectively. Roughly corresponding increases in Ca and Mg were found in the mineral soil, but no increases in P or K were recorded.

Table 6.--Quantities of nutrients in forest floor layers of four loblolly pine plantations before burning, and percentage change immediately after prescribed fire

Nutrient and condition	L layer		F + H layer	
	lb/acre	% change	lb/acre	% change
Calcium				
Preburn	33.9	--	80.3	--
Postburn	15.2	-55	88.4	+10
Magnesium				
Preburn	13.4	--	23.2	--
Postburn	4.5	-66	24.0	+3
Potassium				
Preburn	16.1	--	37.5	--
Postburn	6.3	-61	36.6	-2
Nitrogen				
Preburn	50.8	--	262.5	--
Postburn	27.7	-45	245.5	-6
Phosphorus				
Preburn	6.3	--	17.0	--
Postburn	2.7	-57	18.7	+10

Source: Adapted from Kodama and Van Lear 1980.

These quantities, both lost and transferred to the mineral soil, had a very small effect on total site nutrients, especially on an annual basis. Thus, burning to rapidly release nutrients will not alleviate site nutrient deficiencies caused by infertile soils. Good prescribed burning for hazard reduction and vegetation control will have little influence on nutrient loss or transfer.

In addition to fire intensity and amount of organic material consumed, soil properties often determine the fate and effects of released nutrients. Released nutrients may greatly affect pH and nutrient availability in soils with low exchange capacity such as sands, but their influence will be limited in soils with high exchange capacity, such as clays. Similarly, due to these properties and the differential potential for leaching losses, plant response to burning may be dramatic and short-lived on sand but moderate and extended on clay.

Properly applied hazard-reduction burns that remove only a portion of the L layer and none of the F and H, and do not expose mineral soil, do not increase runoff. Nutrient loss via runoff or erosion, therefore, is not appreciably increased. Soil erosion can be the most obvious consequence of burning. However, in only one of six burning studies by Ralston and Hatchell (1971) (one in the North Carolina Piedmont) did erosion greatly exceed a soil loss of 1 inch (3 cm) in 1,000 years or an annual loss of 0.3 tons/acre (0.67 t/ha). This rate is the soil-loss rate estimated to take place as a normal geologic erosion process.

Burning may indirectly affect nutrient cycling by altering the micro and macro flora and fauna of the forest floor. In long-term prescribed burning programs, the populations of spore-forming microfungi or bacteria are not reduced sufficiently to impair soil processes. Intensive fires, however, may reduce populations

dramatically, temporarily sterilizing the soil. This is most likely to occur in fires where slash has been piled and burned for disposal.

Nitrogen fixation by free-living soil micro-organisms has been found to increase after burning. The amounts fixed range from about 1 lb/acre/year to as high as 12.5 lb/year in some areas. Comparable unburned areas fix less than 0.05 lb annually. Fixation rates by free-living organisms are limited by a lack of an energy source and available P in most forest soils.

Individual low-intensity prescribed fires do not cause important losses of nonvolatile nutrients. However, the long-term cumulative losses of the volatile nutrients, especially N, can be substantial--over 100 pounds per acre during a rotation. Nevertheless, this quantity is small in relation to that removed in a complete harvest of aboveground biomass plus the losses that accompany intensive site preparation. In intense burns, as in uncontrolled wildfires and those used for slash reduction, important decreases of site N and sulfur through volatilization can occur. Losses of nonvolatile elements caused by intense fires are usually associated with ash carried from the site rather than by postfire leaching. Losses of these nonvolatile elements can be large, but even they are smaller than can occur in harvest and site-preparation activities.

### Site Preparation

Site preparation controls competition, removes impediments to planting, and modifies the site nutritionally and physically for improved tree growth. If properly handled, competition control and the removal of planting impediments will not adversely affect the nutrient cycle. However, movement of organic matter and soil to produce an immediate gain in seedling growth and survival may have negative effects on nutrient availability later in the rotation.

One of the most intensive systems for site preparation is shearing, rootraking, and windrowing followed by disking for final competition control. This system tends to concentrate surface soil, roots, forest floor, and logging debris in windrows and to mix any organic residues between the windrows with the mineral soil. On infertile soils, such nutrient displacement can seriously affect the growth of the present rotations as well as of successive ones. The result is superior growth of trees adjacent to windrows compared with those farther from the windrows. In a flatwood site in Florida, Morris and others (1983) found that although windrows occupied only 6 percent of the site, these areas contained from 10 to 40 percent of the total or extractable nutrient reserves in the system. These nutrient displacements were larger than were the removals associated with harvesting the stand. N displacement into windrows was six times the 5.3 lb N/acre (5.9 kg/ha) removed in the bolewood harvest. Displacements of P, K, Ca, and Mg into the windrows also equaled or exceeded those in the harvest removal.

Erosion following soil disturbance transports nutrients in both dissolved and solid forms. Losses from erosion may be greater than those estimated from average nutrient concentrations in sediments due to the selective transportation of organic and mineral-organic particles. Only on sloping sites and those without protective cover, however, should overall nutrient reserves be significantly affected by erosion. Local areas affected by improperly constructed or located roads and skid trails may be severely impacted. Leaching losses of all elements except N are usually small.

Site preparation can significantly affect biological activity and associated nutrient availability. Removing the protective plant canopy and forest floor by shearing, piling, and disking can raise soil temperatures 3.6 to 9 °F (2 to 5 °C)

above those that occur with low-intensity treatments such as chopping. Fewer plants on prepared sites also reduce transpirational losses and result in higher soil moisture. The warmer and wetter conditions coupled with the mixing of forest floor and soil promote greater biological activity. Increased activity is especially important in the conversion of organic N to nitrate N, which is subject to leaching and denitrification. In late summer, the nitrate pool in the surface soil of a sheared, piled, and disked area on the Piedmont of North Carolina was 10 lb/acre (11 kg/ha) compared with less than 1 lb for a chopped-only treatment. Differences of similar magnitude were also found in soil solutions at the 28-inch (70 cm) depth. Use of herbicides and the elimination of vegetation greatly increased nitrate in both surface soil and at the 28-inch depth, adding to the loss of site N.



Nutrient losses from site preparation, especially those of N, may be reduced by maintaining an intact forest floor and by allowing the planted site to revegetate normally, controlling only the most troublesome competing vegetation. These soil covers will moderate the soil temperature and moisture increases associated with intensive site preparation that promote high biological activity. The forest floor and organic residues can also immobilize mineralized N, and the small plants use any excess mineralized N. Nitrogen retained in these pools may be utilized for tree growth later in the rotation.

The mineralization rate of organic N, as influenced by site-preparation intensity, has important implications for tree growth. Models relating N release to the intensity of site preparation and N growth requirements are shown in figure 9. After both high and low intensities of site preparation, there is an initial excess of mineralized N--more than the amount taken up by the small root systems of the young trees. Under




Figure 9.--Theoretical effect of site-preparation intensity on long-term N reserves. High-intensity site preparation (above) includes competition control and mixing of the forest floor and mineral soil. Low-intensity treatment (below) includes only competition control. (Adapted from Burger and Kluender 1982.)

high-intensity preparation, the N surplus is larger and is lost more rapidly than under low-intensity preparation. Late in the rotation, this loss of N reserves causes poor tree growth. Even later, N reserves may be lacking for growth in subsequent rotations. Thus, intensive preparation that changes the rate and cycling pathways of nutrients may provide for rapid early growth but can result in reduced growth over the long term.

### Fertilization

Forest fertilization has become a standard practice for many forest products companies that demand high productivity from loblolly pine sites. Fertilization can modify nutrient cycling through changes in the size of components of the cycle (the pools), and in the transfer rates between pools. For example, fertilization with a soluble nutrient source may influence not only the leaching rate of the applied element but also those of other elements competing for exchange sites in the soil. Additions of nutrients in short supply may also increase the uptake of other nutrients as tree growth accelerates. These two simple examples, among many, can substantially change nutrient cycling on a site, especially if fertilization is accompanied by modifications in site preparation, rotation length, harvesting procedures, and other forest management practices.

In addition to alleviating naturally occurring nutrient deficiencies, a second role for fertilizer is to replace nutrients removed via harvest or as a consequence of site disturbance. Unfortunately, using fertilizers to replace nutrients is more complex than simply substituting for those removed in a one-to-one ratio. Information on nutrient uptake by loblolly pine over long periods is limited, but what exists shows overall efficiency of fertilizer use is low. For applied N, about one-quarter is taken up by trees, one-quarter is held on the site in other components,

and the remainder is lost by leaching and volatilization. Thus, for N, if inputs are needed to balance losses, between 2 and 4 pounds of fertilizer N would have to be applied for every pound removed in management activities.

Phosphorus utilization during the first rotation is similar to that of N. However, unlike N, little P is lost from the site. Little is known about the applied P availability during succeeding rotations. No information is available on the uptake-application ratios of other nutrients in loblolly pine stands.

The point at which nutrient replenishment is required for yield maintenance is not known. When replenishment of most nutrients is attempted, small quantities of nutrients applied several times during a rotation may be better than one large application at the time trees are established. Perhaps even greater knowledge and effort will be required for efficient nutrient maintenance than for fertilization to alleviate obvious nutrient deficiencies.

### Soil-Improving Plants

Wild legumes and other N-fixing plants help maintain the ecosystem's supply of N, and as understory contribute to the cycling of other nutrients from the lower into the upper soil layers and the forest floor. At certain times during the loblolly pine rotation, it may be appropriate to establish soil-improving plants. Ideally, these plants should not have severe competitive or allelopathic effects on the pines, should fix and release important quantities of N, and should be tolerant of the severe site conditions that are often found in loblolly plantations.

In one young loblolly plantation on the North Carolina Coastal Plain, trees were fertilized with

P and grown in stands of lespedeza (Lespedeza cuneata (Dumont.) G. Don and L. thunbergii (DC Naki) established to provide biologically fixed N. Trees with the P plus lespedeza, at age 7, had a volume of 10 cords/acre (89.6 m<sup>3</sup>/ha) compared with 8.4 cords (75.2 m<sup>3</sup>) for trees with P fertilizer alone. During the initial years of lespedeza development, foliage of trees in the lespedeza stands contained higher concentrations of N than did controls or trees receiving only P. As lespedeza biomass declined with stand closure, foliar N concentrations, regardless of treatment, became similar. The lespedeza was estimated to have added 400 lb of N/acre to the site over its 4-year period of maximum growth.

There may be instances when the secondary effect of legumes may be as important as that of the N fixed. Deep-rooted legumes may reduce leaching losses by taking up nutrients from below the rooting zones of young trees. Leguminous litter, high in N, may speed up forest floor decomposition, hastening nutrient release and cycling. Legumes may also be capable of improving the physical properties of soil, thereby increasing the potential soil volume from which trees may extract moisture and nutrients.

### Intensive Forest Management and Productivity

Barring severe erosion or soil and nutrient displacement, it is unlikely that even the most intensive harvesting and silvicultural procedures will transform productive loblolly pine sites into unproductive ones in a rotation or two. Existing management systems, however, are based upon sustained yield with the implication that productivity will be maintained. Yet, there is a dearth of long-term data showing whether present management is meeting sustained-yield objectives.

Despite many limitations and unknowns, nutrient budgets may provide a method for comparing effects of alternative forestry practices on ecosystem nutrients (table 7). Although only a budget for N is presented here, budgets can be prepared for other nutrients and the implications similarly studied. Some nutrient losses during harvest and regeneration are unavoidable, but the quantities removed depend largely on the management techniques adopted and the skill with which they are carried out. For example, employing three short rotations rather than a single 60-year rotation increases stem yield from 87 to 144 tons/acre (195 to 323 t/ha) with only proportionate increases in nutrient removal (Switzer and others 1978). If, however, the entire aboveground portion of the loblolly stand is harvested, N removal increases from 155 to 634 lb/acre (174 to 711 kg/ha) and P from 12 to 59 lb/acre (13 to 66 kg/ha) or more than fourfold during the 60-year

Table 7.--Effects of stem and whole-tree harvest of loblolly pine and site-preparation practices on nitrogen demands at rotations of 16 and 32 years, in pounds per acre

Practice or process	16 years		32 years	
	Whole tree <sup>a</sup>	Stem	Whole tree <sup>a</sup>	Stem
DEMANDS OF N				
Harvest	230	103	382	208
KG, windrow, disk	209	336	209	336
Erosion				
Kg, w, disk (Piedmont)	28	28	28	28
Chop (Piedmont)	5	5	5	5
Prescribed burn	60	60	120	120
Slash burn	0	120	0	136
Leaching				
Disk or bed	32	32	40	40
Chop or herbicide	16	16	20	20
INPUT OF N				
Atmospheric deposition	77	77	154	154
Nitrogen fixation				
No seeding	32	32	46	46
Seeded legumes	400	400	400	400

<sup>a</sup>Excluding roots.

period. When nutrient changes of this magnitude can occur as a result of a management decision, the long-term ability of the ecosystem to support this yield must be examined. If the conclusion is negative, there are three basic remedies: (1) apply fertilizers, (2) change the rotation length, and (3) harvest less biomass. If one or more of these remedies are not adopted, yield reductions in future rotations must be anticipated.

Other causes of nutrient outflow from a site may be as great or greater than that of the harvest. Intensive site-preparation practices such as rostraking and windrowing are one such cause; prescribed burning is a second. Surprisingly, erosion, although conspicuous, seldom causes major nutrient losses, and these losses can be minimized by converting from intensive to conservative site preparation. Similarly, losses from leaching can be reduced by using conservative site-preparation treatments.

By examining various forest management systems and their associated nutrient removals, some idea can be obtained of their overall impact on the ecosystem and its ability to provide nutrients for sustained yield. Several systems with widely differing N removal rates are shown in table 8. Included are both intensive and conservative management systems. Even the most conservative system listed, the 16-year stem-only harvest followed by chopping, causes a small net loss of N without legumes. With a 32-year rotation and chopping, annual loss would be similar, 1 lb/acre. These data indicate that when stem harvest is the largest contributor to N removal, rotation age does not influence nutrient loss on an annual basis. Nutrient losses associated with the stem are uniform because stem biomass of the stand increases fairly uniformly to maturity. If the entire tree is harvested, nutrient losses of the stand allocated over the lengthened rotation are reduced due to stand foliage and branch biomass peaking at an early age.

Table 8.--Changes in system nitrogen with various combinations of rotation length, biomass removal, and site preparation in loblolly pine plantations

Rotation length (years)	Biomass removal	Site treatment	Rotation	Annual
			Pounds/acre	
16	Whole tree	1	-450	-29
16	Whole tree	2	-82	-5
32	Whole tree	1	-579	-18
32	Whole tree	2	-225	-7
16	Stem	3	-15	-1
16	Stem	4	+233	+15
32	Whole tree	5	+27	+1
32	Stem	6	-148	-5

Treatments:

1. KG-windrow, disk, prescribed burn (PB).
2. Treatment 1 plus legumes.
3. Chop.
4. Chop, slash burn, legumes.
5. Chop, PB, legumes.
6. Herbicide, PB, natural regeneration.

Site-preparation practices rank first or second as causes of nutrient drain. Roottraking and disking are sometimes combined with bedding to impose severe nutrient demands on the site. Although good survival and early growth can be obtained with these methods, less nutrient-costly alternatives may give equally good yields over the long term. Chopping and herbicide applications may substitute for intensive procedures, especially if site-to-site adjustments in planting and preparation can be made.

Fertilization can offset nutrient losses, but for N its effect is short term and the proportion of the applied N recovered by trees is often low.

Legumes may be an alternative source of N. However, their utilization is more complex than the application of fertilizer. Seedbed preparation, fertilization, selection of species, and competition are considerations if biologically fixed N is to supplement or supplant N fertilization.

Quantities of nutrients added as fertilizer to compensate for those removed from the site by harvest or lost during regeneration are not of equal value in maintaining site productivity. For example, the N lost in erosion, windrowing, burning, and harvesting is derived mostly from organic compounds that are normally mineralized and made available to higher plants over many years. Organic matter and organic N are essential to maintain the physical, chemical, and biological properties of the soil; therefore, fertilization may be a poor and economically unsound substitute for N losses, especially on infertile sites already low in organic matter.

Based upon the known deficiencies and probability of losses from the loblolly pine ecosystem, the impacts of intensive management practices on site nutrients are in the order of  $N > P > K > Ca >$  other elements. Like N demands, those for P and K under intensive forest management exceed natural inputs. Phosphorus deficiency is less difficult to correct than N because P persists in the soil and trees are able to utilize it in slowly soluble forms. Rock phosphate is a cost-effective, slowly soluble source of P and, when available, may be substituted for triple superphosphate and other soluble forms. When large amounts of biomass are removed and intensive site preparation causes accelerated leaching, K and Ca deficiencies will eventually limit growth on many soils. Potassium can persist because it can be taken up in luxury quantities and cycled by the vegetation with few leaching losses. Calcium, however, is cycled more slowly and when accompanied by N fertilization or high mineralization

of native N is readily leached with the nitrate ion. Its loss could reduce the productivity of base-deficient sandy soils.

As a solution to the problem of excess nutrient removal over that of input, it has been suggested that an "ecological rotation" be matched to site. In this system, rotation length and harvest are related to nutrient input from the atmosphere and from soil mineral weathering. While ecologically sound, this approach ignores the general lack of primary nutrient-containing minerals in soils of the South and the low and variable atmospheric input rates of the different nutrients. Furthermore, most rotation lengths would have to be increased to such an extent that annual production would fall. Forest managers would also lose flexibility of forest production. The best system for minimizing the potential for a long-term decline in yield is a forest manager with a knowledge of and the ability and freedom to apply good nutrient cycling principles.



## Literature Cited

- Burger, J.A.; Kluender, R.A. Site preparation-Piedmont: In: Symposium, loblolly pine ecosystem. 1982 December 8-10; Raleigh, NC. 1982:58-74. [Available from North Carolina State University, Raleigh]
- Kodama, A.E.; Van Lear, D.H. Prescribed burning and nutrient cycling relationships in young loblolly pine plantations. *Southern Journal of Applied Forestry* 4:118-121; 1980.
- Morris, L.A.; Pritchett, W.L.; Swindel, F.G. Displacement of nutrients in windrows during site preparation of a flatwood forest. *Soil Science Society of America Journal* 47:591-594; 1983.
- Ralston, C.W.; Hatchell, G.E. Effects of prescribed burn on physical properties of soil. In: Prescribed burning symposium proceedings. 1971 April 14-16; Charleston, SC. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station; 1971:68-85.
- Switzer, G.L.; Nelson, L.E. Nutrient accumulation and cycling in loblolly pine (*Pinus taeda* L.) plantation ecosystems: the first twenty years. *Soil Science Society of America Proceedings* 36:143-147; 1972.
- Switzer, G.L.; Nelson, L.E.; Hinesley, L.E. Effects of utilization on nutrient regimes and site productivity. In: Proceedings, Complete tree utilization of southern pine. 1978 April; New Orleans, LA. Forest Products Research Society; 1978:91-102.
- Switzer, G.L.; Nelson, L.E.; Smith, W.H. The mineral cycle in forest stands. In: Forest fertilization--theory and practice. TVA, Muscle Shoals, AL; 1968:1-9.

Jorgensen, Jacques R.; Wells, Carol G.

Foresters' primer in nutrient cycling. A loblolly pine management guide, Gen. Tech. Rep. SE-37. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station; 1986. 42 pp.

The nutrient cycle, which includes the input of nutrients to the site, their losses, and their movement from one soil or vegetation component to another, can be modified by site preparation, rotation length, harvest system, fertilization, and fire, and by using soil-improving plants. Included is a report on how alternative procedures affect site nutrients, and provides general principles that can be followed to enhance long-term productivity of loblolly pine.

**Keywords:** Pinus taeda, nitrogen, phosphorus, potassium, rotation length, nitrogen fixing, whole-tree harvesting.

Jorgensen, Jacques R.; Wells, Carol G.

Foresters' primer in nutrient cycling. A loblolly pine management guide, Gen. Tech. Rep. SE-37. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station; 1986. 42 pp.

The nutrient cycle, which includes the input of nutrients to the site, their losses, and their movement from one soil or vegetation component to another, can be modified by site preparation, rotation length, harvest system, fertilization, and fire, and by using soil-improving plants. Included is a report on how alternative procedures affect site nutrients, and provides general principles that can be followed to enhance long-term productivity of loblolly pine.

**Keywords:** Pinus taeda, nitrogen, phosphorus, potassium, rotation length, nitrogen fixing, whole-tree harvesting.



The Forest Service, U.S. Department of Agriculture, is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives—as directed by Congress—to provide increasingly greater service to a growing Nation.

USDA policy does not permit discrimination because of race, color, national origin, sex or religion. Any person who believes he or she has been discriminated against in any USDA-related activity should write immediately to the Secretary of Agriculture, Washington, D.C. 20250.