

Reprinted from

# Forest Ecology and Management

---

Forest Ecology and Management 114 (1999) 321-327

Wildlife linkages: volumes and values of residual  
timber in riparian zones in eastern Texas

Gary Allen Burns<sup>a</sup>, R. Montague Whiting, Jr.<sup>b,\*</sup>,  
George M. LeGrande<sup>c</sup>, James G. Dickson<sup>d</sup>

<sup>a</sup> *Burns Forestry, P.O. Box 1227, Crockett, TX 75835, USA*

<sup>b</sup> *College of Forestry, Stephen F. Austin State University, Nacogdoches, TX 75962, USA*

<sup>c</sup> *College of Forestry, Stephen F. Austin State University, Nacogdoches, TX 75962, USA*

<sup>d</sup> *USDA Forest Service, P.O. Box 7600, SFA Station, Nacogdoches, TX 75962, USA*



ELSEVIER

# Forest Ecology and Management

**Aims and scope.** Forest Ecology *and* Management publishes scientific articles concerned with forest management and conservation, and in particular the application of biological, ecological and social knowledge to the management of man-made and natural forests. The scope of the journal includes all forest ecosystems of the world. A refereeing process ensures the quality and international interest of the manuscripts accepted for publication. The journal aims to encourage communication between scientists in disparate fields who share a common interest in ecology and natural-resource management, and to bridge the gap between research workers and forest managers in the field to the benefit of both. The journal should be of interest to research workers, managers and policy makers in forestry, natural resources, ecological conservation and related fields.

## FOUNDING EDITOR

Laurence L. Roche, Murroe, Ireland

## EDITORS-IN-CHIEF

For the Americas, Australia, New Zealand and the Pacific:

**R.F. Fisher**

Department of Forest Science

Texas A&M University

College Station, TX 77843-2135, USA

For the rest of the world:

**G.M.J. Mohren**

Forest Production Ecology Group

Department of Vegetation Ecology

DLO-Institute for Forestry and Nature Research

P.O. Box 23

6700 AA Wageningen, The Netherlands

## BOOK REVIEWS EDITOR

**Margaret R. Gale**

School of Forestry and Wood Products

Michigan Technological University

1400 Townsend Drive

Houghton, MI 49931, USA

## EDITORIAL ADVISORY BOARD

**G. Abrahamsen**, Agricultural University of Norway, As, Norway

**R. Alfaro**, Canadian Forestry Service, Victoria, B.C., Canada

**F. Andersson**, Swedish University of Agricultural Sciences, Uppsala, Sweden

**P.M.S. Ashton**, Yale University, New Haven, USA

**P. Attiwill**, University of Melbourne, Parkville, Vic., Australia

**S. Boyle**, Oregon State University, Corvallis, USA

**J. Brown**, US Environmental Protection Agency, Corvallis, USA

**I. Burke**, Colorado State University, Fort Collins, CO, USA

**J.C. Calvo**, Institute of Technology, Cartago, Costa Rica

**R.-J. Chen**, University of Hong Kong, Pokfulan Road, Hong Kong

**J.D. Deans**, Institute of Terrestrial Ecology, Penicuik, Midlothian, UK

**R.M. DeGraaf**, USDA Forest Service, University of Massachusetts, Amherst, MA, USA

**S. Diamandis**, Forest Research Institute, Thessaloniki, Greece

**D.P. Dykstra**, CIFOR, Jakarta, Indonesia

**E.P. Farrell**, University College Dublin, Dublin, Ireland

**P.M. Fearnside**, Instituto Nacional de Pesquisas da Amazonia-INPA, Manaus-Amazonas, Brazil

**PH. Freer-Smith**, Forestry Commission, Farnham, UK

**O. Garcia**, ENGREF, Nancy, France

**D. Gilmore**, Canadian Forest Product Ltd., Alberta, Canada

**R.A. Goyer**, Louisiana State University, Baton Rouge, LA, USA

**J.B. Hall**, University College of North Wales, Bangor, UK

**F. Houllier**, Campus International de Baillarguet, Laboratoire

Associe de Modelisation des Plantes (AMAP), Montpellier, France

**B.M. Kumar**, Kerala Agricultural University, Kerala, India

**J.P. Lassoie**, Cornell University, Ithaca, NY, USA

**J.N. Long**, Utah State University, Logan, UT, USA

**A.E. Lugo**, International Institute of Tropical Forestry, Rio Piedras, PR, USA

**J.A. Maghembe**, SADCC / ICRAF Agroforestry Project, Zomba, Malawi

**F. Makeschin**, Institut für Bodenkunde und Standortslehre, Tharandt, Germany

**DC. Malcolm**, University of Edinburgh, Edinburgh, UK

**E. Mälkönen**, Finnish Forest Research Institute, Vantaa, Finland

**M.A.R. Nahuz**, Instituto de Pesquisas Tecnológicas, São Paulo, SP, Brazil

**R. Päivinen**, European Forestry Institute, Joensuu, Finland

**S.G. Pallardy**, University of Missouri, Columbia, MO, USA

**R.F. Powers**, Pacific Southwest Research Station, Redding, CA, USA

**T. Pukkala**, University of Joensuu, Joensuu, Finland

**F.E. Putz**, University of Florida, Gainesville, FL, USA

**L. Rasmussen**, RiSO, Roskilde, Denmark

**D.D. Reed**, Michigan Technological University, Houghton, MI, USA

**R. Sands**, University of Canterbury, Christchurch, NZ

**J.A. Stanturf**, Stoneville, MS, USA

**Q. Sziklai**, University of British Columbia, Vancouver, B.C., Canada

**J.R. Toliver**, USDA Forest Service, Washington, DC, USA

**K. von Weissenberg**, University of Helsinki, Helsinki, Finland

**D. Whitehead**, Manaaki Whenua Landacre Research, Lincoln, New Zealand

**Publication information:** Forest Ecology *and* Management (ISSN 0378-1127). For 1999 volumes 111-122 are scheduled for publication. Subscription prices are available upon request from the Publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where air delivery via SAL mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available on request. Claims for missing issues should be made within six months of our publication (mailing) date.

**Orders, claims, and product enquiries:** please contact the Customer Support Department at the Regional Sales Office nearest you:

**New York:** Elsevier Science, PO Box 945, New York, NY 10159-0945, USA; Tel. (+1) (212) 633 3730, [toll free number for North American customers: 1-888-4ES-INFO (437-4636)]; fax: (+1) (212) 633 3680; e-mail usinfo-f@elsevier.com

**Amsterdam:** Elsevier Science, PO Box 211, 1000 AE Amsterdam, The Netherlands; Tel.: (+31) 20 485-3757; fax: (+31) 20 485 3432; e-mail nlinfo-f@elsevier.nl

**Tokyo:** Elsevier Science, 9-15, Higashi-Azabu 1-chome, Minato-ku, Tokyo 106, Japan; Tel. (+81) (3) 5561 5033; fax (+81) (3) 5561 5047; e-mail: info@elsevier.co.jp

**Singapore:** Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192; Tel. (+65) 434 3727; fax: (+65) 337 2230; e-mail: asiainfo@elsevier.com.sg

**Rio de Janeiro:** Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; phone: (+55) (21) 509 5340; fax: (+55) (21) 507 1991; e-mail: elsevier@campus.com.br [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]

## Wildlife linkages: volumes and values of residual timber in riparian zones in eastern Texas

Gary Allen Burns<sup>a</sup>, R. Montague Whiting, Jr.<sup>b,\*</sup>,  
George M. LeGrande<sup>c</sup>, James G. Dickson<sup>d</sup>

<sup>a</sup> *Bums Forestry, 170. Box 1227, Crockett, TX 75835, USA*

<sup>b</sup> *College of Forestry, Stephen F. Austin State University, Nacogdoches, TX 75962, USA*

<sup>c</sup> *College of Forestry, Stephen F. Austin State University, Nacogdoches, TX 75962, USA*

<sup>d</sup> *USDA Forest Service, P.O. Box 7600, SFA Station, Nacogdoches, TX 75962, USA*

---

### Abstract

In regenerating southern pine, maintenance of riparian zones (RZs) is a major land concession for soil and water protection and wildlife habitat enhancement. However, there are few data quantifying the volume and value of residual timber in such areas. We inventoried merchantable timber in nine RZs of three widths in sapling-class East Texas pine plantations. Present, discounted, and projected volumes and values of residual timber were determined. Average per-acre volumes of narrow, medium, and wide RZs were 337, 1438, and 2542 board feet (Doyle log rule) and 4.6, 8.2, and 7.2 cords, respectively. At US\$ 154.00 and US\$ 57.00 per thousand board feet for pine and hardwood saw timber, respectively, and US\$ 15.00 and US\$ 5.00 per cord for pine and hardwood pulpwood, respectively, average present per-acre values were US\$ 46.41, US\$ 209.93 and US\$ 352.75 for narrow, medium, and wide RZs. Interest rates for the growth of trees from 1981 to 1989 ranged 3.7%–23.5% for pine and 2.2%–6.1% for hardwood. For the 1981-1989 discount period, average per-acre net annual equivalents of narrow, medium, and wide RZs were US\$ 4.52, US\$ 20.46, and US\$ 34.38, respectively. Stumpage values at the time of harvest projected at 7% compound interest for 30 years ranged US\$ 10.66–US\$ 3547.54 per acre and for 80 years ranged US\$ 313.93–US\$ 104499.95 per acre. The impact of these results on wildlife is discussed. © 1999 Elsevier Science B.V. All rights reserved.

---

### 1. Introduction

Increased demand for wood products in the southern United States has resulted in many natural pine-hardwood stands being converted to even-age pine plantations. Strips of mature trees, however, are usually retained along streams when adjacent upland stands are harvested. Such areas are referred to as streamside management zones, stringers, streamers

and/or riparian zones (RZs). Riparian zones can buffer streams from erosion and non-point water pollution. They offer protection to stream banks and beds and act as sediment filters. Wildlife habitat is also enhanced by these corridors and by the increase in habitat diversity (Dickson, 1989).

In most RZs, the moisture regime is favorable for rapid tree growth, the site index is high, and substantial timber resources may be present. However, there are few quantitative data on timber volumes and values or timber-wildlife relationships in RZs. This

---

\*Corresponding author. Tel.: +1-409-4683301.

study focused on volumes and values of residual timber in **RZs** of three widths along first- and second-order intermittent streams in eastern Texas.

## 2. Methods

### 2.1. Study areas

Personnel with the United States Forest Service (USFS), Southern Forest Experiment Station, Nacogdoches, TX, selected three sites in eastern Texas to evaluate the impacts of **RZs** on wildlife (Dickson and Bosch, 1990). Within each site, Forest Service biologists established a narrow (<85 ft), a medium (98-131 ft), and a wide (>164 ft) **RZ** study area, thus, a total of nine study areas. All study areas were in **RZs** that bisected young pine plantations. Within each study **area**, two sample transects, each 656 ft long, were created. Each transect was divided into four equal-length compartments and in all but one study area, a short buffer existed between transects. Buffers ranged 57-209 ft in length (Dickson and Huntley, 1987). For the purpose of this study, the transects were as wide as the **RZs**.

The **RZs** were on property owned by forest industry. When the study areas were established in 1984, all pine plantations were less than 5 years old. The pine plantations resulted when second growth pine-hardwood stands were **clearcut** and the areas mechanically site-prepared and planted to 1-O loblolly pine (*Pinus taeda*) seedlings. Some timber was selectively harvested within the **RZs** and usually more trees per acre were removed from narrow **RZs** than from wide **RZs**. Soils of all the nine study areas were broadly defined as sandy loams.

### 2.2. Field procedures

In each study area, a 100% inventory of all merchantable timber in the transects and in the between-transect buffers was conducted; species, dbh, merchantable height, and product class were recorded for each merchantable tree. All trees were measured using USFS (1985) guidelines. Lengths of the transects and between-transect buffers had been previously measured (Dickson and Huntley, 1987); widths were measured by pacing from the edge of one plantation

perpendicularly across the **RZ** to the edge of the other. At least four such measurements were made along each transect.

To determine growth, age, and form class, prism points were established in the center of every other compartment. Then, rotating clockwise from the bearing of the transect, the first pine and hardwood included in the lo-factor variable radius plot were measured to determine form class and radial growth. Form class was estimated with Wiant form class wedges. Tree age and radial growth data were gathered from increment cores taken at breast height. For determining the age, 3 and 2 years were added to the age at dbh for pines and hardwoods, respectively (USFS, 1985). These data were grouped as either pine or hardwood.

### 2.3. Data analyses

Merchantable trees in each **RZ** were categorized as pulpwood or saw timber using USFS (1985) guidelines. Present, past, and projected merchantable volumes of timber in each **RZ** were calculated. Present volumes were those at the time of the inventory, past volumes were those 8 years previous to the inventory at approximately the time of the prior harvest, and projected volumes were those 8 years following the inventory.

Present volumes were estimated using dbh and height at the time of inventory; the Doyle log rule and the average form class for each species group and product class were also used. For discounted and projected volumes, diameters were estimated by subtracting or adding the average 8-year diameter growth of the appropriate species group to the current diameter of each merchantable tree. The resulting diameter was then paired with the average merchantable height for that diameter class to determine the volume of the tree. Neither **ingrowth** nor mortality were considered.

Timber values were estimated using 10-year average **stumpage** prices (1980-1989) as reported by the Texas Forest Service (1989). The values used were US\$ 5.00 per cord for hardwood pulpwood, US\$ 15.00 per cord for pine pulpwood, US\$ 57.00 per thousand board ft for hardwood saw timber, and US\$ 154.00 per thousand board ft for pine saw timber. Neither fluctuations in timber prices nor inflation were

considered in computing discounted and projected values.

Compound interest rates were computed for the discounted and projected per-acre residual timber values by species product class. The values represent the rate earned on the capital investment in the marketable timber. The formula used to calculate the compound interest rate ( $r$ ) was:

$$r = n\sqrt[n]{\frac{FV}{PV}} - 1$$

where

$n$  number of years in the period,  
 FV future value,  
 PV present value.

The net annual equivalents (NAE), the annual payment that will equal the net present value of an asset during its lifetime, were also determined. The NAE is useful in comparing an investment that yields a periodic return with one that yields an annual return and in comparing alternatives having unequal lives. The formula used, which is identical to the installment payment formula, was:

$$NAE = NPV \left( \frac{i(1+i)}{(1+i)^n - 1} \right)$$

where

NPV net present value,  
 $i$  investment interest rate available,  
 $n$  number of years in the period.

The interest rate ( $i$ ) used was 7% and the period ( $n$ ) was the average number of years elapsed between the

time the adjoining stand was harvested and the data were gathered for this study (i.e. 8 years). The NAEs were computed to include the value of the growth of the merchantable timber.

### 3. Results

Form classes of pine and hardwood saw timber averaged 81 and 77, respectively; pulpwood form classes averaged 82 for pine and 77 for hardwood. Diameter growth ranged from 0.13 to 0.40 in per year for pines and 0.10-0.33 in per year for hardwoods. Average annual diameter growth was 0.30 in for pine and 0.18 in for hardwood. Total tree ages ranged from 36 to 93 years for pine and 34-134 years for hardwood. Average ages for pine and hardwood were 59 and 63 years, respectively.

The narrow RZs had very few merchantable stems and thus little saw timber, especially pine saw timber (Table 1). Virtually all the pine had been harvested and most of the standing pulpwood volume was hardwood. Some saw timber had been selectively harvested from all medium RZs, but all had saw timber volume remaining, thus all had substantially more volume than did the narrow RZs. One wide RZ had no trees harvested and another had only a few trees removed. As a result, the wide RZs had almost twice as much saw timber volume as did the medium ones. All RZs had pulpwood volume and most had much more hardwood than pine pulpwood (Table 1).

As would be expected, the narrow RZs had the lowest average merchantable timber value per acre and the wide RZs had the highest. Present value of the merchantable timber in the narrow RZs averaged US\$ 46.42 per acre (range US\$ 2.25-US\$ 114.20).

Table 1

Average numbers of stems per acre and volume per acre of pine and hardwood saw timber and pulpwood trees in RZs of three widths in eastern Texas, 1989

RZ width	Saw timber						Pulpwood					
	Stems/acre			Vol./acre (bd. ft.)			Stems/acre			Vol./acre (cords)		
	Pine	Hardwood	Total	Pine	Hardwood	Total	Pine	Hardwood	Total	Pine	Hardwood	Total
Narrow	<1	8	8	9	327	336	8	44	52	0.4	4.2	4.6
Medium	6	9	15	747	691	1438	15	69	84	1.5	6.7	8.2
Wide	6	12	18	1648	894	2542	13	67	80	1.2	6.0	7.2

The Doyle log rule was used to determine saw timber volumes.

Table 2  
Estimated stumpage values (US\$/acre) of merchantable timber in RZs of three widths in eastern Texas

RZ width	Saw timber			Pulpwood			Total
	Pine	Hardwood	Subt.	Pine	Hardwood	Subt.	
<b>Present (1989)</b>							
Narrow	1.44	18.66	20.10	5.30	21.02	26.32	46.42
Medium	115.09	39.37	154.46	21.90	33.57	55.47	209.93
Wide	253.79	50.94	304.65	18.15	29.90	48.10	352.75
<b>Discounted (1981-1989)</b>							
Narrow	0.82	13.11	13.93	1.45	16.50	17.95	31.88
Medium	74.38	31.10	105.48	10.80	26.00	36.80	142.28
Wide	163.14	37.76	200.90	11.00	23.78	34.78	235.68
<b>Projected (1989-1997)</b>							
Narrow	2.41	31.10	33.51	11.15	22.22	33.37	66.88
Medium	200.56	57.23	257.79	23.00	38.60	61.60	319.39
Wide	352.15	65.87	417.98	24.40	36.47	60.87	478.85

The average merchantable timber value of the medium RZs was US\$ 209.93 per acre (range US\$ 98.77–US\$ 349.34) and that of the wide RZs was US\$ 352.75 per acre (range US\$ 92.72–US\$ 699.85) (Table 2). Trends in discounted and projected values paralleled those of present values (Table 2). For both sets, lowest values were in the narrow RZs and highest values were in the wide RZs. For an individual RZ, the lowest discounted value was US\$ 1.40 per acre and the highest projected value was US\$ 914.21 per acre (Burns, 1993).

Average discounted compound interest rates (1981–1989) for pine saw timber values were 7.2%, 6.6%, and 8.5% in the narrow, medium, and wide RZs, respectively; average projected compound interest rates (1989–1997) were 6.7%, 7.7%, and 8.8%, respectively. Discounted hardwood saw timber rates were 4.2%, 3.1%, and 3.7%, respectively; projected rates were 8.5%, 4.8%, and 3.1%, respectively.

Net annual equivalents of the total timber value (i.e. original timber volume and the timber growth) for the discount period (1981–1989) ranged US\$ 0.22–US\$ 68.21 per acre. Average values were US\$ 4.52 for the narrow RZs, US\$ 20.46 for the medium RZs, and US\$ 34.38 for the wide RZs. Except for the narrow RZs, most of the value was in the saw timber component.

Stumpage values at the time of the harvest projected at 7% compound interest for 30 years, approximately the length of the rotation used by forest industry, ranged US\$ 10.66–US\$ 3547.55 per acre of RZ. Aver-

age values were US\$ 242.60, US\$ 1083.07, and US\$ 1794.06 per acre for narrow, medium, and wide RZs, respectively. For 80 years, which is similar to a USFS rotation, values ranged US\$ 313.93–US\$ 104 499.95 per acre and averaged US\$ 7148.59, US\$ 31 904.07, and US\$ 52 847.56 per acre for narrow, medium, and wide RZs, respectively.

#### 4. Discussion

Although the width of the RZ has been generalized as an indicator of harvesting intensity, the RZ is merely the area not site prepared and planted to pine. The width of the RZ may not be as important to timber and wildlife values as the harvesting intensity within it. In this study, most narrow RZs had been heavily harvested, leaving few trees with little commercial value. The medium RZs usually had more volume per acre, but most trees present were hardwoods. The wide RZs generally had the most pine timber and thus the highest per-acre value. However, a wide RZ in which most merchantable timber is harvested will contain lower timber volumes and thus values than will a narrow RZ where the timber is left intact.

By leaving merchantable timber in the RZs, the landowner sacrifices present and future income streams. In a Georgia study, Caulfield et al. (1993) found that although RZs comprised only 8.2% of the land area, the opportunity costs were 14.4%. In this

study, the landowner may have sacrificed a NAE of as much as US\$ 68.21 per RZ acre per year in the wide RZs. If such RZs average 8% of the total forested area, this equates to US\$ 5.46 per forested acre per year. This exceeds the current average per-acre value for an unimproved hunting lease in eastern Texas. The value of such a lease on industrial forest lands is approximately US\$ 2.60 per acre per year (R.M. Whiting, unpublished data).

Medium RZs are the transition from the wide to the narrow RZs. In these RZs, the landowner may have sacrificed a NAE of as much as US\$ 34.05 per RZ acre per year. There are no known data relating the proportion of the stand occupied by RZs to the width of the RZs, but if we assume that medium RZs occupy only 6% of the total forested area, this represents US\$ 2.04 per forested acre per year. At the time that the adjacent stands were harvested, that value exceeded the per-acre rate that hunting clubs were paying to lease the lands.

Although the narrow RZs generally had the least merchantable timber, the landowner may still have been sacrificing as much as US\$ 11.13 per RZ acre per year. Narrow RZs had the greatest relative variability in NAEs, with a range-to-mean ratio of 240% compared to ratios of 119% and 172% for the medium and wide RZs, respectively (Bums, 1993).

Obviously landowners sacrifice significant economic returns by retaining RZs, and all else being equal, the wider the RZ, the greater the potential economic loss. However, wide, lightly-harvested or unharvested RZs have a wide range of ecological benefits, which, unfortunately, are less easily quantifiable than financial benefits (Caulfield et al., 1993).

While young pine plantations can provide suitable habitat for many early-succession wildlife species such as white-tailed deer (*Odocoileus virginianus*), northern bobwhite (*Colinus virginianus*), and cottontails (*Sylvilagus floridanus*), they do not provide adequate habitat for species requiring hardwoods, snags, cavity trees, and large down woody material (Thill, 1990). The habitat requirement of such species may be satisfied by wide RZs in which mature forests are maintained. Also, such RZs provide travel corridors connecting forest fragments and may enhance faunal diversity.

In young pine plantations, wide RZs are essential for gray squirrels (*Sciurus carolinensis*) and fox squir-

rels (*S. nigra*) (Dickson and Huntley, 1987) because they usually contain large, mast-bearing trees. In fact, both species may be restricted to RZs until pines in the adjacent plantations bear cones; gray squirrels may be restricted through much of a 30-year rotation.

The mast trees in such RZs are also important to wild turkeys (*Meleagris gallopavo*) (Dickson, 1989) and white-tailed deer (Harper, 1990). Research by Burk et al. (1990) in Mississippi suggested that wild turkeys use RZs for traveling, feeding, roosting, and loafing. The open understory in such areas is particularly attractive to turkeys. Poteet (1990) found that white-tailed deer significantly preferred RZs during fall and winter; he did not evaluate the width of the zones, however.

Other species benefited by wide RZs with mature-forest components include short-tailed shrews (*Blarina brevicauda*) (Dickson and Williamson, 1988), several species of herptiles (i.e. reptiles and amphibians) (Rudolph and Dickson, 1990), and a wide variety of birds. Shrews and herptiles benefit from the intact overstory and midstory, sparse shrub and herbaceous vegetation, and abundant leaf litter (Dickson and Williamson, 1988; Rudolph and Dickson, 1990).

Such RZs also benefit some species of song birds (Whiting, 1978). While bird communities are generally abundant and diverse in young and old forest stands, they may be reduced in dense, middle-aged pine plantations that lack vegetational diversity (Whiting, 1978; Dickson and Bosch, 1990). Bird abundance and diversity are positively correlated with habitat patchiness and foliage diversity (Whiting, 1978) and wide RZs containing unharvested or lightly harvested mature forests usually provide such habitat characteristics. They also provide large, old trees and snags for cavity-nesting birds. As would be expected, the value of such RZs to song birds declines as the adjacent stands age and acquire mature-forest characteristics (Murray and Stauffer, 1995).

Narrow, heavily harvested RZs do favor some species. Some small mammals (i.e. fulvous harvest mice, *Reithrodontomys fulvescens*) need the low-growing vegetation, logs, and logging slash found in narrow RZs. Populations of such species may be higher in the RZs than in the adjacent young pine plantations (Dickson and Williamson, 1988). The understory in narrow RZs is usually more dense than

that in the adjacent young pine plantation. It also may contain more blackberry (*Rubus sp.*) thickets and thus provide escape cover for a variety of species, including northern bobwhite, American woodcock (*Scolopax minor*), and white-tailed deer. However, there are no studies suggesting that narrow RZs are critical habitat for a particular wildlife species.

The wildlife-related values of medium RZs lie somewhere between these extremes. The way they function is very dependent on the harvesting regime used in them. In lightly harvested medium RZs, some squirrels will be present as will some late-successional bird species. Conversely, animal communities in heavily harvested medium RZs may resemble those in narrow RZs (Dickson and Williamson, 1988).

Riparian zones also provide protection for soil and water attributes. However, there is little agreement on the necessary width or the acceptable harvesting regimes in RZs to provide such protection. In fact, medium-width, partially harvested RZs may be adequate for protecting most intermediate streams in the southeastern United States. For example, Pritchett and Fisher (1987) wrote that partial harvests seldom decrease water quality and that directional tree felling can minimize debris entering streams; Hunter's (1990) comments were similar. Conversely, in areas of steep topography, wide, unharvested RZs may be necessary to slow overland water flow and filter large sediment loads (Peare, 1992). Obviously, characteristics of RZs necessary to protect soil and water are dictated by site conditions.

## 5. Conclusions

Currently all states in the nation have regulatory or nonregulatory best management practice (BMP) programs for forestry-related activities (Ice et al., 1997). All BMPs include stream protection, thus directly or indirectly require RZs. However, this study demonstrates that land owners may sacrifice significant economic returns by retaining RZs, with wider RZs having greater potential economic losses.

There are financial considerations for managing timber in RZs, but there are also ecological concerns. Plant communities in RZs are not static, and succession may lead to communities comprised of tree species with low economic values, especially in the

southeastern United States. Likewise, climax or near-climax communities may have lower overall wildlife-related values than do the midsuccessional communities that are currently present in most southern RZs. Unfortunately, there is a paucity of information on managing timber in southern RZs. This study strongly demonstrates the need for such information.

## References

- Burk, J.D., Hurst, G.A., Smith, D.R., Leopold, B.D., Dickson, J.G., 1990. Wild turkey use of streamside management zones in loblolly pine plantations. In: Healy, W.M., Healy, G.B. (Eds.), Proceedings of the Sixth National Wild Turkey Symposium. National Wild Turkey Federation, Edgefield, SC, pp. 84-89.
- Burns, G.A., 1993. Volume and value of residual timber in riparian zones in East Texas pine plantations. M.S. thesis, Stephen F. Austin State University, Nacogdoches, TX, p. 117.
- Caulfield, J., Welker, J., Meldahl, R. 1993. Opportunity costs of streamside management zones on an industrial forest property. In: Brissette, J.C. (Ed.), Proceedings of the Seventh Biennial Southern Silvicultural Research Conference. U.S. For. Serv. Gen. Tech. Rep. SO-93, pp. 639-644.
- Dickson, J.G., 1989. Streamside zones and wildlife in southern U.S. forests. In: Greswell, R.E., Barton, B.A., Kershner, J.L. (Eds.), Practical approaches to riparian resource management: An educational workshop, U.S. Bur. of Land Manage., Billings, MT, pp. 131-133.
- Dickson, J.G., Bosch, M.L., 1990. U.S. Forest Service research and management of non-game birds in the South. In: Wells, N. (Complier), Non-game migratory bird workshop. U.S. Fish and Wildl. Serv., Fort Collins, CO, pp. 24-25.
- Dickson, J.G., Huntley, J.C., 1987. Riparian zones and wildlife in southern forest: The problem and squirrel relationships. In: Dickson, J.G., Maughan, O.E. (Eds.), Managing southern forests for wildlife and fish - a proceeding. U.S. For. Serv. Gen. Tech. Rep. SO-65, pp. 37-39.
- Dickson, J.G., Williamson, J.H., 1988. Small mammals in streamside management zones in pine plantations. In: Szaro, R.C., Severson, K.E., Patton, D.R. (Technical coordinators), Proceedings of the symposium on management of amphibians, reptiles and small mammals in North America. U.S. For. Serv. Gen. Tech. Rep. RM-166, pp. 375-378.
- Harper, K.S. 1990. An evaluation of a habitat suitability index for white-tailed deer in East Texas. MS thesis, Stephen F. Austin State University, Nacogdoches, TX, p. 43.
- Hunter, M.L. Jr., 1990. Wildlife, Forests, and Forestry. Prentice Hall, Engelwood Cliffs, NJ, p. 370.
- Ice, G.C., Stuart, G.W., Waide, J.B., Irland, L.C., Elletson, P.V., 1997. Twenty-five years of the Clean Water Act: How clean are forest practices. J. For. 95(7), 9-13.
- Murray, N.L., Stauffer, D.F., 1995. Nongame bird use of habitat in central Appalachian riparian forests. J. Wildl. Manage. 59, 78-88.

- Peare, D., 1992. The three S's of stream protection. *Tree Farmer* 92, 9.
- Poteet, M.L., 1990. White-tailed deer use of riparian zones and adjacent pine plantations in East Texas. MS thesis, Stephen F. Austin State University, Nacogdoches, TX, pp. 69.
- Pritchett, W.L., Fisher, R.F., 1987. Properties and management of forest soils. 2nd ed. Wiley and Sons, New York, p. 494.
- Rudolph, D.C., Dickson, J.G., 1990. Streamside zone width and amphibian and reptile abundance. *Southwest. Nat.* 35, 472-416.
- Texas Forest Service, 1989. Texas timber price trends. Texas Forest Service, College Station, TX, p. 8.
- Thill, R.E., 1990. Managing southern pine plantations for wildlife. In: Proceedings of the twenty-ninth IUFRO World Congress, Canadian Int. Union For. Res. Organ, pp. 58-68.
- United States Forest Service, 1985. Forest survey inventory work plan. For. Inventory and Anal. Res. Work Unit, So. For. Exp. Stn., Starkville, MS, p. 57.
- Whiting, R.M. Jr., Ph.D, 1978. Avian diversity in various age pine forests in East Texas. Ph.D. dissertation, Texas A& M University, College Station, TX, p. 160.

# Forest Ecology and Management

## Submission of manuscripts.

Manuscripts should be submitted in triplicate. Authors from the Americas, Australia, New Zealand and the Pacific are requested to send their manuscripts to Forest Ecology and Management, Dr. Richard F. Fisher, Department of Forest Science, Texas A&M University, College Station, TX 77843-2135, USA; all other authors are requested to send their manuscripts to the Editorial Office, Forest Ecology and Management, P.O. Box 181,1000 AD Amsterdam, The Netherlands.

**Authors in Japan please note:** Upon request, Elsevier Science Japan will provide authors with a list of people who can check and improve the English of their paper (*before submission*). Please contact our Tokyo office: Elsevier Science Japan, 1-9-15 Higashi-Azabu, Minato-ku, Tokyo 106-0044, Japan; tel. (+81)-3-5561-5033; fax (+81)-3-5561-5047.

**Electronic manuscripts:** Electronic manuscripts have the advantage that there is no need for the rekeying of text, thereby avoiding the possibility of introducing errors and resulting in reliable and fast delivery of proofs. For the initial submission of manuscripts for consideration, hardcopies are sufficient. For the processing of *accepted papers*, electronic versions are preferred. After *final acceptance*, your disk plus two, final and exactly matching printed versions should be submitted together. Double density (DD) or high density (HD) diskettes (3.5 or 5.25 inch) are acceptable. It is important that the file saved is in the native format of the wordprocessor program used. Label the disk with the name of the computer and wordprocessing package used, your name, and the name of the file on the disk. Further information may be obtained from the Publisher.

**Enquiries concerning manuscripts and proofs:** questions arising after acceptance of the manuscript, especially those relating to proofs, should be directed to Elsevier Science Ireland Ltd., Bay 15K, Shannon Industrial Estate, Shannon, Co. Clare, Ireland, tel. (+353-61) 471944; fax(+353-61) 472144.

**Advertising information:** Advertising orders and enquiries can be sent to: **USA, Canada and South America:** Mr Tino de Carlo, The Advertising Department, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010-5107, USA; phone: (+1) (212) 633 3815; fax: (1) (212) 633 3820; e-mail: [t.decarlo@elsevier.com](mailto:t.decarlo@elsevier.com) **Japan:** The Advertising Department, Elsevier Science K.K., 9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106, Japan; phone: (+81) (3) 5561 5033; fax: (+81) (3) 5561 5047. **Europe and ROW:** Rachel Gresle-Farthing, The Advertising Department, Elsevier Science Ltd., The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, UK; phone: (+44) (1865) 843565; fax: (+44) (1865) 843976; e-mail: [r.gresle-farthing@elsevier.co.uk](mailto:r.gresle-farthing@elsevier.co.uk)

**Orders, claims, and product enquiries:** please contact the Customer Support Department at the Regional Sales Office nearest you:

**New York:** Elsevier Science, PO Box 945, New York, NY 10159-0945, USA; Tel.: (+1) (212) 633 3730 [toll free number for North American customers: 1-888-4ES-INFO (437-4636)]; Fax: (+1) (212) 633 3680; E-mail: [usinfo-f@elsevier.com](mailto:usinfo-f@elsevier.com)

**Amsterdam:** Elsevier Science, PO Box 211,1000 AE Amsterdam, The Netherlands; Tel.: (+31) 20 4853757; Fax: (+31) 20 4853432; E-mail: [nlinfo-f@elsevier.nl](mailto:nlinfo-f@elsevier.nl)

**Tokyo:** Elsevier Science, 9-15, Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel.: (+81) (3) 5561 5033; Fax: (+81) (3) 5561 5047; E-mail: [info@elsevier.co.jp](mailto:info@elsevier.co.jp)

**Singapore:** Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192; Tel.: (+65) 434 3727; Fax: (+65) 337 2230; E-mail: [asiainfo@elsevier.com.sg](mailto:asiainfo@elsevier.com.sg)

**Rio de Janeiro:** Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; Tel.: (+55) (21) 509 5340; Fax: (+55) (21) 507 1991; E-mail: [elsevier@campus.com.br](mailto:elsevier@campus.com.br) [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]

**US mailing info:** *Forest Ecology and Management* (0378-1121) is published monthly by Elsevier Science B.V. (Molenwerf 1, Postbus 211,1000 AE Amsterdam). Annual subscription price in the USA is US\$ 2307.00 (valid in North, Central and South America), including air speed delivery. Second class postage rate is paid at Jamaica, NY 11431.

**USA POSTMASTERS:** Send address changes to *Forest Ecology and Management* Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003.

**AIRFREIGHT AND MAILING** in the USA by Publications Expediting Inc., 200 Meacham Avenue, Elmont, NY 11003.

**Forest Ecology and Management has no page charges**

For a full and complete Guide for Authors, please refer to *Forest Ecology and Management*, Vol. 113, No. 1, pp. 105-108. The instructions can also be found on the World Wide Web: access under <http://www.elsevier.nl> or <http://www.elsevier.com>

# LANDSCAPE AND URBAN PLANNING

An International Journal of Landscape Ecology,  
Landscape Planning and Landscape design

## Editor-in-Chief:

J.E. Rodiek, College of Architecture, Texas A & M University, College Station, TX 77843-3137, USA

## AIMS AND SCOPE

A journal concerned with conceptual, scientific and design approaches to land use. By emphasizing ecological understanding and a multi-disciplinary approach to analysis and planning and design, it attempts to draw attention to the interrelated nature of problems posed by nature and human use of land. In addition, papers dealing with ecological processes and interactions within urban areas, and between these areas and the surrounding natural systems which support them, will be considered. Papers in which specific problems are examined are welcome. Topics might include but are not limited to landscape ecology, landscape planning and landscape design. Landscape ecology examines how heterogeneous combinations of ecosystems are structured, how they function and how they change. Landscape planning examines the various ways humans structure their land use changes. Landscape design involves the physical strategies and forms by which land use change is actually directed. *Landscape and Urban Planning* is based on

the premise that research linked to practice will ultimately improve the human made landscape.

## Editorial Advisory Board:

I.D. Bishop, *Parkville, Vic., Australia*,  
E.G. Bolen, *Wilmington, NC, USA*,  
I.D. Bruns, *Schorndorf, Germany*,  
J.B. Byrom, *Edinburgh, UK*,  
T.C. Daniel, *Tucson, AZ, USA*,  
R.M. DeGraaf, *Amherst, MA, USA*,  
J.G. Fabos, *Amherst, MA, USA*,  
S. Gonzalez Alonso, *Madrid, Spain*,  
M. Hough, *Etobicoke, ON, Canada*,  
P. Jacobs, *Montreal, PQ, Canada*,  
D.S. Jones, *Melbourne, Vic., Australia*,  
H. Lavery, *Milton, Qld., Australia*,  
W.M. Marsh, *Flint, MI, USA*,  
D.L. Mitchell, *Dallas, TX, USA*,  
D.G. Morrison, *Athens, GA, USA*,  
J.I. Nassauer, *St. Paul, MN, USA*,  
M. Nelischer, *Guelph, ON, Canada*,  
D.D. Paterson, *Vancouver, BC, Canada*,  
A. Ramos, *Madrid, Spain*,  
P. Shepard, *Claremont, CA, USA*,  
O.R. Skage, *Alnarp, Sweden*,  
R.C. Smardon, *Syracuse, NY, USA*,  
G. Sorte, *Alnarp, Sweden*,  
F. Stearns, *Rhineland, WI, USA*,

A.C. Szaro, *Washington, DC, USA*,  
J.W. Thomas, *La Grande, OR, USA*,  
P.J. Trowbridge, *Ithaca, NY, USA*,  
T.H.D. Turner, *London, UK*,  
N.J. Vroom, *Wageningen, The Netherlands*,  
W.V. Wendler, *College Station, TX, USA*,  
B.-E. Yang, *Seoul, Korea*,  
E.H. Zube, *Tucson, AZ, USA*

ABSTRACTED/INDEXED IN  
Applied Ecology Abstracts,  
Biological Abstracts,  
Current Contents B & S,  
Environmental Periodicals  
Bibliography, Geobase,  
Geographical Abstracts,  
LandSearch.

## 1994 SUBSCRIPTION DATA

Volumes 28-30 (in 9 Issues)

Subscription price:

Dfl. 1080.00 (US \$584.00)

incl. Postage

ISSN 0169-2046



**ELSEVIER**  
SCIENCE PUBLISHERS

**Elsevier Science** Publishers,  
P.O. Box 211, 1000 AE Amsterdam,  
The Netherlands  
Fax: (020) 5803-203

Customers in the USA and Canada:  
Elsevier Science Publishers  
P.O. Box 845, Madison Square Station,  
New York, NY 10160-0757, USA  
Fax: (212) 633-3680

*The Dutch Guilder price quoted applies worldwide, except in the Americas (North, Central and South America). The US Dollar price quoted applies in the Americas only. Non-VAT registered customers in the European Community should add the appropriate VAT rate applicable in their country to the price.*