

S. Kumar · C. Echt · P. L. Wilcox · T. E. Richardson

Testing for linkage disequilibrium in the New Zealand radiata pine breeding population

Received: 13 September 2002 / Accepted: 29 April 2003 / Published online: 30 July 2003
© Springer-Verlag 2003

Abstract Linkage analysis is commonly used to find marker-trait associations within the full-sib families of forest tree and other species. Study of marker-trait associations at the population level is termed linkage-disequilibrium (LD) mapping. A female-tester design comprising 200 full-sib families generated by crossing 40 pollen parents with five female parents was used to assess the relationship between the marker-allele frequency classes obtained from parental genotypes at SSR marker loci and the full-sib family performance (average predicted breeding value of two parents) in radiata pine (*Pinus radiata* D. Don). For alleles (at a marker locus) that showed significant association, the copy number of that allele in the parents was significantly correlated, either positively or negatively, with the full-sib family performance for various economic traits. Regression of parental breeding value on its genotype at marker loci revealed that most of the markers that showed significant association with full-sib family performance were not significantly associated with the parental breeding values. This suggests that over-representation of the female parents in our sample of 200 full-sib families could have biased the process of detecting marker-trait associations. The evidence for the existence of marker-trait LD in the population studied is rather weak and would require further testing. The exact test for genotypic disequilibrium between pairs of linked or unlinked marker loci revealed non-significant LD. Observed genotypic frequencies at several marker loci were significantly different from the expected Hardy-Weinberg equilibrium. The possibilities of utilising marker-trait associations for early selection, among-family selection and selecting parents for the next generation of breeding are also discussed.

Communicated by D.B. Neale

S. Kumar (✉) · C. Echt · P. L. Wilcox · T. E. Richardson
N Z Forest Research Institute, Private bag 3020, Rotorua,
New Zealand.
e-mail: Satisch.Kumar@forestresearch.co.nz
Tel.: +64-7343-5899
Fax: +64-7343-5330

Keywords Linkage disequilibrium · SSR marker · Breeding value · Marker-assisted selection · *Pinus radiata*

Introduction

The availability of genetic markers has facilitated experimental studies to detect quantitative trait loci (QTLs) in many species. Loci affecting quantitative traits can be detected via their linkage with these markers, only if there is linkage disequilibrium (LD) between the marker and the actual QTL. However, in most outcrossed forest-tree populations the association between alleles at a marker locus and the true QTL is likely to be in linkage equilibrium (Strauss et al. 1992). To overcome this limitation, most QTL detection experiments in the forest trees use family structures (e.g. full-sibling detection populations) that generate LD. Using this approach, a number of studies have found putative QTLs influencing various economic traits in conifers (e.g. Knott et al. 1997; Kumar et al. 2000; Sewell et al. 2001).

Simulation studies (e.g. Johnson et al. 2000; Kumar and Garrick 2001) and case studies (e.g. Williams and Neale 1992) have shown that substantial additional genetic gain can be obtained using within-family marker-assisted selection (MAS) in conifers. There are two major areas where MAS could contribute to such gains, namely 'backward' selection (re-selection of parents) and 'forward' selection (selection of individuals within families), in tree-breeding programs. For successful integration of MAS for 'backward' selection and/or 'forward' selection, it is important that marker-QTL associations are stable across a variety of genetic backgrounds. However, high degrees of linkage equilibrium (which means that linkage-phase relationships differ among individuals) makes broader application of marker-assisted selection (MAS) in forest trees very complex. This characteristic generally precludes the use of MAS for across-family selection and requires that MAS for within-family selection be customised for each family.

Using a factorial mating design, Arcade et al. (1996) provided experimental evidence for the existence of marker-trait associations in a broader population of larch. Verhaegen et al. (1998) using a factorial design in *Eucalyptus*, found a significant relationship between the cumulative number of marker alleles in the parents with the full-sib family performance for various traits. Sewell et al. (2001) showed that marker-trait associations were stable across two related pedigrees of loblolly pine. Finding molecular markers such as these will enhance the rate of genetic gain by using such markers in the between-family and within-family selections (e.g. Neale et al. 2002).

Natural and artificial selection, genetic sampling effects due to small effective population sizes, hybridisation and restricted migration of gametes among subdivisions, can produce LD among loci. In addition, LD might arise from selection on epistatic interaction (Lewontin 1974) or genetic drift (Hill and Robertson 1968) caused by the geographic origin of the populations involved in the breeding program. Verhaegen et al. (1998), in a LD study of *Eucalyptus* hybrids, postulated the different natural origins of the populations involved as the cause of the LD in their study. The current breeding population of New Zealand radiata pine has originated from local 'land-race' stocks which in turn have resulted essentially from hybridisation between two distinct natural populations (Monterey and Año Nuevo).

Non-random mating and artificial selection over 3–4 generations have led to the current radiata-pine breeding population. It is expected that provenance-hybridisation, non-random mating and artificial selection could have caused LD in the current New Zealand radiata-pine breeding population. Availability of linkage maps (e.g. Devey et al. 1999; Sewell et al. 1999) constructed using highly polymorphic markers provides an opportunity to test if LD exists between alleles at linked loci in breeding populations that have gone through generations of artificial selection and breeding.

Testing the LD hypothesis by detecting marker-QTL associations in each of the families in a large breeding program, such as in New Zealand, can be prohibitively costly. In this study, we extend the approach, used by Arcade et al. (1996) and Verhaegen et al. (1998) for dominant markers (RAPD) to codominant markers, to test the existence of LD at the population level between a series of marker loci and trait loci for various growth and form traits. The other two objectives of this study were to test for genotypic disequilibrium between pairs of various linked and unlinked marker loci; and to test the existence of disequilibrium at a given marker locus [this is similar to testing for departure from Hardy-Weinberg (H-W) equilibrium].

Materials and methods

Genetic material and female-tester mating design

The concept of "female-testing" assumes that the tested progeny of crosses of candidate pollen parents with four or five females give an accurate estimate of the breeding value or the general combining ability (GCA) of the pollen parents. A field trial consisting of 750 pair-crosses (150 pollen parents mated factorially with each of five female parents) was established in 1994 with six replicates of each pair cross. Each tree was assessed at the age of 6-years for diameter at breast height (mm), stem straightness (1–9 scale; 1 = most crooked, 9 = very straight), branching cluster frequency (1–9 scale; 1 = fewest branch clusters, 9 = most clusters) and wood density (kg/m^3).

To calculate variance components and to predict the parental breeding values (which will be used to study marker-trait association), the following linear model was used:

$$Y_{ijkl} = \mu + R_i + P_j + T_k + PT_{jk} + e_{ijkl}, \quad (1)$$

where Y_{ijkl} is the observation for the i^{th} progeny of the cross between k^{th} tester and j^{th} pollen parent in the i^{th} replicate; μ is an overall mean; R_i is the fixed effect associated with the i^{th} replicate; P_j is the random effect associated with the j^{th} pollen parent; T_k is the random effect associated with the k^{th} female tester; PT_{jk} is the random interaction associated with the j^{th} pollen parent and k^{th} female tester; and e_{ijkl} is the random error.

The random effects, P_j , T_k , PT_{jk} and e_{ijk} , are assumed to have expected values equal to zero and variances σ_e^2 , σ_i^2 , σ_{ct}^2 and σ_{e}^2 , respectively. ASREML software (Gilmour et al. 1997) was used to estimate genetic parameters and to obtain the best linear unbiased prediction (BLUP) of parental breeding values.

Molecular-marker assays

For the purpose of this linkage disequilibrium study, genotyping of the 45 parents (40 pollen parents and 5 female parents) using 34 polymorphic SSR markers was out-sourced to *SignaGen*—Molecular Breeding Solutions (<http://www.SignaGen.com>). All individuals produced reliable 2-allele genotypes at each locus. More detail of the methodology is available on request from *SignaGen*. Biallelic genotypes at each of the 34 loci were obtained for each of the 45 parents and the allele frequencies were obtained.

As shown in Table 1, frequencies of observed alleles varied from 0.01 to 0.82. Most of the alleles were present in very low frequency. The three highest-frequency alleles were first identified and the remaining low-frequency alleles were lumped to form a "synthetic" 4th allele. This resulted in four alleles (three distinct alleles and one synthetic allele) at each locus. If there were originally only four observed alleles at a locus, and one of the alleles had a frequency less than 5%, then it was lumped with the 3rd most-frequent allele and thus resulted in three alleles (two distinct alleles and one synthetic allele) at all such loci. Synthetic alleles were used because if the true rare allele was present only in female parents, it may bias the process of detecting marker-trait association in the breeding population using the female-tester design (Ewens and Spielman 1995; Pritchard and Rosenberg 1999). Biallelic genotypes of all 45 parents were obtained after forming the "synthetic" alleles at each marker locus and were used for all further calculations.

LD between a marker and a trait locus

Analysis of variance (ANOVA) is commonly used to study the association between marker genotypes and trait loci in segregating populations. Arcade et al. (1996) and Verhaegen et al. (1998) used one-way ANOVA to investigate the statistical relationship between the expected RAPD marker-'band present' allele-frequency in the offspring, and family performance. The expected frequency of the

Table 1 Number of alleles and their frequencies (calculated from a sample size of 45 individuals) at each marker loci. The linkage groups to which a marker was assigned to are based on a QTL mapping pedigree

Locus abbreviation	No. of alleles	Range of allele frequencies	No. of alleles after lumping	Linkage group
NZPR0119	9	0.01–0.34	4	15
NZPR0129	21	0.01–0.24	4	6
NZPR0138b	14	0.01–0.23	4	3
NZPR0143	9	0.02–0.25	4	3
NZPR0157	12	0.02–0.23	4	1
NZPR0248b	10	0.01–0.35	4	15
NZPR0290	23	0.01–0.15	4	1
NZPR0407	15	0.01–0.27	4	11
NZPR0413	14	0.01–0.25	4	4
NZPR0440	4	0.03–0.73	3	11
NZPR0443	8	0.01–0.53	4	7
NZPR0476	7	0.01–0.82	3	16
NZPR0495	9	0.01–0.29	4	11
NZPR0581	7	0.01–0.35	4	10
NZPR0599	10	0.01–0.19	4	16
NZPR0706	7	0.01–0.37	4	1
NZPR0826	6	0.01–0.56	4	6
NZPR0947	3	0.08–0.70	3	15
NZPR1078	4	0.01–0.76	4	3
PR4.6	14	0.01–0.32	4	1
RIPPT0079a	6	0.01–0.35	4	7
RIPPT0699	18	0.01–0.40	4	1
RIPPT0914	7	0.02–0.61	4	2
RIPPT0984	22	0.01–0.20	4	10
UCC0705	4	0.03–0.48	3	7
UCC0719	5	0.01–0.69	3	16
UCC0735	8	0.01–0.60	4	2
UCC0759	16	0.01–0.24	4	6
UCC0784	10	0.01–0.45	4	8
UCC0802	14	0.01–0.34	4	3
UCC0844	11	0.01–0.32	4	3
UCC0922	14	0.01–0.39	4	6
UCC1017	15	0.01–0.21	4	4
UCC1149	9	0.04–0.35	4	6

"band present" allele-frequency in the offspring was obtained from the parental genotypes of the family. Because RAPDs are dominant markers, they grouped full-sib families into either two or three frequency classes, depending on whether the marker was polymorphic in one or both parents. Using codominant markers in the present study, it was possible to group full-sib families in up to five expected frequency classes of an allele, depending upon the marker genotypes of the parents (see Fig. 1).

Lande and Thompson (1990) suggested the use of the regression of phenotypic records of the quantitative trait on the number of favourable marker alleles at a marker locus to infer the presence of linkage disequilibrium. Luo (1998), using simulation studies, compared the power of ANOVA and regression approaches, and reported that regression analysis is more powerful than the ANOVA, particularly when the QTL has low heritability. In this study, the concept of Arcade et al. (1996) and Verhaegen et al. (1998) was fitted using a linear regression approach. For each of the alleles at 34 marker loci, the linear model was written as:

$$F_i = \mu + \gamma_{ik} + e_{ik}, \quad (2)$$

where F_i is the expected performance (defined as the average breeding value of the two parents; breeding value estimates were obtained from the female-tester design as explained earlier) of i^{th} full-sib family; μ is the general mean; γ_{ik} is the expected frequency (calculated from the parental marker genotypes) of allele j at the k^{th} locus in the full-sib progeny; e_{ik} is the random error. A total of 524 regression analyses were conducted for various combinations of alleles at different marker loci and trait phenotypes. The significance of the regression coefficient was used to infer LD at the population level. As the mating design structure involved only five female parents, some spurious associations between marker and the pair-cross effects are expected. Thus, the analysis of regression of

parental breeding values or general combining ability (GCA) on marker allele-frequency was also undertaken to infer LD in the breeding population. The sample sizes within each of the allele-frequency class could vary and thus to account for this unbalanced structure, the threshold levels for testing the significance of the regression coefficient was obtained using the permutation method (Churchill and Doerge 1994).

LD between pairs of marker loci

Genotypic data on pairs of loci was used to calculate the LD between pairs of marker loci. When working with genotypic data from natural populations, it is not reasonable to assume that random-mating occurs, and gamete frequencies are seldom known. Hill (1974) proposed a method to calculate LD coefficients when the population undergoes the random union of gametes. As some of the single-locus results for our population indicate departures from H-W frequencies (see below), an assumption that the random union of gametes can not be assumed. Weir (1979) proposed a composite measure of LD, which is appropriate under conditions of the non-random union of gametes and incomplete identification of genotypes. To test the genotypic LD between a pair of loci, the null hypothesis that the genotypes at one locus are independent from genotypes at the other locus was assumed.

The analysis was implemented using GENEPOP software (Raymond and Rousset 1995). Using this method, a contingency table based on the observed genotypic data, is prepared for each pair of loci and an unbiased estimate of the P -value of the probability test, or the exact test is performed using a Markov Chain method. The P -value of the test is calculated as the sum of the probabilities of all contingency tables (with the same marginal

Using a factorial mating design, Arcade et al. (1996) provided experimental evidence for the existence of marker-trait associations in a broader population of larch. Verhaegen et al. (1998) using a factorial design in *Eucalyptus*, found a significant relationship between the cumulative number of marker alleles in the parents with the full-sib family performance for various traits. Sewell et al. (2001) showed that marker-trait associations were stable across two related pedigrees of loblolly pine. Finding molecular markers such as these will enhance the rate of genetic gain by using such markers in the between-family and within-family selections (e.g. Neale et al. 2002).

Natural and artificial selection, genetic sampling effects due to small effective population sizes, hybridisation and restricted migration of gametes among subdivisions, can produce LD among loci. In addition, LD might arise from selection on epistatic interaction (Lewontin 1974) or genetic drift (Hill and Robertson 1968) caused by the geographic origin of the populations involved in the breeding program. Verhaegen et al. (1998), in a LD study of *Eucalyptus* hybrids, postulated the different natural origins of the populations involved as the cause of the LD in their study. The current breeding population of New Zealand radiata pine has originated from local 'land-race' stocks which in turn have resulted essentially from hybridisation between two distinct natural populations (Monterey and Año Nuevo).

Non-random mating and artificial selection over 3–4 generations have led to the current radiata-pine breeding population. It is expected that provenance-hybridisation, non-random mating and artificial selection could have caused LD in the current New Zealand radiata-pine breeding population. Availability of linkage maps (e.g. Devey et al. 1999; Sewell et al. 1999) constructed using highly polymorphic markers provides an opportunity to test if LD exists between alleles at linked loci in breeding populations that have gone through generations of artificial selection and breeding.

Testing the LD hypothesis by detecting marker-QTL associations in each of the families in a large breeding program, such as in New Zealand, can be prohibitively costly. In this study, we extend the approach, used by Arcade et al. (1996) and Verhaegen et al. (1998) for dominant markers (RAPD) to codominant markers, to test the existence of LD at the population level between a series of marker loci and trait loci for various growth and form traits. The other two objectives of this study were to test for genotypic disequilibrium between pairs of various linked and unlinked marker loci; and to test the existence of disequilibrium at a given marker locus [this is similar to testing for departure from Hardy-Weinberg (H-W) equilibrium].

Materials and methods

Genetic material and female-tester mating design

The concept of "female-testing" assumes that the tested progeny of crosses of candidate pollen parents with four or five females give an accurate estimate of the breeding value or the general combining ability (GCA) of the pollen parents. A field trial consisting of 750 pair-crosses (150 pollen parents mated factorially with each of five female parents) was established in 1994 with six replicates of each pair cross. Each tree was assessed at the age of 6-years for diameter at breast height (mm), stem straightness (1–9 scale; 1 = most crooked, 9 = very straight), branching cluster frequency (1–9 scale; 1 = fewest branch clusters, 9 = most clusters) and wood density (kg/m^3).

To calculate variance components and to predict the parental breeding values (which will be used to study marker-trait association), the following linear model was used:

$$Y_{ijkl} = \mu + R_i + P_j + T_k + PT_{jk} + e_{ijkl}, \quad (1)$$

where Y_{ijkl} is the observation for the i^{th} progeny of the cross between k^{th} tester and j^{th} pollen parent in the i^{th} replicate; μ is an overall mean; R_i is the fixed effect associated with the i^{th} replicate; P_j is the random effect associated with the j^{th} pollen parent; T_k is the random effect associated with the k^{th} female tester; PT_{jk} is the random interaction associated with the j^{th} pollen parent and k^{th} female tester; and e_{ijkl} is the random error.

The random effects, P_j , T_k , PT_{jk} and e_{ijk} , are assumed to have expected values equal to zero and variances σ_e^2 , σ_t^2 , σ_{ct}^2 and σ_{et}^2 , respectively. ASREML software (Gilmour et al. 1997) was used to estimate genetic parameters and to obtain the best linear unbiased prediction (BLUP) of parental breeding values.

Molecular-marker assays

For the purpose of this linkage disequilibrium study, genotyping of the 45 parents (40 pollen parents and 5 female parents) using 34 polymorphic SSR markers was out-sourced to *SignaGen*—Molecular Breeding Solutions (<http://www.SignaGen.com>). All individuals produced reliable 2-allele genotypes at each locus. More detail of the methodology is available on request from *SignaGen*. Biallelic genotypes at each of the 34 loci were obtained for each of the 45 parents and the allele frequencies were obtained.

As shown in Table 1, frequencies of observed alleles varied from 0.01 to 0.82. Most of the alleles were present in very low frequency. The three highest-frequency alleles were first identified and the remaining low-frequency alleles were lumped to form a "synthetic" 4th allele. This resulted in four alleles (three distinct alleles and one synthetic allele) at each locus. If there were originally only four observed alleles at a locus, and one of the alleles had a frequency less than 5%, then it was lumped with the 3rd most-frequent allele and thus resulted in three alleles (two distinct alleles and one synthetic allele) at all such loci. Synthetic alleles were used because if the true rare allele was present only in female parents, it may bias the process of detecting marker-trait association in the breeding population using the female-tester design (Ewens and Spielman 1995; Pritchard and Rosenberg 1999). Biallelic genotypes of all 45 parents were obtained after forming the "synthetic" alleles at each marker locus and were used for all further calculations.

LD between a marker and a trait locus

Analysis of variance (ANOVA) is commonly used to study the association between marker genotypes and trait loci in segregating populations. Arcade et al. (1996) and Verhaegen et al. (1998) used one-way ANOVA to investigate the statistical relationship between the expected RAPD marker-'band present' allele-frequency in the offspring, and family performance. The expected frequency of the

Table 3 Summary statistics, at the individual-tree level, and estimates of narrow-sense heritability (h^2) of various traits: diameter at breast height (DBH), straightness (STR), branch cluster frequency (BR) and wood density (WD)

Statistics	DBH (mm)	STR (1–9 scale)	BR (1–9 scale)	WD (kg/m ³)
Mean	174	6.11	5.76	353
Minimum	63	1	1	271
Maximum	300	9	9	439
h^2	0.14	0.17	0.20	0.68

Table 4 Significant markers and the variance explained for different traits

Trait	Marker	R ² (%) for the positive allele	R ² (%) for the negative allele
DBH	NZPR0440	30.0	47.8
	UCC0784	25.4	23.6
BR	NZPR0143	41.8	46.3
	NZPR0138b	42.8	33.6
STR	NZPR0290	21.5	24.8
	NZPR0706	19.7	14.6
WD	UCC0719	9.7	14.20




Fig. 1 Association of the alleles 1 and 3 with the family performance for branching (BR) at the marker locus NZPR0143. The other alleles at this locus did not show any association.

Expected frequency of the allele in the full-sib families is shown on the X-axis and the expected family performance is shown on the Y-axis

correction, none of the LD tests between marker loci remained significant. As the majority of the marker loci were mapped with a recombination fraction of more than 0.05, one would not expect to detect strong genotypic disequilibrium among various pairs of loci.

LD between a marker and a trait locus

Marker-trait associations were evaluated for various traits measured at 6-years of age: diameter at breast height (DBH), stem straightness (STR), branching cluster frequency (BR) and wood density (WD). Summary statistics along with estimates of narrow-sense heritability of various traits, obtained from complete trial data, are shown in Table 3. At the individual-tree level, the average DBH, STR, BR and WD were 174, 6.11, 5.76 and 353 respectively. Estimates of narrow-sense heritabilities were 0.14, 0.17, 0.20 and 0.68 for DBH, STR, BR and WD respectively.

Molecular markers that showed significant (at the 1% experiment-wise threshold level) associations with the full-sib family performance for a number of traits, are shown in Table 4. In such cases, at each marker locus, the alleles showing positive, negative or no effect on the

family performance were clearly evident. For example, Fig. 1 shows that at locus NZPR0143 the average family performance for BR increases as the expected frequency of allele 1 increases in the progeny. The average family performance for the BR score was found to be -0.38, -0.25, 0.03, 0.019 and 0.38 when the expected frequency of allele 1 in the families was 0, 0.25, 0.50, 0.75 and 1.0, respectively. At locus NZPR0143 allele 3 shows an association with a negative QTL allele for branching. For allele frequency of 0, 0.25, 0.50, 0.75 and 1.0, the expected family performance was 0.18, -0.05, -0.25, -0.51 and -0.48 (Fig. 1). The molecular marker (NZPR0143) showing association with the BR was mapped in the same region that showed significant association with BR in a QTL-mapping pedigree (Wilcox et al. 1998). For WD, the average expected family performance increased from -5.24 to 2.91 kg/m³ when the frequency of allele 3 increased from 0 to 1. Similarly, the average WD of families decreased from 2.11 to -6.17 kg/m³ with a change of allele-2 frequency from 0 to 1 (Fig. not shown).

As the design used in this study involved only five female parents and 40 male parents, there is a possibility that the significant associations found at different marker loci might be influenced by the fact that these alleles are

Table 5 Estimated frequency of the favourable and unfavourable alleles in male and female populations, at the selected marker loci

Locus	Frequency of the positive-effect allele in		Frequency of the negative-effect allele in	
	Males	Females	Males	Females
UCC0784	0.22	0.30	0.18	0.40
NZPR0440	0.14	0.40	0.76	0.50
NZPR0143	0.41	0.50	0.20	0.33
NZPR0138b	0.44	0.36	0.24	0.17
NZPR0290	0.14	0.25	0.64	0.75
NZPR0706	0.35	0.50	0.21	0.10
UCC0719	0.16	0.50	0.71	0.50

segregating only in the female population. Any marker allele that is in high frequency in the over-represented sub-population (i.e. five female testers in our case) would show association with the phenotype (Ewens and Spielman 1995; Pritchard and Rosenberg 1999). To study this aspect, the frequency of the alleles showing significant association with full-sib family performance, was calculated for male and female populations separately. Results shown in Table 5 indicated that the frequency of favourable alleles was generally higher in the female population, but no such trend was observed for unfavourable alleles. These results reflected that the evidence of LD found using full-sib family performance could have been influenced by the mating-design structure.

To resolve this issue further, we investigated the relationship between the predicted breeding value (or GCA) of a parent to its genotype at the marker loci that showed a significant LD with the full-sib family performance. Most of the markers showing significant association with the full-sib family performance in the initial analysis (see Table 4) were found to show non-significant associations at the parental GCA level. However, allele 3 and allele 2 at the marker loci NZPR0143 and UCC0719, respectively, were significantly ($P = 0.0004$ and 0.0541 , respectively) associated with the predicted breeding values for BR and WD, respectively. After applying Bonferroni correction, allele 3 at the marker loci NZPR0143 remained significant at 20% experiment-wise threshold.

It is interesting to note that all marker-trait associations found using full-sib family performance could not be repeated when the parental GCA was regressed on the marker-allele frequencies. There could be various explanations for these results. The number of significant tests found (7 out of 524) were not much different from that would be expected (5 out of 524) due to chance alone at a probability level of 0.01. Thus, most of the significant marker-trait associations could be due to the genetic-sampling effects. Mating design structure could also have contributed to these results. Over-representation of five female parents in our sample of 200 full-sib families might have contributed to the observed significant marker-trait associations (e.g. Ewens and Spielman 1995; Pritchard and Rosenberg 1999). These results clearly indicate that the existence of LD, found using full-sib performance (as was the case in Arcade et al. 1996 and Verhaegen et al. 1998), could be verified using

GCA effects of parents involved in those full-sib crosses. We suggest this should be a statistical validation of the association found at the full-sib family level. Our results suggest that the evidence of marker-trait LD, in the population of 45 parents that we studied, is rather weak. Various candidate genes influencing wood property traits, along with the markers showing putative associations with trait loci in this pilot study, are now being evaluated using a larger set of the New Zealand radiata pine-breeding population parents.

Marker-assisted selection for breeding

Various authors have advocated the use of molecular markers in advanced generation tree-breeding programmes (e.g. O'Malley and McKeand 1994; Johnson et al. 2000; Kumar and Garrick 2001). A prominent feature of advanced generation programmes is the between-and-within-family selection, where full-sib families are culled based on the mid-parent GCA values and the best offspring are chosen for breeding from the remaining families. In the New Zealand radiata pine-breeding programme, various new elite (or breed) populations are being formed for various economic traits (e.g. structural and appearance timber grades) by selecting 24 parents from the main breeding population (Jaywickrama and Carson 2000). These parents are crossed factorially to create full-sib families where advanced generation selections are made for breeding and deployment purposes. Marker-trait associations found using full-sib family performance and parental GCA values, as described in this study, would indicate the QTLs of 'general value' that can be used for between-and-within-family selection in elite populations. Detecting LD in small elite populations, however, might not be generalised for the wider radiata pine populations. The method described here would assist finding QTLs that are specific to the best existing parental combinations. The most heralded benefit of MAS in forestry is the use of marker-trait associations for early selection. The marker-trait LD found using the approach presented in this study could be used for early selection of individuals for mass propagation or for clonal testing.

Acknowledgements We thank the New Zealand Radiata Pine Breeding Company for giving access to phenotypic data. We thank Dr. Christophe Plomion and Dr. Rowland Burdon for helpful

discussion during the completion of this project. We also thank anonymous referees for very helpful and constructive comments. The experiments comply with the current laws of New Zealand where the experiments were performed. SSR markers with prefixes RIPPT and UCC are exclusive property of International Paper Company. We thank them for the use of these markers for this study.

References

- Arcade A, Faivre-Rampant P, Le Guerroue B, Paques LE, Prat D (1996) Quantitative traits and genetic markers: analysis of a factorial mating design in larch. In: Ahuja MR, Boerjan W, Neale D (eds) Somatic cell genetics and molecular genetics of trees. Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 211–216
- Churchill GA, Doerge RW (1994) Empirical threshold values for quantitative trait mapping. *Genetics* 138:963–971
- Devey ME, Sewell MM, Uren TL, Neale DB (1999) Comparative mapping in loblolly pine and radiata pine using RFLP and microsatellite markers. *Theor Appl Genet* 99:656–652
- Ewens WJ, Spielman RS (1995) The transmission/disequilibrium test: history, subdivision, and admixture. *Am J Hum Genet* 57:455–464
- Gilmour AR, Thompson R, Cullis BR, Welham SJ (1997) ASREML. NSW Agriculture, Orange, NSW, Australia
- Haldane JBS (1954) An exact test for randomness of mating. *J Genet* 52:631–635
- Hill WG (1974) Estimation of linkage disequilibrium in random mating populations. *Heredity* 33:229–239
- Hill WG, Robertson A (1968) Linkage disequilibrium in finite populations. *Theor Appl Genet* 38:226–231
- Jayawickrama, KJS, Carson MJ (2000) A breeding strategy for the New Zealand Radiata Pine Breeding Cooperative. *Silvae Genet* 49:82–90
- Johnson GR, Wheeler NC, Strauss SH (2000) Financial feasibility of marker-aided selection in Douglas-fir. *Can J For Res* 30:1942–1952
- Knott SA, Neale DB, Sewell MM, Haley CS (1997) Multiple-marker mapping of quantitative trait loci in an outbred pedigree of loblolly pine. *Theor Appl Genet* 94:810–820
- Kumar S, Garrick DJ (2001) Genetic response to within-family selection using molecular markers in some radiata pine breeding schemes. *Can J For Res* 31:779–785
- Kumar S, Spelman RJ, Garrick DJ, Richardson TE, Lausberg M, Wilcox PL (2000) Multiple-marker mapping of wood density loci in an outbred pedigree of radiata pine. *Theor Appl Genet* 100:926–933
- Lande R, Thompson R (1990) Efficiency of marker-assisted selection in the improvement of quantitative traits. *Genetics* 124:743–756
- Lewontin RC (1974) The genetic basis of evolutionary change. Columbia Univ. Press, New York
- Louis EJ, Dempster ER (1987) An exact test for Hardy-Weinberg and multiple alleles. *Biometrics* 43:805–811
- Luo ZW (1998) Detecting linkage disequilibrium between a polymorphic marker locus and a trait locus in natural populations. *Heredity* 80:198–208
- Neale DB, Sewell MM, Brown GR (2002) Molecular dissection of the quantitative inheritance of wood property traits in loblolly pine. *Ann For Sci* 59:595–605
- O'Malley DM, McKeand SE (1994) Marker-assisted selection for breeding value in forest trees. *For Genet* 1:207–218
- Pritchard JK, Rosenberg NA (1999) Use of unlinked genetic markers to detect population stratification in association studies. *Am J Hum Genet* 65:220–228
- Raymond M, Rousset F (1995) GENEPOP: population genetics software for exact tests and ecumenicism. *J Hered* 86:248–249
- Rousset F, Raymond M (1995) Testing heterozygote excess and deficiency. *Genetics* 140:1413–1419
- Sewell MM, Sherman BK, Neale DB (1999) A consensus map for loblolly pine (*Pinus taeda* L.). I. Construction and integration of individual linkage maps from two outbred three-generation pedigrees. *Genetics* 151:321–330
- Sewell MM, Bassoni DL, Megraw RA, Davis, MF, Wheeler, NC, Tuskan, GA, Neale DB (2001) Mapping QTLs for wood property traits in loblolly pine (*Pinus taeda* L.) with application towards marker-assisted breeding (Abstract). Wood, Breeding and Biotechnology Workshop, Bordeaux, 11th–14th, June 2001
- Strauss SH, Lande R, Namkoong G (1992) Limitations of molecular marker-aided selection in forest-tree breeding. *Can J For Res* 22:1050–1061
- Verhaegen D, Plomion C, Poitel M, Costa P, Kremer A (1998) Quantitative trait dissection analysis in *Eucalyptus* using RAPD markers 2. Linkage disequilibrium in a factorial design between *E. urophylla* and *E. grandis*. *For Genet* 5:61–69
- Weir BS (1979) Inferences about linkage disequilibrium. *Biometrics* 35:235–254
- Weir BS, Cockerham CC (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution* 38:1358–1370
- Wilcox PL, Richardson TE, Corbett GC, Lee JR, Lausberg M, Carson SD (1998) Results from Phase-II QTL detection experiments. New Zealand Forest Research Institute, Report No. 61
- Williams CG, Neale DB (1992) Conifer wood quality and marker-aided selection: a tool for the improvement of forest-tree species. *Can J For Res* 22:1009–1017