

U.S. Department of Agriculture

Forest Service Research Note SE-297 July 1980

ECTOMYCORRHIZAL COLONIZATION OF CONTAINER-GROWN NORTHERN RED OAK AS AFFECTED BY FERTILITY

by John L. Ruehle[1]

ABSTRACT.—The effects of different fertility regimes on *Pisolithus tinctorius* ectomycorrhizal development on northern red oak grown in containers was tested in the greenhouse. Artificial infestation of a milled pine bark-sewage sludge growing medium produced the best ectomycorrhizal development (63 percent) when a nutrient solution with 100 μg/ml N and 20 μg/ml P was used every 3 weeks. A commercially available nutrient solution (NP-500 μg/ml) produced the largest seedlings, those rated suitable for planting, but caused a marked reduction in ectomycorrhizal development (Pt = 20 percent).

The results indicate that the production of large and vigorous northern red oak seedlings in containers with abundant *Pisolithus* ectomycorrizae is strongly dependent on the levels of available nutrients in the growing medium.

Keywords: *Pisolithus tinctorius, Quercus rubra.*

Northern red oak (*Quercus rubra* L.) grows best in upland coves and middle and lower slopes having a fine-textured soil and topography favoring a high water table (Core 1971). Intense competition makes it difficult to reproduce red oak stands on the better sites. There is a high mortality of newly established oak seedlings in this region (Clark and Watt 1971). Although reducing competition helps artificial oak regeneration, mowing and use of herbicides are not feasible in many areas in the uplands within the range of northern red oak (Russell 1971).

Conventional nursery stock has not demonstrated the ability to grow fast enough to compete with woody vegetation on forest sites (Johnson 1979). The advent of containerization in recent years offers a possible means of improving early seedling growth of oaks. Initial trials by Johnson (1974) and Wendel (1979) showed growth rates of containerized northern red oak to be too slow to be competitive. Johnson speculated that a larger container and a longer propagation period offered the best opportunities for improving the growth of outplanted tublings.

The use of ectomycorrhizal fungi ecologically adapted to the planting site in container operations may also improve survival and early seedling growth (Ruehle and Marx 1979). These symbiotic fungi assist the plant in water and mineral uptake by increasing the effective absorbing area of the feeder roots. Good nonmycorrhizal oak seedlings can be grown in containers in the greenhouse because conditions are controlled to minimize stress of all kinds (Tinus and McDonald 1979). However, such seedlings often fail to survive or grow well after planting, especially during a period of environmental stress. Pine seedlings colonized with *Pisolithus tinctorius* survive and grow better on routine reforestation sites than seedlings colonized with *Thelephora terrestris* or other fungi (Marx and others 1977a). Since this symbiont also colonizes oaks, it would be a good candidate for tailoring the roots of oaks before outplanting (Marx 1979). Previous work (Ruehle, unpublished data) has shown that northern red

[1]Ruehle is Plant Pathologist, Institute for Mycorrhizal Research and Development, USDA Forest Service, Forestry Sciences Laboratory, Athens, Georgia 30602.

oak became colonized with *P. tinctorius* when inoculum was added to the growing medium. However, the colonization was erratic and usually quite low (below 5 percent) in these tests where trees received a commercially available soluble fertilizer (NPK at 500 p/m each concentration) every 3 weeks. Before field trials can be conducted to evaluate the performance of containerized seedlings colonized with specific ectomycorrhizae selected to adapt the seedlings to the site, greenhouse experiments must be conducted to determine the optimum growing medium, inoculation techniques, and fertility levels needed to produce acceptable plantable seedlings. Current cultural practices for growing containerized oaks are often not suitable for colonization of roots by certain ectomycorrhizal fungi. Frequent watering of seedlings in containers with poor drainage (growing medium that is too fine often becomes waterlogged) prevents feeder-root development. Lack of feeder roots prevents adequate ectomycorrhizal development. Frequent application of high concentrations of water-soluble fertilizer also prevents mycorrhizal development by many symbiotic fungi (Marx and others 1977b).

The study was conducted to determine the best fertility regime for developing *P. tinctorius* ectomycorrhizae on northern red oak grown in containers.

METHODS AND MATERIALS

Milled pine bark used as a growing medium for containerized southern pine seedlings promotes extensive secondary-root development and a high percentage of ectomycorrhizae (Ruehle and Marx 1977), and was used in this study. Processed sewage sludge was added to milled pine bark because it permits less frequent application of soluble fertilizer. Sewage sludge used in this study was obtained from two primary plants in Athens, Georgia, screened and dried 48 hours at 80°C. The nutrient content and organic matter of this sludge was discussed by Berry and Marx (1976). The milled pine bark was fumigated with methyl bromide and then mixed 4:1 v/v with the processed sewage sludge to create the basic growing medium for this study. Inoculum of *P. tinctorius* (isolate 185) prepared according to standard procedures (Marx and Bryan 1975) was dried at room temperature to approximately 20 percent moisture, and mixed with the growing medium at a 1:15 v/v ratio.

Fifty plywood boxes were constructed with screen bottoms. Each was large enough to hold five Super-45 Spencer-LeMaire Rootrainers® (Spencer-LeMaire Industries Ltd., Edmonton, Alberta, Canada). Each Rootrainer had three cavities; each cavity was 575 cm^3. After each box was filled with Rootrainers (15 cavities/box), 200 cm^3 of growing medium was added to each cavity. The growing medium mixed with inoculum was then added (350 cm^3) to each cavity to fill it to within 3 cm of the top.

A mixed lot of northern red oak seed collected in western North Carolina was Clorox® treated, sorted by flotation, and stored in plastic bags in a refrigerator for 2 months. The seed was planted (one per cavity) in March 1978. Additional seed were planted in trays to provide transplants for cavities lacking seedlings. Three weeks after seeding, those cavities lacking seedlings received transplants from the trays.

A nutrient solution modified from one employed by Marx and others (1977b) and a Rapid-gro® solution (control) was used in this test to provide various levels of N and P (table 1). Treatments assigned to each group of 10 boxes are listed in table 2.

Table 1.—Modified nutrient solutions for fertility study on northern red oak grown in containers

Element	Source	Concentration
		μg/ml
N	NH_4NO_3	12.5, 25, 50, or 100
P	Na_2HPO_4	5 or 20
K	KCl	40
Ca	$CaCl_2$	40
Mg	$MgSO_4 \cdot 7H_2O$	20
Fe	Chelated Fe 10%	5.5
Mo	$NaMoO_4 \cdot 7H_2O$	.001
Cu	$CuSO_4 \cdot 5H_2O$	.006
Bo	H_3Bo_3	.09
Zn	$ZnSO_4 \cdot 7H_2O$	.1
Mn	$MnCl_2 \cdot 4H_2O$	.7

The solutions were added to saturation (≃ 50 ml) to each cavity at 2, 5, 8, 11, 14, and 17 weeks after germination of seedlings. The treatments were completely randomized on the greenhouse bench with each treatment replicated 10 times; each replicate was a box of 15 seedlings.

After 20 weeks, five randomly selected seedlings were removed from each box and the growing medium was gently washed from the roots.

Table 2.—Treatments for fertility study on northern red oak grown in containers, by nutrient concentration

Treatment	Concentration		
	N	P	K
	.....μg/ml.....		
MNS[1]			
1	12.5 (N_1)	5 (P_1)	40
2	25 (N_2)	5 (P_1)	40
3	50 (N_4)	20 (P_4)	40
4	100 (N_8)	20 (P_4)	40
Rapidgro®	500	500	500

[1]MNS—Modified nutrient solution with various levels of NPK.

Height, root-collar diameter, and fresh top and root weight were determined for each seedling. The roots were visually assessed for degree of ectomycorrhizal development.[2]

Tops and roots of 10 seedlings per replicate were ovendried at 80°C for 60 hours and weighed. Leaves were removed and analyzed for total N by the Kjeldahl method, and for the other elements by dry-ash methods (Wells and others 1973).[3]

RESULTS

Seedlings receiving Rapidgro® produced greater height and root-collar diameter than those receiving the modified nutrient solutions (fig. 1). There was no difference in the two parameters among seedlings receiving the modified nutrient solutions. Top dry weight followed the same pattern between treatments with seedlings receiving the Rapidgro® solution producing four times more top dry weight than those receiving other nutrient solutions (fig. 2). Seedlings fertilized with the Rapidgro® solution also had the greatest amount of root dry weight. Seedlings receiving N_8P_4 nutrient solution had significantly greater root weight than those receiving treatment 1 or Rapidgro®.

[2]Ectomycorrhizal assessments were made by Donald H. Marx, Institute for Mycorrhizal Research and Development, Forestry Sciences Laboratory, Athens, Georgia.

[3]Analyses of foliar samples were performed by Carol G. Wells, USDA Forest Service, Forestry Sciences Laboratory, Research Triangle Park, N.C.



Figure 1.—Shoot height and root-collar diameter of northern red oak seedlings at 15 weeks under five fertility regimes. Bars or circles bearing the same letter are not significantly different (P = 0.05).



Figure 2.—Shoot and root dry weight under five fertility regimes. Bars bearing the same letter are not significantly different (P = 0.05).

The percentage of N in the foliage related well with the increasing amounts of N in the nutrient solutions (table 3). The level of N in the foliage of seedlings receiving Rapidgro® was quite high, probably approaching the luxuriant consumption range. The level of foliar P was slightly affected by fertility regime but levels of K, Ca, Mg, and Fe were not.

The highest percentage of ectomycorrhizae developed on seedlings in N_4P_4 and N_8P_4 treatments (table 4); significantly less formed on seedlings receiving the highest levels of N and P (Rapidgro®). These findings agree with those of

Table 3.—Foliar analysis of 15-week-old northern red oak grown under five fertility regimes in containers,[1] by nutrient

Treatment[2]	N	P	K	Ca	Mg	Fe
	.....	.....	.....	*Percent*	.....	.....
N_1P_1	0.89[d3]	0.46[b]	1.18[a]	0.93[a]	0.23[a]	0.01[a]
N_2P_1	.92[cd]	.47[b]	1.25[a]	.91[a]	.22[a]	.01[a]
N_4P_4	1.03[c]	.53[a]	1.23[a]	1.02[a]	.25[a]	.01[a]
N_8P_4	1.17[b]	.44[b]	1.12[a]	1.00[a]	.22[a]	.01[a]
Rapidgro®	2.02[a]	.42[b]	1.00[a]	.92[a]	.25[a]	.01[a]

[1]Values are means of 10 replicates; each replicate was a composite of leaves from 10 seedlings.

[2]Concentrations of nitrogen as μg/ml – N_1 = 12.5, N_2 = 25, N_4 = 50, N_8 = 100; phosphorus as μg/ml – P_1 = 5, P_4 = 20; Rapidgro® – NPK = 500 μg/ml each.

[3]Means within columns not followed by the same letter are significantly different (P = 0.05).

Table 4.—Percentage of seedlings with ectomycorrhizae and degree of ectomycorrhizal development by *Pisolithus tinctorius* (Pt)[1]

Treatment[2]	Seedlings with Pt	Pt ectomycorrhizae
	..... *Percent*	.....
N_1P_1	99	37.0[b3]
N_2P_1	100	40.7[b]
N_4P_4	100	58.6[c]
N_8P_4	100	63.5[c]
Rapidgro®	89	20.7[a]

[1]Values are means of five seedlings from each of 10 replicates.

[2]Concentrations of nitrogen as μg/ml – N_1 = 12.5, N_2 = 25, N_4 = 50, N_8 = 100; phosphorus as μg/ml – P_1 = 5, P_4 = 20; Rapidgro® – NPK = 500 μg/ml each.

[3]Means not followed by the same letter are significantly different (P = 0.05).

Marx and others (1977b) working with *P. tinctorius* on pine seedlings. Naturally occurring ectomycorrhizae (mostly *Thelephora terrestris*) formed approximately 5 to 10 percent of the total mycorrhizal development on all seedlings.

CONCLUSIONS

Northern red oak seedlings will grow to plantable size in containers filled with 730 cm^3 of pine bark-sludge growing medium in 3½ to 4 months when fertilized with Rapidgro® every 3 weeks. However, ectomycorrhizal development by *Pisolithus tinctorius* is less than satisfactory under this fertility regime. More *Pisolithus* ectomycorrhizae developed with a balanced nutrient solution with lower concentrations of N and P. Treatments 3 (N_4P_4) and 4 (N_8P_4) produced seedlings with excellent ectomycorrhizal development under the conditions of this study. Unfortunately, seedlings under these fertility regimes were still slightly chlorotic (2.5 GY/7.5/5/9) compared to the dark-green foliage on plants receiving Rapidgro® (7.5 GY/4.5/5).[4] Seedlings under treatment 3 or 4 also lacked appropriate stem height and root collar diameter. Tinus and McDonald (1979) state that container-grown bur oak (*Quercus macrocarpa* Michx.) should be 20 to 25 cm tall before outplanting. Northern red oak probably should be at least this tall before planting, and possibly up to 35 to 40 cm, the height of seedlings receiving Rapidgro®.

Work is being continued to find the fertility regime that will produce northern red oak in containers having the robust tops and roots of seedlings receiving abundant N and P and the *P. tinctorius* colonization of those receiving less amounts of N and P. The use of sewage sludge also contributes approximately 10 to 20 μg/ml of N and P to the fertility regime when used in the growing medium. A chemical analysis of the growing medium before and after the addition of nutrient solutions needs to be done to define the actual levels of available N and P.

[4]Foliage-color rating with Munsell color charts for plant tissue (hue/value/chroma).

LITERATURE CITED

Berry, Charles R., and Donald H. Marx.
1976. Sewage sludge and *Pisolithus tinctorius* ectomycorrhizae: Their effect on growth of pine seedlings. For. Sci. 22:351–358.

Clark, F. Bryan, and Richard F. Watt.
1971. Silvicultural methods for regenerating oaks. Proc. Oak Symp., p. 37–43. [Morgantown, W. Va., August 1971.] Northeast. For. Exp. Stn., Upper Darby, Pa.

Core, Earl L.
1971. Silvical characteristics of the five upland oaks. Proc. Oak Symp., p. 19–22. [Morgantown, W. Va., August 1971.] Northeast. For. Exp. Stn., Upper Darby, Pa.

Johnson, Paul S.
1974. Containerization of oak seedlings for the oak-hickory region: A progress report. Proc. North Am. Containerized For. Tree Seedling Symp., p. 197–199. [Denver, Colo., August 1974.] Great Plains Agric. Counc. Publ. 68.

Johnson, R. L.
1979. Adequate oak regeneration: A problem without a solution. Proc. 7th Annu. Hardwood Symp., Hardwood Res. Counc., p. 59–65. [Cashiers, N.C., May 1979.] USDA For. Serv., Southeast. Area, State and Priv. For., Atlanta, Ga.

Marx, Donald H.
1979. Synthesis of ectomycorrhizae by different fungi on northern red oak. USDA For. Serv., Res. Note SE-282, 8 p. Southeast. For. Exp. Stn., Asheville, N.C.

Marx, Donald H., and William C. Bryan.
1975. Growth and ectomycorrhizal development of loblolly pine seedlings in fumigated soil infested with the fungal symbiont *Pisolithus tinctorius*. For. Sci. 21:245–254.

Marx, Donald H., William C. Bryan, and Charles E. Cordell.
1977a. Survival and growth of pine seedlings with *Pisolithus* ectomycorrhizae after two years on reforestation sites in North Carolina and Florida. For. Sci. 23:363–373.

Marx, D. H., A. B. Hatch, and Joseph F. Mendicino.
1977b. High soil fertility decreases sucrose content and susceptibility of loblolly pine roots to ectomycorrhizal infection by *Pisolithus tinctorius*. Can. J. Bot. 55:1569–1574.

Ruehle, John L., and Donald H. Marx.
1977. Developing ectomycorrhizae on containerized pine seedlings. USDA For. Serv., Res. Note SE-242, 8 p. Southeast. For. Exp. Stn., Asheville, N.C.

Ruehle, John L., and Donald H. Marx.
1979. Fiber, food, fuel, and fungal symbionts. Science 206:419–422.

Russell, Thomas E.
1971. Seeding and planting upland oaks. Proc. Oak Symp., p. 49–54. [Morgantown, W. Va., August 1971.] Northeast. For. Exp. Stn., Upper Darby, Pa.

Tinus, Richard W., and Stephen E. McDonald.
1979. How to grow tree seedlings in containers in greenhouses. USDA For. Serv., Gen. Tech. Rep. RM-60, 256 p. Rocky Mt. For. and Range Exp. Stn., Fort Collins, Colo.

Wells, Carol G., D. M. Crutchfield, N. M. Berenyi, and Charles B. Davey.
1973. Soil and foliar guidelines for phosphorus fertilization of loblolly pine. USDA For. Serv., Res. Note SE-110, 15 p. Southeast. For. Exp. Stn., Asheville, N.C.

Wendel, George W.
1979. Growth and survival of planted northern red oak seedlings in West Virginia. USDA For. Serv., Tree Plant. Notes 30:16–19.

The use of trade, firm, or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture or the Forest Service of any product or service to the exclusion of others that may be suitable.

This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife--if they are not handled or applied properly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.

The Forest Service, U.S. Department of Agriculture, is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives—as directed by Congress—to provide increasingly greater service to a growing Nation.

USDA policy does not permit discrimination because of race, color, national origin, sex or religion. Any person who believes he or she has been discriminated against in any USDA-related activity should write immediately to the Secretary of Agriculture, Washington, D.C. 20250.