



Growth and Seed Production of Sawtooth Oak (*Quercus acutissima*) 22 Years After Direct Seeding

J.C.G. Goelz and D.W. Carlson

SUMMARY

Sawtooth oak (*Quercus acutissima* Carruth.) was direct seeded at two locations, one with a poorly drained clay soil and the other with a well-drained silty clay loam. For comparison, Nuttall oak (*Q. nuttallii* Palmer) was direct seeded on the poorly drained clay soil. On the well-drained silty clay loam, sawtooth oak was 18 ft taller and 2.4 inches larger in d.b.h. at age 22 than on the poorly drained clay soil. On the clay, sawtooth oak grew faster than Nuttall oak, but survival was lower. Almost all sawtooth oaks were producing acorns on both sites; however, no developing acorns were found on the Nuttall oaks. Sawtooth oak is a viable alternative for planting on a wide range of Mississippi Delta forest types. Because sawtooth oak has a poorer form than Nuttall oak, its primary use is as a source of wildlife food. Sawtooth oak could be included in plantings with the multiple objectives of timber production and wildlife use because it grows well and could potentially be used for pulpwood.

Keywords: Acorn, Mississippi Delta, Nuttall oak, plantation, wildlife.

INTRODUCTION

Sawtooth oak was introduced into the United States in 1862 from its natural range in eastern Asia. It has been widely planted in the United States as a food tree for wildlife because it produces acorns as early as the fifth or sixth growing season and produces seed crops almost every year thereafter (Mercer 1969, Olson 1974). It has also been used to reclaim spoil banks (McMinn and Crane 1984). Most plantings are descended from trees grown on the University of Georgia campus, Athens, GA, and at the National Plant Introduction Station, Savannah, GA (USDA SCS 1965).

Sawtooth oak has been successfully planted on upland areas of the Coastal Plain, Piedmont, and Appalachian regions (Schoenike 1971, Sullivan and Young 1961). This study was begun in 1970 to describe the development and seed production of sawtooth oak on bottomland sites in the Mississippi Delta. Francis and Johnson (1985) reported results of this study after 14 growing seasons.

METHODS

Two sites with different soil characteristics were selected and direct seeded. The Huntington Point site, in Bolivar County, Mississippi, has a Commerce-Convent soil association (fine-coarse-silty, mixed, thermic Aeric Fluvaquents) with a silty clay loam surface layer underlain by fine sandy loam. The soil has a pH of 7.4. Southern bottomland oak species are generally unsuitable for soils with a pH in excess of 7.5 (Baker and Broadfoot 1979, Broadfoot 1976). In rare years when the Mississippi River overflows its levee, the site floods to a depth

up to 6 ft for as long as 2.5 months; however, the soil is well drained when not flooded. The Delta Experimental Forest site in Washington County, Mississippi, has a Sharkey-Alligator soil association (very fine, montmorillonitic, acid, thermic Vertic Haplaquepts) with clayey texture throughout and a pH of 5.2. Although the area rarely floods from overflow water, this soil is generally saturated during the winter and spring and often possesses a perched water table during the dormant season because water infiltration from winter rain is slow.

Both sites were within larger areas cleared for plantation establishment of other species at the same time and were mechanically cleared and disked before seeding. During April of 1970, two to four acorns were planted 1 to 2 inches deep in spots spaced 10 ft by 10 ft. The Huntington Point site had 80 seed spots of sawtooth oak, whereas the Delta Experimental Forest site had 96 seed spots each of sawtooth and Nuttall oaks in alternating rows. The seed spots were hoed twice during the first year, and the sites were mowed between the rows in the second and third years. Because the sawtooth oak alternated with Nuttall oak and the Nuttall oak was smaller, the sawtooth oak measurements represent sizes that are probably larger than those occurring in a monospecific plantation of sawtooth oak.

During the winter after the 22d growing season, the sites were remeasured for diameter at breast height

(d.b.h.) with a diameter tape, and heights were taken with a clinometer. If more than one tree occupied a seed spot, the largest tree diameter was measured. Acorn production was determined in July 1992 by using binoculars to observe the presence of developing acorns. Diameters and heights of the Nuttall and sawtooth oaks on the Delta Experimental Forest site and the sawtooth oak at Huntington Point were compared using a t-test. Survival of Nuttall and sawtooth oaks on the Delta Experimental Forest site was compared by calculating χ^2 from a 2x2 contingency table.

RESULTS AND DISCUSSION

Heights and diameters were significantly different ($p < 0.001$) between species at the Delta Experimental Forest site and between locations for sawtooth oak. Average tree height, diameter, growth, and survival and acorn production percentages are presented in table 1. There are moderate differences in average diameter and height between sawtooth oak and Nuttall oak at the Delta Experimental Forest site. The early differences noted by Francis and Johnson (1985) have been maintained or have increased over the most recent 8-year period. Recent average height growth has been the same for the

Table 1.— Comparison of average height, diameter, growth, survival, and acorn production at 22 years after planting for sawtooth oak on two sites and Nuttall oak on one site

Item for Comparison*	Sawtooth oak		Nuttall oak
	Huntington Point	Delta Experimental Forest	Delta Experimental Forest
Height (ft)	66	48	43
Diameter (inches)	8.7	6.3	5.1
Height growth (ft)	32	20	20
Diameter growth (inches)	3.6	2.3	2.1
Survival (percent)	53	60	70
Acorn production (percent)	95	96	0

*Growth data represent growth over eight growing seasons.

two species, but the difference between average diameters for the two species has increased. Figure 1 is a view of the sawtooth oak plantation at Huntington Point. Figure 2 shows the contrast between sawtooth oaks and Nuttall oaks at the Delta Experimental Forest site.

Sawtooth oak grows much better on the well-drained soil of the Huntington Point site than on the heavy clay soil at the Delta Experimental Forest site (table 1). At Huntington Point, sawtooth oak has grown an average of 4 ft in height per year and 0.45 inch in d.b.h. per year over the last 8 years. Although this represents excellent growth, Shumard oak (*Quercus shumardii* Buckl.) grows as well as or slightly better than sawtooth oak on similar well-drained sites (Kennedy and Krinard 1985). Because sawtooth oak performs similarly to these other oak species on two contrasting sites, it would be an acceptable choice on either heavy clays or well-drained soils.



Figure 2.— *The plantation at Delta Experimental Forest, Mississippi, showing sawtooth oak growing on the left and Nuttall oak, on the right.*



Figure 1.— *The sawtooth oak plantation growing at Huntington Point, Mississippi.*

Survival of sawtooth oak was less than that of Nuttall oak at the Delta Experimental Forest site (table 1, $p < 0.05$). Direct seeding of Nuttall oak is probably more successful than for any other southern oak; the survival of sawtooth oak is comparable to most other oak species. Direct seeding of sawtooth oak requires that more acorns be planted per acre to achieve stocking equal to that of Nuttall oak. McMinn and Crane (1984) reported 90 percent survival at 5 years for sawtooth oak seedlings planted on a spoil bank. The direct seeded sawtooth oaks may have had greater mortality than those of planted seedlings because of seed mortality or predation or greater amounts of competing vegetation.

Almost all sawtooth oaks at both sites were producing acorns (table 1). Ten percent fewer sawtooth oaks were producing acorns 8 years ago (Francis and Johnson 1985). None of the Nuttall oaks bore developing acorns (table 1). Although Olson (1974) suggests that Nuttall oak may produce acorns as early as age 5, no acorns were observed at age 14 or 22 at the Delta Experimental Forest site. Although no data were recorded for quantity or quality of acorn production, many of the sawtooth oaks possessed large developing crops (fig. 3), and there were numerous sawtooth oak seedlings growing in the vicinity of the plantations.



Figure 3.— A branch of a sawtooth oak showing developing acorns. The picture was taken in July.

The d.b.h. (fig. 4a) and height (fig. 4b) distributions of sawtooth oak were much wider than those for Nuttall oak. Several sawtooth oaks had died back at some time during stand development. Other sawtooth oaks were apparently poor phenotypes because they were much

smaller than neighboring trees, although there was no evidence of dieback. The greater variability of sawtooth oak may indicate an opportunity to thin these stands. Thinning from below would leave superior, faster growing trees, shorten rotations, and, presumably, increase seed production.




Figure 4.— Distributions of sawtooth oak by, a, diameter at breast height (d.b.h.); b, height; Delta Experimental Forest (DEF), Mississippi site, and Huntington Point (HP), Mississippi site and of Nuttall Oak on the DEF site.



Figure 5.— *The branches of sawtooth oak as seen here are persistent and relatively large.*

Although sawtooth oak growth and seed production are greater than those of Nuttall oak, stem quality is poorer (fig. 5). Lower limbs are retained longer than those of Nuttall oak, and stem crook is more common. Sawtooth oak has rougher bark than Nuttall oak, and some stem crook or top dieback may be due to greater loads of vines. Sawtooth oak may not be suitable for quality sawlog production without pruning and vine control. However, sawtooth oak is planted primarily as a food tree for wildlife, and thus log quality may be less important than for other oak species.

When timber production is one of the objectives of management, sawtooth oak may be suitably grown in combination with other, later seeding oaks. Sawtooth oak would provide early acorn production for wildlife and later could be discriminated against while marking for pulpwood thinnings to provide growing space for the more commercially valuable, later seeding oaks. If sawtooth oak was seeded with another species of similar growth rate, intimate mixtures or alternating rows could be used. If either species has a much greater growth rate, then at least the slower growing species should be seeded in blocks of three or more successive rows to lessen inter-specific competition.

LITERATURE CITED

- Baker, J.B.; Broadfoot, W.M. 1979. A practical field method of site evaluation for commercially important southern hardwoods. Gen. Tech. Rep. SO-26. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 51 p.
- Broadfoot, W.M. 1976. Hardwood suitability for and properties of important Midsouth soils. Res. Pap. SO-127. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 84 p.
- Francis, J.R.; Johnson, R.L. 1985. Direct-seeded sawtooth oaks (*Quercus acutissima* Carruth.) show rapid growth on diverse sites. *Tree Planters' Notes*. 36(3): 3-5.
- Kennedy, H.E., Jr.; Krinard, R.M. 1985. Shumard oaks successfully planted on high pH soils. Res. Note SO-321. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 3 p.
- McMinn, J.W.; Crane, W.H. 1984. Five year performance of selected woody species on an upper Coastal Plain spoil bank. *Southern Journal of Applied Forestry*. 8: 207-209.
- Mercer, J., Jr. 1969. Sawtooth oak holds promise as wildlife plant for the Southeast. *Soil Conservation*. 34: 178.
- Olson, D.F., Jr. 1974. *Quercus* L. Oak. In: Schopmeyer, C.S., tech. coord.. *Seeds of woody plants in the United States*. Agric. Handb. 450. Washington, DC: U.S. Department of Agriculture, Forest Service: 692-703.
- Schoenike, R.E. 1971. Sawtooth oak—promising exotic tree species for the Piedmont. *For. Bull.* 6. Clemson, SC: Clemson University, Department of Forestry. 2 p.
- Sullivan, E.G.; Young, W.C. 1961. An exotic oak *Quercus acutissima* for wildlife food planting. In: Conner, E., Jr., ed. *Proceedings of the 15th annual conference of the Southwest Association of Game and Fish Commissions; 1961 October 22-25; Nashville, TN*. Nashville, TN: Southwest Association of Game and Fish Commissions: 136-141.
- U.S. Department of Agriculture, Soil Conservation Service, 1965. *Planting guide for sawtooth oak (Quercus acutissima Carruthers)*. 4-L-20355. Washington, DC: U.S. Department of Agriculture, Soil Conservation Service.

This publication reports research involving pesticides. It does not contain recommendations for their use, nor does it imply that the uses discussed here have been registered. All uses of pesticides must be registered by appropriate State and/or Federal agencies before they can be recommended.

CAUTION: Pesticides can be injurious to humans, domestic animals, desirable plants, and fish or other wildlife—if they are not handled or applied properly. Use all herbicides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and their containers.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at (202) 720-2600 (Voice and TDD).

To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, DC 20250, or call 1-800-245-6340 (voice) or (202) 720-1127 (TDD). USDA is an equal employment opportunity employer.