

## INTERACTION OF FLAVANOIDS WITH PEPTIDES AND PROTEINS AND CONFORMATIONS OF DIMERIC FLAVANOIDS IN SOLUTION

Tsutomu **Hatano**,<sup>a</sup> Takashi **Yoshida**,<sup>a</sup>  
and Richard W. Hemingway\*

<sup>a</sup> Faculty of Pharmaceutical Sciences  
Okayama University  
**Tsushima**, Okayama 700-8530  
JAPAN

\*Southern Research Station  
USDA Forest Service  
Pineville, Louisiana 71360  
USA

### 1. INTRODUCTION

Although the physiological roles of tannins and related polyphenols in plants have not yet been clarified, their ability to form complexes with proteins or related biopolymers has been correlated with some protection of the plants from predators such as animals, insects, and microbes.<sup>1</sup> Similarly, commercial uses of tannins, especially in the leather and brewing industries, are also based on their binding with **proteins**.<sup>2</sup> Pharmacological properties of tannins have been investigated based on recent advances in the structural study of tannins in medicinal plants, and various actions of tannins including antitumor and antiviral effects have been **revealed**.<sup>3</sup> These effects are attributed to interactions with certain biomolecules in organisms, too.


Most of these effects are dependent on the chemical structures or molecular shapes of **tannins**.<sup>3</sup> Therefore, clarification of the molecular conformations of tannins and related polyphenols is requisite to understanding the process of molecular interaction of tannins with the biomolecules. Here we summarize our reports on the conformational analyses of **dimeric** flavanoids related to condensed tannins and interaction of these flavanoids **with** peptides. Because molecular



conformations of oligomeric **flavanoids**<sup>4</sup> in the free phenolic form in solution with the biologically significant solvent (water) had not yet been characterized, we focused our efforts on definition of the conformations of the natural free phenols in aqueous solutions.

## 2. CONFORMATIONS IN ORGANIC SOLVENTS

Among various proanthocyanidins, we used **catechin-(4 $\alpha$ →8)-catechin** and **catechin-(4 $\alpha$ →8)-epicatechin** for the conformational analysis, since the presence of high rotational barriers around their interflavan **linkages**<sup>5</sup> resulted in sharp NMR spectra for the two conformers seen for each of the two compounds. However, complete assignment of these spectra had not been attempted. These **dimeric pro-cyanidins** were synthesized as shown in figure 1.<sup>6</sup> The **2 $\alpha$ ,3 $\beta$ -dihydroquercetin**, prepared by Karchesy at Oregon State University, was treated with sodium **borohydride** and the resulting hydroxyl group at C-4 of the reduction product was removed in acidic conditions to give a cation. Nucleophilic A-rings of either **(+)-catechin** or **(-)-epicatechin** then attacked the cation to form the **dimeric pro-cyanidins**. These two dimers, which have **(+)-catechin** as the upper unit and either **(+)-catechin** or **(-)-epicatechin** as the lower unit, were then available without structural ambiguity.



**Figure 1.** Syntheses of **catechin-(4 $\alpha$ →8)-catechin** and **catechin-(4 $\alpha$ →8)-epicatechin**.

The  $^1\text{H}$  NMR spectrum of **catechin-(4 $\alpha$ →8)-epicatechin** in acetone-*c*6 is shown in figure 2. Steric hindrance minimizes rotation around the **interflavan** bond, and two relatively stable conformational forms were observed in the NMR spectra. An expanded spectrum shows the signals in the aromatic region at around **6 ppm** are A-ring and D-ring protons. Signals at lower field of the aromatic region are composed of four sets of three protons each forming the ABC **3-spin** system characteristic of catechol rings. These are attributable to protons of B-ring and E-ring of this **dimeric** procyanidin. Therefore, the aromatic protons indicate the presence of two conformers, one of which shows signals due to the minor conformer with significant **upfield** shifts relative to those of the major conformer. The **upfield** shift of these B- and E-ring protons indicates that the minor conformer adopts a conformation where the two aromatic rings are close to each other enabling these anisotropic effects.

To establish the spatial positions of the constituent flavan units around the interflavan bonds of the **dimeric** procyanidins, definition of the conformations of heterocyclic C- and F-rings that affect locations of B- and E-rings in each dimer is required. Coupling constants of heterocyclic C- and F-ring protons in the  $^1\text{H}$  NMR spectra of the two dimers, **catechin-(4 $\alpha$ →8)-catechin** and **catechin-(4 $\alpha$ →8)-catechin**, were analyzed with the aid of spectral simulation. The coupling constants for the upper units of both dimers were similar to those for the half-chair form of the heterocyclic ring of the monomeric catechin in which the B-ring is equatorial. However, the coupling constants for the lower units are different from the constituent monomers.

For the lower residue of **catechin-(4 $\alpha$ →8)-epicatechin**, estimation of the  $J_{2,3}$  coupling constant is difficult, since H-2 appears as a broad singlet and H-3 proton





Figure 2.  $^1\text{H}$  NMR spectrum of **catechin-(4 $\alpha$ →8)-epicatechin**.




**Figure 3.** Conformations and coupling patterns for F-ring H-3 of catechin-(4 $\alpha$ →8)-epicatechin in acetone- $d_6$  containing  $D_2O$ .

shows a multiplet signal due to coupling with the two H-4 protons as shown in figure 3. However, spectral simulation using PCPMR<sup>7</sup> led to the assignment that the coupling constant between H-2 and H-3 is 1.2 Hz, and  $J_{3,4a}$  and  $J_{3,4b}$  are 4.2 and 2.4, respectively, when recorded in acetone- $d_6$ . These F-ring coupling constants are consistent with neither the half-chair conformation with an equatorial phenyl ring nor the reverse half-chair conformation with an axial phenyl ring. An energetically unfavorable skewed-boat conformation was thus suggested for the F-ring, even though the observed coupling may be the result of an “averaged” coupling over the time frame of the NMR experiment\* for reasons described below.

In  $D_2O$ , catechin-(4 $\alpha$ →8)-epicatechin showed a coupling pattern (fig. 4) somewhat different from that in acetone- $d_6$  as shown in figure 3. Obviously, the observed signal pattern of F-ring H-3 is far from that for the energetically preferable half-chair or reverse half-chair conformation, and the simulated signal patterns indicated that the F-ring adopts the C3 sofa conformation.


On the other hand, the dimer catechin-(4 $\alpha$ →8)-catechin showed a signal pattern of F-ring H-3 as shown in figure 5. The observed signal pattern is closer to the calculated signal pattern for the half-chair heterocyclic ring with an equatorial phenyl group. The calculated signal pattern for the reverse chair form with an axial phenyl group is far from the measured one, indicating that this form does not participate strongly in the conformation of this compound.



**Figure 4.** Conformations and coupling patterns for F-ring H-3 of catechin-(4 $\alpha$ →8)-epicatechin in D<sub>2</sub>O.

In addition, the molecular shapes of proanthocyanidins are strongly affected by steric hindrance around their interflavan bonds, so it is necessary to define the dihedral angles between the two flavan units of **dimeric** procyanidins. Spatial positions of the two halves in each of the two dimers were analyzed based on <sup>1</sup>H-<sup>1</sup>H long-range COSY<sup>9</sup> and NOESY measurements.


The <sup>1</sup>H-<sup>1</sup>H long-range COSY spectrum of catechin-(4 $\alpha$ →8)-epicatechin shown in figure 6 enabled the assignments of the B- and E-ring protons to each conformer by cross peaks due to the coupling between C-ring H-2 and B-ring H-2' and H-6', and between F-ring H-2 and E ring H-2' and H-6'. In addition, this measurement permitted assignment of the orientation of D-ring relative to the upper half of the dimer in the following way. The <sup>1</sup>H-<sup>1</sup>H long-range COSY spectrum showed strong cross peaks between C-ring H-4 and H-6 and H-8 of A-ring where the A-ring plane is at 90 degrees to the C4-H4 bond. The cross peak between D-ring H-6 and C-ring H-4 of the major conformer therefore indicates that the bond C4-H4 and the D-ring plane form an angle of about 90 degrees. On the other hand, the minor conformer showed a much weaker cross peak for the coupling between the C-ring H-4 and D-ring H-6. Therefore, the dihedral angle between C4-H4 and the D-ring plane in the minor conformer is larger or smaller than 90 degrees tending to either 180 or 0 degrees when the cross' peak is absent.



**Figure 5.** Conformations and coupling patterns for F-ring H-3 of **catechin-(4 $\alpha$ →8)-catechin** in  $D_2O$ .

In support of the above conclusions, the NOESY spectrum of **catechin-(4 $\alpha$ →8)-epicatechin** shown in figure 7 indicates that the C- and E-rings of this molecule are spatially close to each other. A cross peak due to NOE correlation between C-ring H-4 and E-ring H-2' of the major conformer indicated that the major conformer of this **dimeric** procyanidin adopts a conformation in which the  $\alpha$ -oriented E-ring of the lower unit locates near the  $\beta$ -oriented H-4 of the upper unit. Although the spectrum also showed many correlations between the protons of one conformer and the corresponding protons of the other conformer, these cross peaks are not due to NOE correlations but to the conformational exchange. For example, the H-2' of the lower half of one conformer shows a correlation with H-2' of the lower half of another conformer.

The presence of the NOE between C-ring H-4 and E-ring H-2' in the molecule of **catechin-(4 $\alpha$ →8)-epicatechin** was confirmed by measurement of a NOE difference spectrum (fig. 8). Irradiation of H-2' of the lower residue of the major conformer caused increase of the peak area of C-ring H-4 along with that of F-ring H-2. Therefore, the orientation of the lower residue is that shown in figure 7, where the E-ring of the lower residue locates on the same side as that of H-4 of the upper residue. Irradiation of H-2' of the lower unit of the major conformer also caused a negative peak for the corresponding proton of the minor conformer. The appearance of the negative peak of the corresponding proton of another conformer is explained by conformational exchange accompanied by saturation transfer.




**Figure 6.**  $^1\text{H}$ - $^1\text{H}$  Long-range COSY spectrum of **catechin-(4 $\alpha$ →8)-epicatechin** in **acetone- $d_6$**  containing **D $_2$ O**.

Analogous spatial correlations between the protons of C- and E-rings in one conformer of the **dimeric** procyanidin, **catechin-(4 $\alpha$ →8)-catechin** were also observed in the NOESY and NOE difference spectra.

### 3. CONFORMATIONS IN WATER


The  $^1\text{H}$  NMR spectra of **catechin-(4 $\alpha$ →8)-epicatechin** in three different solvents, acetone- $d_6$ , **dioxane- $d_8$**  and **D $_2$ O**, differed as shown in figure 9. The chemical shifts of several aromatic protons in the spectrum in **dioxane- $d_8$**  varied from those in the spectrum in acetone- $d_6$ , whereas relative abundances of the two conformers in



**Figure 7.** NOESY Spectrum of catechin-(4 $\alpha$ →8)-epicatechin in acetone & containing D<sub>2</sub>O.

these two solvents were almost the same. In D<sub>2</sub>O, the signals of the major conformer in the organic solvents almost disappeared, and only the signals of another conformer were clearly observed. The aromatic protons of the major conformer in D<sub>2</sub>O (i.e., the minor conformer in the organic solvents) shifted upfield noticeably relative to those of the other conformer. These changes are attributable to the anisotropic effects between B- and E-rings where the two aromatic rings overlap each other.

The composition of the two conformers of catechin-(4 $\alpha$ →8)-epicatechin was dependent on the water content in the organic solvent. As shown in figure 10, the abundance of the major conformer in the organic solvents increases in the presence of a small amount of D<sub>2</sub>O and then rapidly decreases upon further addition of D<sub>2</sub>O. The relative proportion of the two conformers was estimated from the peak area in the <sup>1</sup>H NMR spectra. This change in rotamer population can be explained in the following way. The more "extended" conformer (i.e., the major conformer in




**Figure 8.** NOE difference spectrum of catechin-(4 $\alpha$ →8)-epicatechin in acetone- & containing D<sub>2</sub>O.

acetone-*d*<sub>6</sub>) is considered to be energetically preferable in the organic solvents. On the other hand, in D<sub>2</sub>O, the dimer prefers the more “compact” conformation that is stabilized by hydrophobic or  $\pi$ - $\pi$  interactions. These two conformations, which have been discussed for catechin-(4 $\alpha$ →8)-epicatechin, are represented by those shown in figure 11.<sup>10</sup>

Interestingly, for catechin-(4 $\alpha$ →8)-catechin, the relative abundance of the more “extended” conformer where the C- and E-rings are close to each other is smaller than that of the more “compact” conformer even in organic solvents such as acetone-*d*<sub>6</sub> and dioxane-*d*<sub>8</sub>. In D<sub>2</sub>O, this dimer exclusively adopts the more “compact” conformation.


#### 4. INTERACTIONS BETWEEN POLYPHENOLS AND PEPTIDES CONTAINING PROLINE RESIDUES

Utilization of tannins for leather-tanning is based on the affinity of tannins for collagen, which is rich in proline and hydroxyproline residues. The abundant prolyl residues in salivary proteins also are considered to contribute substantially




**Figure 9.** Solvent dependent changes of the  $^1\text{H}$  NMR spectrum of catechin-(4 $\alpha$ →8)-epicatechin.

to preferential complexation with tannins resulting in the astringency so important to the flavor of fine red wines.<sup>2,11</sup> The benefits of traditional uses of medicinal plants containing high levels of tannins on such problems as skin diseases or digestive disorders may also be attributable to the interaction of tannins with **proline**-rich proteins. Participation of **proline** residues in the complexation between pentagalloylglucose and proline-rich **peptides** has been shown.<sup>12</sup>




**Figure 10.** Effects of D<sub>2</sub>O content on relative abundances of two conformers of catechin-(4α→8)-epicatechin.



**Figure 11.** Conformations of catechin-(4α→8)-epicatechin in solutions a) More "extended", major conformation in acetone-d<sub>6</sub> and dioxane-d<sub>8</sub>. b) More "compact", minor conformation in D<sub>2</sub>O.

Our studies started with the analysis of the simplest combination, catechin and proline (or hydroxyproline), using NOESY experiments as a tool for clarifying the positions in each of the two molecules involved in molecular interaction. Catechin has been regarded as too small to interact to form precipitates with proteins.<sup>13</sup> However, when even a small amount of catechin was added to an aqueous solution (containing 10 percent of CD<sub>3</sub>OD) of poly-L-proline (m.w. 10,000–30,000)

(9 mM for catechin and 120 mM for a monomeric **proline** residue), a white precipitate was formed. The interaction of catechin with a local portion of the **polypeptide** chain was proposed as the mechanism of this interaction." NMR experiments on mixtures of catechin and **proline** in  $\text{H}_2\text{O}$  (containing 10 percent  $\text{CD}_3\text{OD}$ ) with the NOESYHG pulse sequence' showed cross peaks of intermolecular **NOEs** between catechin B-ring protons and the **proline**  $\text{C}_\gamma$  protons (fig. 12). The  $\text{C}_\beta$  and  $\text{C}_\delta$  protons of **proline** showed weaker interactions with aromatic protons of both the A- and B-rings of catechin. On the other hand, the NOESY spectrum of a mixture of catechin and **cis-4-hydroxy-L-proline** showed significant cross peaks between the  $\text{C}_\delta$  protons of hydroxyproline and B-ring protons of catechin. The spectrum also showed correlations between a hydroxyproline  $\text{C}_\delta$  proton ( $\alpha$ -oriented




**Figure 12.** Association of (+)-catechin with **proline**, hydroxyproline, and **dipeptides** containing **proline**. The arrows indicate the sites that showed NOE interaction with **catechin** protons.

proton) to catechin H-6 and H-8. Importantly, noticeable changes in the  $^1\text{H}$  or  $^{13}\text{C}$  chemical shifts, which would suggest the presence of strong hydrogen bonding effects, were not observed. Therefore, hydrophobic interaction seems to be the dominant feature of these intermolecular associations. The aromatic rings of catechin and the aliphatic regions of **proline** or hydroxyproline come in close contact, but the complex is not so tight that we see significant changes in chemical shifts.

As mentioned above, polyphenols are believed to preferentially interact with **proline** residues in peptides. However, catechin showed intermolecular NOEs with the other amino acid residues rather than **proline** when dipeptides containing a **proline** residue, shown in figure 12, were used as targets for intermolecular recognition. An aqueous solution (containing 10 percent  $\text{CD}_3\text{OD}$ ) of a mixture of **Pro-Gly** and (+)-**catechin** showed intermolecular NOE between the glycine  $\text{C}_\alpha$  protons and the (+)-**catechin** H-8 proton. **Pro-Val** showed intermolecular NOE with **catechin** between methyl signals of the valine residue and the catechin H-8 as well as the H-2' proton. **Pro-Phe** showed intermolecular NOE with catechin between the phenyl ring of the **peptide** and the catechin B-ring. **Gly-Pro** showed two sets of  $^1\text{H}$  and  $^{13}\text{C}$  signals, which are attributable to the isomers concerning the amino (imino) nitrogen of **proline**. The major isomer, which showed a 2H singlet-like signal for glycine  $\text{C}_\alpha$  protons, was assigned the trans-configuration. The minor isomer, which showed two doublets of the corresponding protons whose chemical shifts are separated due to the effect of the neighboring **proline** carbonyl carbon, was assigned the cis-configuration. The NOESY spectrum of a mixture of catechin and **Gly-Pro** showed significant cross peaks between the glycine protons of the cis-isomer and the catechin H-6 and H-8. On the other hand, the *trans-isomer* showed only very weak cross peaks with the B-ring protons of catechin. These results indicate that two aromatic rings of the catechin molecule preferentially associate with hydrophobic moieties in **peptides** other than the **proline** residues. Self-association of (+)-**catechin** in water was also shown by an NOE correlation of H-2' with H-8 of another molecule of catechin, even in competition with these peptides.

NOESY experiments on the interactions of polyphenols with two oligopeptides containing **proline** residues suggested the importance of the molecular conformations of both the polyphenol and **peptide** in addition to the hydrophobicity, to the specificity of their complexation. Bradykinin is a nonapeptide, **Arg-Pro-Pro-Gly-Phe-Ser-Pro-Phe-Arg**, containing three prolyl residues. This **peptide** exhibits various physiological effects including the regulation of blood pressure, and conformational studies on it in various solvents have been conducted. "Complexation of this **peptide** with polyphenols has also been investigated." Our NOESY study on the interaction of bradykinin and (+)-**catechin** showed NOEs of the **proline**  $\text{C}_\delta$  protons, the Ser  $\text{C}_\beta$  methylene protons, and the phenyl protons of the **phenylalanyl** residues with the catechin A-ring protons. The phenylalanyl residue also showed correlations with catechin B-ring protons. The amino acid residues in bradykinin showing these NOEs are contained in the sequence. **Phe<sub>5</sub>-Ser<sub>6</sub>-Pro<sub>7</sub>-Phe<sub>8</sub>**. The interaction of this same sequence in bradykinin with micelles has also been reported.<sup>15</sup>

**Gly-Pro-Gly** is a tetrapeptide with inhibitory effects on the activity of dipeptidyl peptidase IV. This compound has been reported to inhibit the entry of



**Figure 13.** Observed NOE interaction of flavanoids with Gly-Pro-Gly-Gly in water.

HIV into cells.<sup>19</sup> Although the *cis*-isomer of the **proline** nitrogen is present, its content is low, so the mixture can be treated practically as *trans*-isomer. Differentiation of the three sets of methylene protons of the glycine residues in the **peptide** was substantiated by the  $^1\text{H}$ - $^1\text{H}$  and  $^1\text{H}$ - $^{13}\text{C}$  long-range COSY analyses.

The NOESY spectrum of the mixture of (+)-**catechin** and Gly-Pro-Gly-Gly in water (containing 10 percent  $\text{CD}_3\text{OD}$ ) showed correlations of catechin H-6 and H-8 with the  $\text{C}_\alpha$  methylene protons of the C-terminal glycine residue, and of the catechin H-5 proton with methylene protons of the N-terminal glycine residue, suggesting that the catechin A-ring preferentially associates with the C-terminal and the B-ring with the N-terminal glycine residues.<sup>20</sup> Intermolecular NOEs, which suggest participation of the **proline** residue in the complexation, were not observed. **Proline** probably plays a role in the restriction of the relative spatial positions of the two glycine residues.

Intermolecular **NOEs** were also observed for the combination of Gly-Pro-Gly-Gly and **catechin-(4 $\alpha$ →8)-catechin**. The association sites in the molecules of the polyphenol and **peptide** are, however, different from those observed in catechin. The NOESY spectrum showed intermolecular NOE between **H-8<sub>A</sub>** (the upper unit of the procyanidin dimer) and the methylene protons of the N-terminal glycine residue and between the E-ring protons, **H-5<sub>E</sub>** and **H-2<sub>E</sub>** of the dimer with the amide proton of the C-terminal glycine residue. These cross peaks suggest an association as exemplified by figure 13.

## 5. CD SPECTRAL ANALYSIS OF POLYPEPTIDE CONFORMATION CHANGE<sup>6</sup>

The **NMR** results described above indicated that complexation is directed to conformationally accessible hydrophobic regions; therefore, the molecular shapes


of both the polyphenol and polypeptide are important to selectivity. The NMR measurements of mixtures of flavans and bradykinin did not show significant changes in the chemical shifts of proton signals. This fact suggested interaction without conformational change of either the polyphenol or the peptide. Because circular dichroism (CD) is known to be sensitive to, the conformational changes of peptides," we used CD to verify whether the conformational changes in the peptide molecule occur with addition of polyphenols or not. When bradykinin was used as the peptide for the complexation, CD spectral measurements did not show any significant spectral changes upon the addition of either catechin or catechin-(4 $\alpha$ →8)-catechin.

However, the CD spectrum of cytochrome c showed a noticeable change in the visible region when either catechin or catechin-(4 $\alpha$ →8)-catechin was added to the solution of cytochrome c (fig. 14). This CD spectral change can, however, be attributed to the reduction of Fe<sup>3+</sup> to Fe<sup>2+</sup> of the heme prosthetic group based on the change in the visible spectrum (fig. 14). Most proteins in aqueous media, including cytochrome c, adopt conformations where the hydrophobic and hydrophilic side chains of the constituent amino acid residues are inside and outside, respectively. The reduction of the heme group by the polyphenols must be explained in terms of the interaction at the hydrophobic "inside" region in the cytochrome c molecule. Further study on the interaction of flavans with cytochrome c is now in progress.

## 6. CONCLUSIONS

Complete assignment of the <sup>1</sup>H and <sup>13</sup>C NMR spectra of the two conformational isomers for each of the dimeric proanthocyanidins catechin-(4 $\alpha$ →8)-catechin and catechin-(4 $\alpha$ →8)-epicatechin, in the natural free phenolic form and in the biologically significant solvent water, has been accomplished." Knowledge of the assignments of the proton spectra, together with the application of various two-dimensional NMR experiments, has permitted the definition of the shapes of the heterocyclic C and F rings in the upper and lower units, the dihedral angles between the upper and lower flavan units, and the effect of changes in solvent composition on the relative proportions and shapes of the two conformers of these dimeric proanthocyanidins in organic and water solvents." In organic solvents, the more extended conformation is favored for catechin-(4 $\alpha$ →8)-epicatechin; and a similar extended conformation, although not the major isomer, is present in nearly equal proportion to a more compact conformer for catechin-(4 $\alpha$ →8)-catechin. However, in water the more compact conformers dominate. These "unfavorable" conformations seem to be stabilized by hydrophobic (or  $\pi$ - $\pi$ ) interactions between the upper and lower unit catechol rings.

The importance of hydrophobic interactions was further highlighted in studies of the interaction of proanthocyanidins with oligopeptides. Even though trimeric and higher molecular weight proanthocyanidins are commonly assumed to be responsible for the precipitation of proteins, when even low mole ratios of the monomeric flavan-3-ol (+)-catechin was combined with polyproline, a white precipitate was obtained.<sup>20</sup> Evidence for hydrophobic interaction was seen in NOESYHG experiments, but no noticeable changes in <sup>1</sup>H or <sup>13</sup>C chemical shifts



**Figure 14.** Spectral changes of cytochrome c upon the addition of flavans a) CD spectra (in  $\text{H}_2\text{O}-\text{CH}_3\text{OH}$ , 9:1, v/v). b) Visible spectra (in  $\text{H}_2\text{O}-\text{CH}_3\text{OH}$ , 9:1, v/v).

were seen that would suggest strong hydrogen bonding in mixtures of (+)-catechin with L-proline or 4-hydroxy-L-proline. Studies of a series of dimeric peptides with various amino acid residues coupled with proline once more highlighted the significance of hydrophobic interaction. In association of catechin with dimeric peptides, stronger cross peaks were seen between either the A- or B-ring protons with the glycine, valine, or phenylalanine protons than to the prolyl unit protons. Once more, no measurable changes in chemical shifts suggesting strong hydrogen

bonding were observed. It is important to note that intermolecular self-association of catechin also resulted in strong cross peaks in these experiments.

Addition of catechin or its dimer to Gly-Pro-Gly-Gly did not show association with the prolyl residue but rather to the conformationally accessible methylenes of **glycine**.<sup>20</sup> No significant changes in NMR chemical shifts or CD spectra that would suggest a "tight" binding or strong hydrogen bonding effects were noted. Studies of the interaction of catechin and **dimeric** procyanidins with bradykinin and cytochrome c also showed no significant change in the conformation of the **peptides**. Rather, association of the polyphenol with the presumed biologically active center of bradykinin and reduction of the **heme** group that resides within the interior of the cytochrome c polymer suggested that hydrophobic interaction dominates in the association of polyphenols with proteins in water. No significant changes in NMR chemical shifts that might suggest strong hydrogen bonding were observed. **The** evidence of all these experiments suggests strong interaction of polyphenols with hydrophobic centers where both self-association and preference for accessible hydrophobic regions of oligomeric **peptides** dominate.

## ACKNOWLEDGMENTS

This work was funded by USDA NRICGP Grant No. 94-03395.

## REFERENCES

1. Schultz, J.C.; Hunter, M.D.; Appel, M. Antimicrobial activity of polyphenols mediates **plant-herbivore** interactions. *In*: Hemingway, R.W., Laks, P.E. (eds.), **Plant polyphenols—synthesis, properties, significance**. Plenum Press, New York, p. 621 (1992).
2. **Haslam**, E. **Practical polyphenolics: from structure to molecular recognition and physiological action**. Cambridge University Press, Cambridge (1998).
3. Okuda, T.; Yoshida, T.; Hatano, T. Hydrolyzable tannins and related polyphenols. *Fortschr. Chem. Org. Naturst.* **66:1** (1995).
4. **Steynberg**, J.P.; Vincent Brandt, E.; Ferreira, D.; Helfer, C.A.; **Mattice**, L.; Gomik, D.; Hemingway, R.W. Conformational analysis of oligomeric flavanoids 2. Methyl ether acetate derivatives of profisetinidins. *Magn. Res. Chem.* **33:611** (1995).
5. **Haslam**, E. Symmetry and promiscuity in proanthocyanidin **biochemistry**. *Phytochemistry* **16:1625**(1977).
6. **Botha**, J.J.; Ferreira, D.; Roux, D.G. Synthesis of condensed tannins. Part 4. A direct **biomimetic** approach to [4,6]- and [4,8]-**biflavonoids**. *J. Chem. Soc. Perkin Trans. 1.* :1235 (1981).
7. PCPMR Spectral Simulation Program, Version 1. Serena Software, Bloomington, IN.
8. **Tobiason**, F.L.; Kelley, S.; Midland, M.M.; Hemingway, R.W. Temperature dependence of (+)-catechin pyran ring proton coupling constants as measured by NMR and modeled using GMMX search methodology. *Tetrahedron Letts.* **38(6):985** (1997).
9. Ziegler, I? A practical guide to 1-D and 2-D experiments for AC/AM Systems, Bruker Instruments, Billerica, MA (1991).
10. Hatano, T.; Hemingway, R.W. Conformational isomerism of phenolic procyanidins: preferred conformations in organic solvents and water. *J. Chem. Soc., Perkin Trans. 2.* :1035 (1997).
11. Hagerman, A.; Butler, L.G. The specificity of proanthocyanidin-protein interactions. *J. Biol. Chem.* **256:4494** (1981).

12. Murray, N.J.; Williamson, M.P.; Lilley, T.H.; **Haslam**, E. Study of the interaction between salivary proline-rich proteins and a polyphenol by  $^1\text{H-NMR}$  spectroscopy. *Eur. J. Biochem.* **219:923** (1994).
13. Bate-Smith, E.C. Haemanalysis of tannins: The concept of relative astringency. *Phytochemistry* **12:907** (1973).
14. Tilstra, L.F.; Cho, D.; Bergmann, W.R.; **Mattice**, W.L. Interaction of condensed tannins with biopolymers. *In*: Hemingway, R.W.; Karchesy, J.J. (eds.). Chemistry and **significance** of condensed tannins. Plenum Press, New York, p. 335 (1989).
15. Denys, L.; Bothner-By, A.A.; Fisher, G.H.; Ryan, J.W. Conformational diversity of bradykinin in aqueous solution. *Biochemistry* **21:6531** (1982).
16. Young, **J.K.**; Hicks, R.P. NMR and molecular modeling investigations of the neuropeptide bradykinin in three different solvent systems: DMSO, **9:1 dioxane/water**, and in the presence of **7.4 mM** lyso phosphatidylcholine micelles. *Biopolymers* **34:611** (1994).
17. Cann, J.R.; Liu, Z.; Stewart, J.M.; Gera, L.; Kotovych, **G.** A CD and an NMR study of multiple bradykinin conformations in aqueous trifluoroethanol solutions. *Biopolymers*. **34:869** (1994).
18. **Haslam**, E. Natural polyphenols (vegetable tannins) as drugs: possible modes of action. *J. Nat. Prod.* **59:205** (1996).
19. Callebaut, C.; **Krust**, B.; Jacotot, E.; Hovamessian, **A.G.** T cell activation antigen, CD26, as a cofactor for entry of HIV in **CD4\*** cells. *Science* **262:2045** (1993).
20. Hatano, T.; Hemingway, R.W. Association of **(+)-catechin** and **catechin-(4 $\alpha$ - $\rightarrow$ 8)-catechin** with oligopeptides containing **proline** residues. *J. Chem. Soc., Chem. Commun.* **:2537** (1996).
21. Das, **T.K.**; Mazumdar, S.; Mitra, S. **Heme** CD as a probe for monitoring local structural changes in **heme-proteins**: alkaline transition in hemeproteins. *Proc. Indian Acad. Sci.* **107:497** (1996).

