



United States
Department of
Agriculture

Forest Service

Southern Forest
Experiment Station

New Orleans,
Louisiana


Proceedings Reprint



SMALL MAMMAL COMMUNITIES OF MATURE PINE HARDWOOD STANDS IN THE OUACHITA MOUNTAINS

Tappe, Philip A.; Thill, Ronald E.; Krystofik, Joseph J.; Heidt,
Gary A.

In: Baker, James B., **comp.** Proceedings of the symposium on
ecosystem management research in the Ouachita Mountains:
pretreatment conditions and preliminary findings; 1993 October
26-27; Hot Springs, AR. Gen. Tech. Rep. SO-112. New
Orleans, LA: U.S. Department of Agriculture, Forest Service,
Southern Forest Experiment Station: 74-81.



Southern
Forest
Experiment
Station

1994



Small Mammal Communities of Mature Pine-Hardwood Stands in the Ouachita Mountains¹

Philip A. Tappe, Ronald E. Thill, Joseph J. Krystofik, and Gary A. Heidt²

ABSTRACT

A study was conducted on the Ouachita and Ozark National Forests in Arkansas to evaluate the effects of alternative pine-hardwood reproduction cutting methods on small mammal abundance and diversity. Pretreatment characteristics of small mammal communities on 20 late-rotation mixed pine-hardwood stands in four physiographic zones of the Ouachita Mountain region of Arkansas are presented. Each physiographic zone (block) contained one replication of five treatments (four future treatments and an untreated control). The most commonly captured small mammal species were *Peromyscus* spp., *Blarina carolinensis*, and *Ochrotomys nuttalli*. Capture success varied between years but most likely reflected changes in probabilities of capture of individual animals and not fluctuations in community composition. Small mammal species richness, diversity, evenness, and relative abundance did not differ between physiographic zones or future treatments.

INTRODUCTION

There is currently a paucity of information concerning the implications of even- and uneven-aged mixed pine-hardwood management for small mammals. In extensive literature searches on the topic of silvicultural effects on wildlife habitat (covering the years 1953 through 1990), Harlow and Van Lear (1981, 1987) and NCASI (1993) did not cite a single paper specifically addressing the effects of reproduction cutting methods in mixed pine-hardwood stands on small mammal communities. Coupled with increasing concerns over the impacts of silvicultural practices on wildlife, research on the effects of alternative silvicultural practices is particularly warranted for small mammal communities. Small mammals are the primary prey base for many mammalian and avian predators. Mycophagous species facilitate dispersal of fungal spores that form root-inhabiting ectomycorrhizae required by most higher plants for adequate nutrient procurement, enhanced water absorption, and protection from root pathogens (Maser and others 1978). Consumption of pine seeds by some species may adversely affect regeneration success (Pank 1974, Smith and Aldous 1947). In addition, fossorial species may significantly influence hydrological processes on forested watersheds (Ursic and Esher 1988).

The objective of this study is to evaluate the effects of alternative pine-hardwood reproduction cutting methods on small mammal abundance and diversity. Pretreatment characteristics of small mammal communities in late-rotation stands in the Ouachita Mountain region of Arkansas are presented in this paper.

METHODS

Study Areas and Treatments

Twenty late-rotation mixed pine-hardwood stands in four physiographic zones of the Ouachita Mountain region were selected for study. These stands are located in the Ouachita National Forest and Ozark-St. Francis National Forest and are characterized by: size ranges from 14.2 to 16.2 ha; predominantly south, southeast, or southwest aspects; and slopes of 5 to 20 percent. Locations and habitat characteristics of the stands are described by Thill and others (1994).

Four replications of five treatments, blocked by physiographic zone, were randomly assigned to 20 stands. These treatments consist of an untreated control and four reproduction cutting methods: clearcut, shelterwood, group selection, and single-tree

¹ Paper presented at the Symposium on Ecosystem Management Research in the Ouachita Mountains: Pretreatment Conditions and Preliminary Findings, Hot Springs, AR, October 26-27, 1993.

² Assistant professor, School of Forest Resources, University of Arkansas at Monticello, Monticello, AR 71656; supervisory research wildlife biologist, Southern Forest Experiment Station, Nacogdoches, TX 75962; research specialist, School of Forest Resources, University of Arkansas at Monticello, Monticello, AR 71656; professor, Department of Biology, University of Arkansas at Little Rock, Little Rock, AR 72204; respectively.

selection. All **silvicultural treatments**, except for the clearcut, provide for the **retention** of **overstory hardwoods**. These treatments were implemented in the **summer** of 1993. For a detailed description of treatments and physiographic **zones**, see **Baker (1994)**.

Small **Mammal** Surveys

Small mammals were trapped prior to treatment installation during December of 1991 and 1992 using Sherman **live**traps (7.62 cm by 8.89 cm by 22.86 cm). These traps are of sufficient **size** to capture eastern **woodrats** (*Neotoma floridana*), **flying squirrels** (*Glaucomys volans*), and smaller mammals. Eighty trap stations per stand were located at 15-m intervals **along permanent** transects established for small mammal trapping, habitat sampling, and biodiversity surveys. To ensure **adequate coverage** of each study area, each stand was subdivided into 50-m wide, parallel bands. One randomly selected transect was **located within** each band so that no transect was closer than 30 m **from** another transect. The number of transects and **their** lengths varied by stand size and shape. To minimize potential edge influences, no traps were placed closer than 50 m from the stand boundaries. A more detailed description of **transect** establishment is given by **Thill** and others (1994). In 1991, one trap **was placed at each** station. Sampling effort was increased to **two traps** per station in 1992 to ensure ample opportunities for **multiple captures** per trap station.

Traps were baited with commercial horse and mule feed and checked for 10 consecutive days. A wad of cotton was placed in each trap for nesting material to minimize **trap** mortality. **Captured** mammals were marked and released at **the** site of capture **after** recording species, sex (when possible), and location (station number) of capture. A tally was also **kept** on the number of empty/sprung traps so that total available trap nights could be computed. Because of insufficient discriminating **physical** characteristics available **from** field observations, animals of the genus *Peromyscus* (including *P. leucopus*, *P. gossypimus*, *P. maniculatus*, and *P. atwateri*) were not identified to **the** species level.

Data **Analyses**

A number of diversity measures were computed for each stand. These included species richness (i.e., the number of species encountered), Shannon's **diversity** index (Shannon and Weaver 1963), and species **evenness** (i.e., the distribution of individuals among species). **In** addition, an index of relative species abundance was obtained by computing **small mammal captures** per 100 trap nights, excluding recaptures and **after** correcting for sprung/empty traps. Though sampling **intensity** was doubled in 1992, very few multiple captures **occurred**. Thus, to ensure that data were comparable between **years**, 1992 **trap** nights were based on the number of trap stations as opposed to the number of individual traps. **Diversity**, evenness, **richness**, and relative abundance of small mammals were compared by year using Witcoxon matched-pairs signed-ranks tests and by physiographic zone and future treatment using one-way analysis of variance (**ANOVA**) and Tukey's HSD. **In** addition, small mammal community composition was evaluated among physiographic zones and among **future** treatments **using** Sorensen's Similarity Index (**Mueller-Dombois** and **Ellenberg** 1974). This index is computed as: $SI = 200C/(A + B)$, where **A** = the number of species in zone or **treatment A**, **B** = the number of species in zone or **treatment B**, and **C** = the number of species that **areas A and B** have in common. Similarity can range **from** 0 percent (no species in common) to 100 percent (identical species composition). So that community composition could be readily compared **between** any two blocks or **treatments**, an index **value** was computed for each combination of both physiographic zones (6 **combinations**) and **future** treatments (10 **combinations**).

RESULTS

After correcting for sprung traps and **differential** sampling intensity, 2 1,165 **trap** nights were **accumulated** over 2 years across **all** stands. A total of 502 small mammals of 10 species were captured (table 1). Trapping success across years was 2.37 new captures per 100 **trap** nights. **The** most commonly **captured** species were *Peromyscus* spp., *Blarina carolinensis*, and *Ochrotomys nuttalli*, comprising 94 percent of all animals **captured** (table 1).

Differences by **Year**

Capture success was **different** between 1991 and 1992. **Total** captures varied significantly between years ($Z = -3.6773$, $P < 0.001$), and numbers of individuals **within** the species were **different** for the three most **commonly** caught **species** (table 1). New captures per **100 trap nights** also **differed between years for the three most commonly captured species** (table 1): *Peromyscus* spp. ($Z = -2.6880$, $P = 0.007$), *Blarina carolinensis* ($Z = -2.7253$, $P = 0.006$), and *Ochrotomys nuttalli* ($Z = -3.0102$, $P = 0.003$).

Because of these differences in capture success, measures of diversity also **differed** between years. Species richness among stands differed by year ($Z = -2.1993$, $P = 0.028$), averaging 3.35 (SE = 0.24) for 1991 and 2.55 (SE = 0.20) for 1992. Species diversity also differed by year ($Z = -2.3146$, $P = 0.021$), averaging 0.90 (SE = 0.08) for 1991 and 0.71 (SE = 0.07) for 1992. However, species evenness did not differ between years ($Z = -0.8213$, $P = 0.412$), averaging 0.73 (SE = 0.06) for 1991 and 0.75 (SE = 0.06) for 1992.

Table 1- Species, numbers of captured individuals, and relative abundance (new captures per 100 trap nights) of small mammals captured by year in Ecosystem Management research stands in the Ouachita Mountains of Arkansas

Species	1991		1992		Total captures	Total captures/100 trap nights
	Captures	Captures/100 trap nights	Captures	Captures/100 trap nights		
<i>Peromyscus</i> spp.*	170	1.62	84	0.79	254	1.20
<i>Blarina carolinensis</i>	127	1.21	31	0.29	158	0.75
<i>Ochrotomys nuttalli</i>	51	0.49	9	0.08	60	0.28
<i>Neotoma floridana</i>	8	0.08	5	0.05	13	0.06
<i>Reithrodontomys fulvescens</i>	5	0.05	1	0.01	6	0.03
<i>Glaucomys volans</i>	3	0.03	2	0.02	5	0.02
<i>Microtus pinetorum</i>	2	0.02	1	0.01	3	0.01
<i>Orzomys palustris</i>	1	0.01	0	0.00	1	co.01
<i>Sigmodon hispidus</i>	1	0.01	0	0.00	1	co.01
<i>Mus musculus</i>	0	0.00	1	0.01	1	co.01
Total	368	3.51	134	1.26	502	2.37

* Include *S. P. leucopus*, *P. gossypinus*, *P. maniculatus*, and *P. attwateri*.

Because habitat characteristics did not change appreciably between years, differences in relative abundance and diversity were probably related to differences in the inherent probabilities of capture of individual animals and not to yearly fluctuations in the actual abundance and/or composition of small mammal species. Species that were captured only 1 year were likely present both years, but their populations may have been so low that capture was unlikely given our sampling intensity. When richness, diversity, and evenness values were computed for pooled 1991 and 1992 data, mean values across all stands were 3.70 (SE = 0.21) for species richness, 0.98 (SE = 0.07) for species diversity, and 0.75 for species evenness.

Differences by Physiographic Zones

Physiographic zones were compared for each year and for pooled data across years. In 1991, 1992, and 1991-92 combined, no differences between zones were found for relative abundance, richness, diversity, or evenness (tables 2, 3). Likewise, few habitat parameters were significantly different by physiographic zone (Thill and others 1994). Small mammal community similarity between physiographic zones ranged from 61.5 percent to 87.5 percent (table 4), and averaged 74.1 percent. However, *Peromyscus* spp., *Blarina carolinensis*, and *Ochrotomys nuttalli* were present in all zones. Because all other species comprised only 6 percent of the animals captured, these areas may be more similar than indicated by Sorensen's index.

Differences by Future Treatments

Groups of stands targeted for future treatments were also compared for each year and for pooled data across years. In 1991 and 1991-92, no differences between future treatments were found for relative abundance, richness, diversity, or evenness (Tables 5 and 6). In 1992, there were no differences between future treatments for relative abundance, richness, or diversity; however, the mean evenness value for the group of stands to receive the shelterwood treatment differed from all other groups of stands except the control group (tables 5, 6). Likewise, only one of 69 habitat parameters (volume of down pine logs, decay class 3) differed significantly among future treatments (Thill and others 1994). Small mammal community similarity between future treatments ranged from 57.1 percent to 92.3 percent (table 4) and averaged 78.5 percent. Similar to physiographic zones, *Peromyscus* spp., *Blarina carolinensis*, and *Ochrotomys nuttalli* were present in all treatment areas. In addition, the next most abundant species, *Neotoma floridana* (table 1), was also present in all treatment areas. Thus, these areas may also be more similar in small mammal composition than indicated by Sorensen's index.

Table 2- Relative abundance (new captures per 100 trap nights) of small mammals and the three most commonly captured species ($\bar{x} \pm SE$) in Ecosystem Management research stands in the Ouachita Mountains of Arkansas by physiographic zone, 1991-92

Species	Year	Zone				F	P*
		North	South	East	West		
All species	1991	3.21 \pm 1.13	3.27 \pm 1.23	4.88 \pm 0.91	2.15 \pm 1.35	0.637	0.60
	1992	1.29 \pm 0.44	1.11 \pm 0.22	1.46 \pm 0.47	1.15 \pm 0.36	0.173	0.91
	Pooled	2.25 \pm 0.66	2.19 \pm 0.69	3.17 \pm 0.75	1.95 \pm 0.71	0.584	0.63
<i>Peromyscus</i> spp.†	1991	2.01 \pm 0.80	1.99 \pm 1.32	1.46 \pm 0.37	0.98 \pm 0.41	0.351	0.79
	1992	0.88 \pm 0.33	0.63 \pm 0.15	0.93 \pm 0.39	0.69 \pm 0.24	0.245	0.86
	Pooled	1.45 \pm 0.45	1.31 \pm 0.67	1.19 \pm 0.27	0.84 \pm 0.26	0.346	0.79
<i>Blarina carolinensis</i>	1991	0.30 \pm 0.09	0.75 \pm 0.27	2.53 \pm 1.03	1.35 \pm 1.68	2.193	0.13
	1992	0.22 \pm 0.06	0.29 \pm 0.09	0.34 \pm 0.21	0.30 \pm 0.10	0.306	0.82
	Pooled	0.26 \pm 0.06	0.52 \pm 0.15	1.44 \pm 0.61	0.83 \pm 0.40	1.847	0.16
<i>Ochrotomys nuttalli</i>	1991	0.72 \pm 0.29	0.34 \pm 0.17	0.81 \pm 0.33	0.08 \pm 0.05	2.022	0.15
	1992	0.07 \pm 0.07	0.07 \pm 0.07	0.15 \pm 0.09	0.08 \pm 0.05	0.514	0.68
	Pooled	0.40 \pm 0.18	0.19 \pm 0.10	0.48 \pm 0.20	0.08 \pm 0.11	1.695	0.19

*One-way ANOVA F-value.

†Probability associated with one-way ANOVA F-value.

‡Includes *P. leucopus*, *P. gossypinus*, *P. maniculatus*, and *P. attwateri*.

Table 3- Species richness, diversity and evenness ($\bar{x} \pm SE$) in Ecosystem Management research stands in the Ouachita Mountains of Arkansas by physiographic zone, 1991-92

Variable	Year	Zone				F*	P†
		North	South	East	West		
Richness	1991	3.20 \pm 0.20	3.40 \pm 0.51	3.20 \pm 0.21	3.60 \pm 0.87	0.133	0.94
	1992	2.20 \pm 0.37	2.80 \pm 0.37	2.60 \pm 0.40	2.60 \pm 0.51	0.362	0.78
	Pooled	3.60 \pm 0.24	3.83 \pm 0.48	3.40 \pm 0.25	4.00 \pm 0.63	0.346	0.19
Diversity‡	1991	0.88 \pm 0.10	0.88 \pm 0.20	0.88 \pm 0.12	0.89 \pm 0.23	0.028	0.99
	1992	0.55 \pm 0.17	0.84 \pm 0.13	0.76 \pm 0.09	0.70 \pm 0.19	0.694	0.57
	Pooled	0.96 \pm 0.14	1.02 \pm 0.21	0.89 \pm 0.13	1.06 \pm 0.05	0.214	0.89
Evenness	1991	0.77 \pm 0.09	0.76 \pm 0.12	0.76 \pm 0.09	0.65 \pm 0.17	0.210	0.89
	1992	0.61 \pm 0.17	0.84 \pm 0.04	0.87 \pm 0.07	0.67 \pm 0.17	0.936	0.45
	Pooled	0.74 \pm 0.09	0.72 \pm 0.13	0.73 \pm 0.08	0.81 \pm 0.05	0.173	0.91

*One-way ANOVA F-value.

†Probability associated with one-way ANOVA F-value.

‡Shannon's diversity index (Shannon and Weaver 1963).

Table 4- Sorensen's similarity indices (*Mueller-Dombois and Ellenberg 1974*) for small mammal communities in Ecosystem *Management research stands* in the *Ouachita Mountains of Arkansas* by future treatments and physiographic zones, 1991-92

Analysis	Comparison	Similarity index (percent)
Future treatments	Clearcut-single tree selection	92.3
	Clearcut-shelterwood	90.9
	Control-single tree selection	85.7
	Control-shelterwood	83.3
	Single tree selection-shelterwood	83.3
	Control-clearcut	76.9
	Group selection-clearcut	76.9
	Group selection-single tree selection	71.4
	Group selection-shelterwood	66.7
	Group selection-control	57.1
Physiographic zones	South-West	87.5
	North-East	80.0
	North-South	76.9
	North- West	76.9
	South-East	61.5
	East-West	61.5

DISCUSSION

Small mammals play an important role in several ecological processes of forested communities and can be greatly influenced by forest management activities. To effectively evaluate the impacts of imposing specific treatments on forest stands, pretreatment conditions relative to small mammal abundance and community composition should be as similar as possible. In this study, no differences were found in small mammal relative abundance, richness, diversity, or evenness by physiographic region, and with only one exception, no differences were found in these same parameters by **future** treatment. Thus, future analyses of posttreatment data should not be confounded by pretreatment differences in respect to small mammal community composition and stand location (physiographic zone).

Though differences in trap types and trapping methodologies prevent direct comparisons with other studies, all stands sampled in this study appeared to be characterized by a relatively low density and diversity of small mammals. Several studies have shown that small mammal abundance and diversity is influenced by successional vegetation **patterns** and structural habitat characteristics (Goodwin and Hungerford 1979; Kirkland 1977, 1990; **McComb** and Noble 1980; Mengak and others **1989a**, 1989b). In general, early successional **seres** are characterized by higher small mammal abundance and diversity than later seres, as well as differences in species composition.

Small mammal abundance, diversity, and species composition is often positively related to understory cover and down woody material. Stands in this study generally had very little down woody material (averaging 3.3 percent ground coverage), and the percentage of cover was low (averaging 0.3 to 2.3 percent) for woody plants, forbs, and graminoids (Thill and others 1994). Numerical and compositional responses of small mammals following treatments will most likely reflect associated increases in the above habitat parameters. The magnitude and temporal characteristics of these responses will probably vary by treatment, and total numbers of some species may increase substantially. Though increasing sampling intensity during **pretreatment** trapping did not increase trapping success, retaining two traps per station will probably be necessary to sample increased small mammal populations following certain treatments.

The three most commonly captured species in this study represented three different **trophic** groups: insectivores (*Blarina carolinensis*), **granivores** (*Ochrotomys nuttalli*), and omnivores (*Peromyscus* spp.). If sufficient numbers of additional species

Table 5- Relative abundance (captures per 100 trap nights) of small mammals and the three most commonly captured species ($\bar{x} \pm SE$) in Ecosystem Management research stands in the Ouachita Mountains of Arkansas by future treatment, 1991-92

Species	Year	Treatment					F	P [†]
		Clearcut	Shelterwood	Single-tree selection	Group selection	Control		
All species	1991	1.06 ± 0.23	3.90 ± 1.62	4.771 1.22	3.54 ± 1.34	4.36 ± 0.88	1.437	0.27
	1992	0.83 ± 0.15	1.30 ± 0.67	1.65 ± 0.31	1.39 ± 0.33	1.31 ± 0.35	0.931	0.46
	Pooled	0.83 ± 0.15	2.60 ± 0.93	3.21 ± 0.83	2.47 ± 0.84	2.83 ± 0.73	1.47s	0.23
<i>Peromyscus</i> spp. [‡]	1991	0.53 ± 0.16	2.57 ± 1.61	1.43 ± 0.62	2.06 ± 1.00	1.41 ± 0.32	0.70	0.60
	1992	0.37 ± 0.13	0.88 ± 0.50	1.04 ± 0.29	0.78 ± 0.29	0.84 ± 0.28	0.614	0.66
	Pooled	0.46 ± 0.10	1.73 ± 0.64	1.25 ± 0.32	1.42 ± 0.34	1.13 ± 0.23	0.942	0.43
<i>Biarina carolinensis</i>	1991	0.46 ± 0.12	0.74 ± 0.37	2.16 ± 1.08	0.47 ± 0.18	2.32 ± 0.11	1.308	0.23
	1992	0.18 ± 0.07	0.19 ± 0.08	0.28 ± 0.12	0.42 ± 0.09	0.36 ± 0.11	1.280	0.32
	Pooled	0.32 ± 0.08	0.46 ± 0.21	1.22 ± 0.62	0.43 ± 0.10	1.33 ± 0.11	1.298	0.29
<i>Ochrotomys nuttalli</i>	1991	0.23 ± 0.18	0.48 ± 0.36	0.67 ± 0.31	0.72 ± 0.42	0.33 ± 0.13	0.4%	0.73
	1992	0.00 ± 0.00	0.14 ± 0.06	0.14 ± 0.09	0.10 ± 0.10	0.03 ± 0.03	0.676	0.62
	Pooled	0.12 ± 0.09	0.31 ± 0.18	0.41 ± 0.18	0.41 ± 0.23	0.19 ± 0.09	0.629	0.63

*One-way ANOVA F-value.

†Probability associated with one-way ANOVA F-value.

‡Includes *P. leucopus*, *P. gossypinus*, *P. maniculatus*, and *P. atriventer*.

Table 6- Species richness, diversity, and evenness ($\bar{x} \pm SE$) in Ecosystem Management research stands in the Ouachita Mountains of Arkansas by future treatment, 1991-92

Variable	Year	Treatment					F [*]	P [†]
		Clearcut	Shelterwood	Single-tree selection	Group selection	Control		
Richness	1991	3.3 ± 0.25	2.75 ± 0.25	3.50 ± 0.50	3.00 ± 0.71	4.25 ± 0.73	1.152	0.37
	1992	2.00 ± 0.00	2.50 ± 0.87	3.00 ± 0.41	2.73 ± 0.25	2.50 ± 0.29	0.647	0.64
	Pooled	3.25 ± 0.25	3.40 ± 0.40	4.25 ± 0.48	3.50 ± 0.29	4.23 ± 0.73	1.062	0.41
Diversity [‡]	1991	1.03 ± 0.06	0.82 ± 0.21	0.97 ± 0.06	0.76 ± 0.26	0.93 ± 0.26	0.328	0.89
	1992	0.62 ± 0.04	0.62 ± 0.04	0.62 ± 0.04	0.86 ± 0.12	0.73 ± 0.14	1.167	0.36
	Pooled	1.04 ± 0.11	0.70 ± 0.11	1.13 ± 0.12	0.99 ± 0.07	0.87 ± 0.23	0.360	0.13
Evenness	1991	0.71 ± 0.17	0.71 ± 0.17	0.81 ± 0.08	0.59 ± 0.20	0.62 ± 0.11	0.8775	0.30
	1992	0.34 ± 0.20	0.34 ± 0.20	0.84 ± 0.05	0.86 ± 0.06	0.80 ± 0.10	4.212	0.02
	Pooled	0.88 ± 0.04	0.70 ± 0.16	0.84 ± 0.05	0.86 ± 0.06	0.59 ± 0.10	1.1%	0.33

*One-way ANOVA F-value. Means within rows followed by unlike letters are statistically different ($P < 0.05$).

†Probability associated with one-way ANOVA F-value.

‡Shannon's diversity index (Shannon and Weaver 1963).

are captured during posttreatment sampling, changes in community structure relative to **trophic** groups will be of interest, particularly in respect to fungi- and seed-consuming species due to their potential effects on natural regeneration.

ACKNOWLEDGMENTS

We gratefully acknowledge the assistance of the many people who participated in small mammal trapping, including USDA Forest Service Ouachita and Ozark National Forest district personnel, University of Arkansas at Little Rock biology students, J.F. **Taulman**, and R.W. **Perry**. We **also** thank B.R. Lockhart and R.A. Williams for reviewing an earlier **draft** of this manuscript.

LITERATURE CITED

- Baker, J.B. 1994. An overview of stand-level ecosystem management research in the **Ouachita/Ozark** National Forests. In: Proceedings of the Symposium on Ecosystem Management Research in the Ouachita Mountains: Pretreatment Conditions and Preliminary Findings, 1993. October 26-27: Hot Springs, AR Gen. Tech. Rep, New Orleans LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station.
- Goodwin, John G., Jr.; Hungerford, C. Roger. 1979. Rodent population densities and food habits in Arizona ponderosa pine forests. Res. Pap. RM-214. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station. 12 p.
- Harlow, Richard F.; Van Lear, David H. 1981. Silvicultural effects on wildlife habitat in the South (an annotated bibliography): **1953-1979**. Tech. Pap. 14. Clemson, SC: Clemson University Department of Forestry. 30 p.
- Harlow, Richard F.; Van Lear, David H. 1987. Silvicultural effects on wildlife habitat in the South (an annotated bibliography): **1980-1985**. Tech. Pap. 17. Clemson, SC: Clemson University Department of Forestry. 42 p.
- Kirkland, Gordon L., Jr. 1977. Responses of small mammals to the clearcutting of **northern** Appalachian forest. Journal of Mammalogy 58: 600-609.
- Kirkland, Gordon L., Jr. 1990. Patterns of initial small mammal community change after clearcutting of temperate North American forests. Oikos. **59(3)**: 313-320.
- Maser, Chris; Anderson, Ralph G.; Cromack, Kermit, Jr. [and others]. 1979. Dead and down woody material. In: Thomas, Jack Ward, tech ed. Wildlife habitats in managed forests: the Blue Mountains of Oregon and Washington. Agric. Handb. 553. Washington, DC: U.S. Department of Agriculture: 78-95. Chapter 6. In cooperation with: **The** Wildlife Management Institute; U.S. Department of the Interior, Bureau of Land Management.
- McComb**, W.C.; Noble, R.E. 1980. Small mammal and bird use of some unmanaged and managed forest stands in the mid-South. In: Proceedings of the 34th Annual Conference of the Southeastern Association of Fish and Wildlife Agencies. 482-49 **1**.
- Mengak, Michael T.; Guynn, David C., Jr.; Van Lear, David H. 1989a. Ecological implications of loblolly pine regeneration for small mammal communities. Forest Science **35(2)**: 503-514.
- Mengak, Michael T.; Van Lear, David H.; Guynn, David C., Jr. 1989b. Impacts of loblolly pine regeneration on selected wildlife habitat components. In: Miller, J.H., **comp.** Proceedings of the 5th biennial southern silvicultural research conference. Gen. Tech. Rep. **SO-74**. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station: 612-618.
- National Council of the Paper Industry for Air and Stream Improvement, Inc. 1993. Forestry, wildlife, and habitat in the East (an annotated bibliography), 1986-1990. In: *Sweeny*, S.R.; Wigley, T.B., **comps.** Tech. Bull. 65 1.3 17 p.
- Pank**, L.F. 1974. A bibliography on seed-eating mammals and birds that **affect** forest regeneration. **Spec.** 174. U.S. Department of the Interior, Fish and Wildlife Service. 28 p.
- Mueller-Dombois, D.; Ellenberg, H. 1974. Aims and methods of vegetation ecology. New York: John Wiley and Sons.

- Smith, C.F.; **Aldous, S.E.** 1947. The influence of mammals and birds in retarding artificial and natural regeneration of coniferous forests of the United States. *Journal of Forestry*. 45: **361-369**.
- Shannon, C.E.; Weaver, W. 1963. *The mathematical theory of communication*. Urbana, IL: University of Illinois Press. 177 p.
- Thill, R.E.**; **Tappe, P.A.**; **Koerth, N.E.** 1994. Wildlife habitat conditions in mature pine-hardwood stands in the **Ouachita/Ozark** National Forests. In: *Proceedings of the Symposium on Ecosystem Management Research in the Ouachita Mountains: Pretreatment Conditions and Preliminary Findings*. 1993. October 26-27: Hot Springs, AR. Gen. Tech. Rep. New Orleans LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station.
- Ursic, S.J.**; **Esher, R.J.** 1988. Influence of small mammals on **stormflow** responses of pine-covered **catchments**. *Water Resources Bulletin*. 24: 133-139.