

Forests and woodlands cover an extremely large area of the United States, 333 million ha or approximately 36 percent of the Nation's land area (Oswalt and others 2019). These forests possess the capacity to provide a broad range of goods and services for current and future generations, to safeguard biological diversity, and to contribute to the resilience of ecosystems, societies, and economies (USDA Forest Service 2011). Their ecological roles include supplying large and consistent quantities of clean water, preventing soil erosion, and providing habitat for a broad diversity of plant and animal species. Their socioeconomic benefits include wood products, nontimber goods, recreational opportunities, and pleasing natural beauty. Both the ecological integrity and the continued capacity of these forests to provide ecological and economic goods and services are of concern, however, in the face of a long list of threats, including insect and disease infestation, drought, fragmentation and forest conversion to other land uses, catastrophic fire, invasive species, and the effects of climate change.

Natural and anthropogenic stresses vary among biophysical regions and local environments; they also change over time and interact with each other. These and other factors make it challenging to establish baselines of forest health and to detect important departures from normal forest ecosystem functioning (Riitters and Tkacz 2004). Monitoring the health of forests is a critically important task, however, reflected within the Criteria and Indicators for

the Conservation and Sustainable Management of Temperate and Boreal Forests (Montréal Process Working Group 1995), which the Forest Service, U.S. Department of Agriculture, uses as a forest sustainability assessment framework (USDA Forest Service 2004, 2011). The primary objective of such monitoring is to identify ecological resources whose condition is deteriorating in subtle ways over large regions in response to cumulative stresses, a goal that requires consistent, large-scale, and long-term monitoring of key indicators of forest health status, change, and trends (Riitters and Tkacz 2004). This is best accomplished through the participation of multiple Federal, State, academic, and private partners.

Although the concept of a healthy forest has universal appeal, forest ecologists and managers have struggled with how exactly to define forest health (Teale and Castello 2011), and there is no universally accepted definition. Most definitions of forest health can be categorized as representing either an ecological or a utilitarian perspective (Kolb and others 1994). From an ecological perspective, the current understanding of ecosystem dynamics suggests that healthy ecosystems are those that are able to maintain their organization and autonomy over time while remaining resilient to stress (Costanza 1992), and that evaluations of forest health should emphasize factors that affect the inherent processes and resilience of forests (Edmonds and others 2011, Kolb and others 1994, Raffa and others 2009). On the other hand, the utilitarian

CHAPTER 1.

Introduction

KEVIN M. POTTER

perspective holds that a forest is healthy if management objectives are met, and that a forest is unhealthy if these objectives are not met (Kolb and others 1994). Although this definition may be appropriate when a single, unambiguous management objective exists, such as the production of wood fiber or the maintenance of wilderness attributes, it is too narrow when multiple management objectives are required (Edmonds and others 2011, Teale and Castello 2011). Teale and Castello (2011) incorporate both ecological and utilitarian perspectives into their two-component definition of forest health: first, a healthy forest must be sustainable with respect to its size structure, including a correspondence between baseline and observed mortality; second, a healthy forest must meet the landowner's objectives, provided that these objectives do not conflict with sustainability.

This national report, the 20th in an annual series sponsored by the Forest Health Monitoring (FHM) program of the Forest Service, attempts to quantify the status of, changes to, and trends in a wide variety of broadly defined indicators of forest health. The indicators described in this report encompass forest insect and disease activity, wildland fire occurrence, drought, tree mortality, and coarse woody materials, among others. The previous reports in this series are Ambrose and Conkling (2007, 2009), Conkling (2011), Conkling and others (2005), Coulston and others (2005a, 2005b, 2005c), and Potter and Conkling (2012a, 2012b, 2013a, 2013b, 2014, 2015a, 2015b, 2016, 2017, 2018, 2019,

2020). Visit <https://www.fs.fed.us/foresthealth/publications/fhm/fhm-annual-national-reports.shtml> for links to each of these reports in their entirety and for searchable lists of links to chapters included in the reports.

This report has three specific objectives. The first is to present information about forest health from a national perspective, or from a multi-State regional perspective when appropriate, using data collected by the Forest Health Protection (FHP) and Forest Inventory and Analysis (FIA) programs of the Forest Service, as well as from other sources available at a wide extent. The chapters that present analyses at a national scale, or multi-State regional scale, are divided between section 1 and section 2 of the report. Section 1 presents results from the analyses of forest health data that are available on an annual basis. Such repeated analyses of regularly collected indicator measurements allow for the detection of trends over time and help establish a baseline for future comparisons (Riitters and Tkacz 2004). Section 2 presents longer-term forest health trends, in addition to describing new techniques for analyzing forest health data at national or regional scales (the second objective of the report). While in-depth interpretation and analysis of specific geographic or ecological regions are beyond the scope of these parts of the report, the chapters in sections 1 and 2 present information that can be used to identify areas that may require investigation at a finer scale.


The second objective of the report is to present new techniques for analyzing forest health data as well as new applications of established techniques, often applied to longer timescales, presented in selected chapters of section 2. Examples in this report are chapters 6 and 7. Chapter 6 presents the results of analyses of nearly 20 years of FIA coarse woody material (CWM) data across spatial scales to assess change over time in CWM biomass, diameter and decay class, and species composition. Chapter 7, meanwhile, uses case studies to demonstrate the potential of rFIA, an open-source R package developed to improve the accessibility of FIA data and offer enhanced flexibility in estimation strategies, to advance forest health evaluation and monitoring.

The third objective of the report is to present results of recently completed Evaluation Monitoring (EM) projects funded through the FHM national program. These project summaries, presented in section 3, determine the extent, severity, and/or cause of forest health problems (FHM 2019), generally at a finer scale than that addressed by the analyses in sections 1 and 2. Each of the six chapters in section 3 contains an overview of an EM project, key results, and contacts for more information.



































When appropriate throughout this report, authors use the Forest Service revised ecoregions for the conterminous United States and Alaska

(Cleland and others 2007, Spencer and others 2002) as a common ecologically based spatial framework for their forest health assessments (fig. 1.1). Specifically, when the spatial scale of the data and the expectation of an identifiable pattern in the data are appropriate, authors use ecoregion sections, larger-scale provinces, or smaller-scale subsections as assessment units for their analyses. Bailey's hierarchical system bases the two broadest ecoregion scales, domains and divisions, on large ecological climate zones, while each division is broken into provinces based on vegetation macro features (Bailey 1995). Provinces are further divided into sections, which may be thousands of km² in area and are expected to encompass regions similar in their geology, climate, soils, potential natural vegetation, and potential natural communities (Cleland and others 1997). Subsections are nested within sections as the smallest level in the hierarchy. This hierarchical system does not address either Hawaii or Puerto Rico beyond including each in a unique, single ecoregion province (Bailey 1995). A set of Hawaii ecoregions based on moisture and elevational characteristics was developed for use in FHM national reports (Potter 2020) because a finer-scale and ecologically oriented spatial assessment framework was needed to estimate the impacts of a destructive forest disease (ch. 2) and of forest fire occurrences (ch. 3) (fig. 1.2, table 1.1).

(B)



Conterminous States ecoregion provinces

-  211: Northeastern Mixed Forest
-  M211: Adirondack-New England Mixed Forest—Coniferous Forest—Alpine Meadow
-  212: Laurentian Mixed Forest
-  221: Eastern Broadleaf Forest
-  M221: Central Appalachian Broadleaf Forest—Coniferous Forest—Meadow
-  222: Midwest Broadleaf Forest
-  223: Central Interior Broadleaf Forest
-  M223: Ozark Broadleaf Forest
-  231: Southeastern Mixed Forest
-  M231: Ouachita Mixed Forest—Meadow
-  232: Outer Coastal Plain Mixed Forest
-  234: Lower Mississippi Riverine Forest
-  242: Pacific Lowland Mixed Forest
-  251: Prairie Parkland (Temperate)
-  255: Prairie Parkland (Subtropical)
-  M242: Cascade Mixed Forest—Coniferous Forest—Alpine Meadow
-  261: California Coastal Chaparral Forest and Shrub
-  M261: Sierran Steppe—Mixed Forest—Coniferous Forest—Alpine Meadow
-  262: California Dry Steppe
-  M262: California Coastal Range Open Woodland—Shrub—Coniferous Forest—Meadow
-  263: California Coastal Steppe—Mixed Forest—Redwood Forest
-  313: Colorado Plateau Semi-Desert
-  M313: Arizona-New Mexico Mountains Semi-Desert—Open Woodland—Coniferous Forest—Alpine Meadow
-  315: Southwest Plateau and Plains Dry Steppe and Shrub
-  321: Chihuahuan Semi-Desert
-  322: American Semi-Desert and Desert
-  331: Great Plains—Palouse Dry Steppe
-  M331: Southern Rocky Mountain Steppe—Open Woodland—Coniferous Forest—Alpine Meadow
-  332: Great Plains Steppe
-  M332: Middle Rocky Mountain Steppe—Coniferous Forest—Alpine Meadow
-  M333: Northern Rocky Mountain Forest-Steppe—Coniferous Forest—Alpine Meadow
-  M334: Black Hills Coniferous Forest
-  341: Intermountain Semi-Desert and Desert
-  M341: Nevada-Utah Mountains Semi-Desert—Coniferous Forest—Alpine Meadow
-  342: Intermountain Semi-Desert
-  411: Everglades

Alaska ecoregion provinces

-  121: Arctic Tundra
-  M122: Bering Tundra
-  M131: Bering Taiga
-  M132: Intermontane Boreal
-  133: Alaska Range Transition
-  M134: Coastal Mountains Transition
-  M241: Coastal Rainforest
-  M243: Aleutian Meadows




Figure 1.2—Ecoregions, and ecoregion subunits, for Hawaii, developed based on moisture zones and elevation (see box 1.1). Ecoregion subunits are shown in the same color by ecoregion. See table 1.1 for the names of the ecoregion subunits listed on the map.

Table 1.1—The six ecoregions and 34 ecoregion subunits for the State of Hawaii

Ecoregion	Subunit	Ecoregion	Subunit
AL: Alpine	Alh: Alpine-Hawai'i	MW: Montane Wet	MWh-hp: Montane Wet-Hawai'i-Hilo-Puna
LW: Lowland Wet	LWh-hp: Lowland Wet-Hawai'i-Hilo-Puna		MWh-ka: Montane Wet-Hawai'i-Ka'ū
	LWh-kh: Lowland Wet-Hawai'i-Kohala-Hāmākua		MWh-kh: Montane Wet-Hawai'i-Kohala-Hāmākua
	LWk: Lowland Wet-Kaua'i		MWh-ko: Montane Wet-Hawai'i-Kona
	LWm-e: Lowland Wet-Maui-East		MWk: Montane Wet-Kaua'i
	LWm-w: Lowland Wet-Maui-West		MWl: Montane Wet-Lāna'i
	LWo: Lowland Wet-O'ahu		MWm-e: Montane Wet-Maui-East
			MWm-w: Montane Wet-Maui-West
			MWmo: Montane Wet-Moloka'i
			MWo: Montane Wet-O'ahu
LLD: Lowland/Leeward Dry	LLDh: Lowland/Leeward Dry-Hawai'i	SA: Subalpine	SAh: Subalpine-Hawai'i
	LLDka: Lowland/Leeward Dry-Kaho'olawe		SAm: Subalpine-Maui
	LLDk: Lowland/Leeward Dry-Kaua'i		
	LLDI: Lowland/Leeward Dry-Lāna'i		
	LLDm: Lowland/Leeward Dry-Maui		
	LLDmo: Lowland/Leeward Dry-Moloka'i		
	LLDn: Lowland/Leeward Dry-Ni'ihau		
	LLDo: Lowland/Leeward Dry-O'ahu		
ME: Mesic	MEh: Mesic-Hawai'i		
	MEk: Mesic-Kaua'i		
	MEl: Mesic-Lāna'i		
	MEm-e: Mesic-Maui-East		
	MEm-w: Mesic-Maui-West		
	MEmo: Mesic-Moloka'i		
	MEo: Mesic-O'ahu		

THE FOREST HEALTH MONITORING PROGRAM

The national FHM program is designed to determine the status, changes, and trends in indicators of forest condition on an annual basis and covers all forested lands through a partnership encompassing the Forest Service, State foresters, and other State and Federal agencies and academic groups (FHM 2019). The FHM program utilizes data from a wide variety of data sources, both inside and outside the Forest Service, and develops analytical approaches for addressing forest health issues that affect the sustainability of forest ecosystems. The FHM program has four major components (fig. 1.3):

- **Detection Monitoring**—nationally standardized aerial and ground surveys to evaluate status and change in condition of forest ecosystems (sections 1 and 2 of this report)
- **Evaluation Monitoring**—projects to determine the extent, severity, and causes of undesirable changes in forest health identified through Detection Monitoring (section 3 of this report)
- **Research on Monitoring Techniques**—work to develop or improve indicators, monitoring systems, and analytical techniques, such as urban and riparian forest health monitoring, early detection of invasive species, multivariate analyses of forest health indicators, and spatial scan statistics (section 2 of this report)




Figure 1.3—The design of the Forest Health Monitoring program (FHM 2003).

- **Analysis and Reporting**—synthesis of information from various data sources within and external to the Forest Service to produce issue-driven reports on status and change in forest health at national, regional, and State levels (sections 1, 2, and 3 of this report)

The FHM program, in addition to national reporting, generates regional and State reports, often in cooperation with FHM partners, both within the Forest Service and in State forestry and agricultural departments. For example, the FHM regions cooperate with their respective State partners to produce the annual Forest Health Highlights report series, available on the FHM website at <https://www.fs.fed.us/foresthealth/protecting-forest/forest-health-monitoring/monitoring-forest-highlights.shtml>. Other examples include Steinman (2004) and Harris and others (2011).

The FHM program and its partners also produce reports and journal articles on monitoring techniques and analytical methods (see <https://www.fs.fed.us/foresthealth/publications/fhm/fhm-publications.shtml>). The emphases of these publications include forest health data (Potter and others 2016, Siry and others 2018, Smith and Conkling 2004); soils as an indicator of forest health (O'Neill and others 2005); urban forest health monitoring (Bigsby and others 2014; Cumming and others 2006, 2007; Lake and others 2006); remote sensing of forest disturbances (Chastain and others 2015, Rebbeck and others 2015); health conditions in national forests (Morin and others 2006); crown conditions (Morin and others 2015; Randolph 2010a, 2010b, 2013; Randolph and Moser 2009; Schomaker and others 2007); indicators of regeneration (McWilliams and others 2015); vegetation diversity and structure (Schulz and

Gray 2013, Schulz and others 2009, Simkin and others 2016); forest lichen communities (Jovan and others 2012, Root and others 2014); down woody materials in forests (Woodall and others 2012, 2013); drought (Vose and others 2016); ozone monitoring (Rose and Coulston 2009); patterns of nonnative invasive plant occurrence (Guo and others 2015, 2017; Iannone and others 2015, 2016a, 2016b, 2018; Jo and others 2018; Oswalt and others 2015; Riitters and others 2018a, 2018b); assessments of forest risk or tree species vulnerability to exotic invasive forest insects and diseases (Koch and others 2011, 2014; Krist and others 2014; Potter and others 2019a, 2019b; Vogt and Koch 2016; Yemshanov and others 2014); spatial patterns of landcover and forest fragmentation (Guo and others 2018; Riitters 2011; Riitters and Costanza 2019; Riitters and Wickham 2012; Riitters and others 2012, 2016, 2017); impacts of deer browse on forest structure (Russell and others 2017); broad-scale assessments of forest biodiversity (Guo and others 2019; Potter 2018; Potter and Koch 2014; Potter and Woodall 2012, 2014); predictions and indicators of climate change effects on forests and forest tree species (Fei and others 2017, Heath and others 2015, Potter and Hargrove 2013); and the overall forest health indicator program (Woodall and others 2010).

For more information about the FHM program, visit the FHM website at <https://www.fs.fed.us/foresthealth/protecting-forest/forest-health-monitoring/>. Among other resources,

this website includes links to all past national forest health reports (<https://www.fs.fed.us/foresthealth/publications/fhm/fhm-annual-national-reports.shtml>), and annual State Forest Health Highlights reports (<https://www.fs.fed.us/foresthealth/protecting-forest/forest-health-monitoring/monitoring-forest-highlights.shtml>).

DATA SOURCES

Forest Service data sources in this edition of the FHM national report include FIA annualized Phase 2 survey data (Bechtold and Patterson 2005, Burrill and others 2018, Woodall and others 2010); FHP national Insect and Disease Survey forest mortality and defoliation data for 2019 (FHP 2020); Moderate Resolution Imaging Spectroradiometer (MODIS) Active Fire Detections for the United States data for 2019 (USDA Forest Service 2020); tree canopy cover data generated from the 2011 National Land Cover Database (NLCD) (Homer and others 2015) through a cooperative project between the Multi-Resolution Land Characteristics Consortium and Forest Service Geospatial Technology and Applications Center (GTAC) (Coulston and others 2012); and FIA's publicly available Environmental Monitoring and

Assessment Program (EMAP) hexagons (Brand and others 2000). Other sources of data include Parameter-elevation Regression on Independent Slopes Model (PRISM) climate mapping system data (PRISM Climate Group 2020), and Alaskan forest and shrub cover derived from the 2011 NLCD. For more information about the FIA program, which is a major source of data for several FHM analyses, see box 1.1.

FHM REPORT PRODUCTION

This FHM national report, the 20th in a series of such annual documents, is produced by forest health monitoring researchers at the Eastern Forest Environmental Threat Assessment Center (EFETAC) in collaboration with North Carolina State University cooperators in the Forest Health Monitoring Research Group (<https://go.ncsu.edu/foresthealth>). A unit of the Southern Research Station of the Forest Service, EFETAC was established under the Healthy Forests Restoration Act of 2003 to generate the knowledge and tools needed to anticipate and respond to environmental threats. For more information about the research team and about threats to U.S. forests, please visit <https://www.forestthreats.org/about>.

Box 1.1


The Forest Inventory and Analysis (FIA) program collects forest inventory information across all forest land ownerships in the United States and maintains a network of more than 130,000 permanent forested ground plots across the conterminous United States, southeastern Alaska, Hawaii, and the Caribbean and Pacific territories with a sampling intensity of approximately one plot/2428 ha (one plot per 6,000 acres). Forest Inventory and Analysis Phase 2 encompasses the annualized inventory measured on plots at regular intervals, with each plot surveyed every 5 to 7 years in most Eastern States, but with plots in the Rocky Mountain and Pacific Northwest regions surveyed once every 10 years (Reams and others 2005). The standard 0.067-ha plot (see figure) consists of four 7.315-m (24-foot) radius subplots (approximately 168.6 m² or 1/24th acre), on which field crews measure trees at least 12.7 cm (5 inches) in diameter. Within each of these subplots is nested a 2.073-m (6.8-foot) radius microplot (approximately 13.48 m² or 1/300th acre), on which crews measure trees smaller than 12.7 cm (5 inches) in diameter. A core-optional variant of the standard design includes four “macroplots,” each with a radius of 17.953 m or 58.9 feet (approximately 0.1012 ha or 1/4 acre) that originates at the center of each subplot (Burrill and others 2018).

Forest Inventory and Analysis Phase 3 plots previously represented a subset of these Phase 2 plots, with one Phase 3 plot for every 16 standard FIA Phase 2

plots. In addition to traditional forest inventory measurements, data for a variety of important ecological indicators were from Phase 3 plots, including tree crown condition, lichen communities, down woody material, soil condition, and vegetation structure and diversity, whereas data on ozone bioindicator plants were collected on a separate grid of plots (Woodall and others 2010, 2011). Most of these additional forest health indicators were measured as part of the Forest Health Monitoring Detection Monitoring ground plot system prior to 2000¹ (Palmer and others 1991). The FIA program recently updated its sampling techniques with flexible spatial and temporal intensities for some of these ecosystem health indicators (including down woody material, vegetation diversity and structure, and crown conditions) to improve field operation efficiency, address emerging user demands, and adjust to evolving forest health science (Castillo and Alvarez 2020). This “Phase 2 Plus Program/ Ecosystem Indicator Program” (P2+) sampling scheme facilitates the collection of a national core set of indicator information on more plots for less cost than the original indicator protocols, with sampling based on a systematic subsample that can change in response to budgetary

¹ U.S. Department of Agriculture Forest Service. 1998. Forest Health Monitoring 1998 field methods guide. Research Triangle Park, NC: U.S. Department of Agriculture Forest Service, Forest Health Monitoring program. 473 p. On file with: Forest Health Monitoring program, 3041 Cornwallis Rd., Research Triangle Park, NC 27709.

fluctuations without compromising long-term analytical capabilities. The enhanced indicator protocols collect less-detailed information on each sampled plot than on the previous Phase 3 plots, but substantially more plots are sampled, increasing the statistical power of forest health analyses and improving the reliability of estimates in important national assessments (Castillo and Alvarez 2020).



Box 1.1 figure—The Forest Inventory and Analysis mapped plot design. Subplot 1 is the center of the cluster with subplots 2, 3, and 4 located 120 feet away at azimuths of 360°, 120°, and 240°, respectively (Burrill and others 2018).

LITERATURE CITED

- Ambrose, M.J.; Conkling, B.L., eds. 2007. Forest Health Monitoring 2005 national technical report. Gen. Tech. Rep. SRS-104. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 76 p.
- Ambrose, M.J.; Conkling, B.L., eds. 2009. Forest Health Monitoring 2006 national technical report. Gen. Tech. Rep. SRS-117. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 118 p. <https://doi.org/10.2737/SRS-GTR-117>.
- Bailey, R.G. 1995. Descriptions of the ecoregions of the United States. 2d ed. Misc. Publ. No. 1391. Washington, DC: U.S. Department of Agriculture Forest Service. Map; presentation scale 1:7,500,000. 108 p.
- Bechtold, W.A.; Patterson, P.L., eds. 2005. The enhanced Forest Inventory and Analysis program—national sampling design and estimation procedures. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 85 p.
- Bigsby, K.M.; Ambrose, M.J.; Tobin, P.C.; Sills, E.O. 2014. The cost of gypsy moth sex in the city. *Urban Forestry & Urban Greening*. 13(3): 459–468. <https://doi.org/10.1016/j.ufug.2014.05.003>.
- Brand, G.J.; Nelson, M.D.; Wendt, D.G.; Nimerfro, K.K. 2000. The hexagon/panel system for selecting FIA plots under an annual inventory. In: McRoberts, R.E.; Reams, G.A.; Van Deusen, P.C., eds. *Proceedings of the first annual Forest Inventory and Analysis symposium*. Gen. Tech. Rep. NC-213. St. Paul, MN: U.S. Department of Agriculture Forest Service, North Central Research Station: 8–13.
- Burrill, E.A.; Wilson, A.M.; Turner, J.A. [and others]. 2018. The Forest Inventory and Analysis Database: database description and user guide for Phase 2 (version 8.0). Washington, DC: U.S. Department of Agriculture Forest Service. 946 p. <http://www.fia.fs.fed.us/library/database-documentation/>. [Date accessed: August 23, 2019].
- Castillo, P.S.B.; Alvarez, M. 2020. Forest Inventory and Analysis: fiscal year 2018 business report. FS-1153. Washington, DC: U.S. Department of Agriculture Forest Service. 71 p. <https://www.fia.fs.fed.us/library/bus-org-documents/docs/17973%20FS%20FIA%20Fiscal%20Year%202018%20Business%20Reportv3%20508.pdf>. [Date accessed: September 15, 2020].
- Chastain, R.A.; Fisk, H.; Ellenwood, J.R. [and others]. 2015. Near-real time delivery of MODIS-based information on forest disturbances. In: Lippitt, C.D.; Stow, D.A.; Coulter, L.L., eds. *Time sensitive remote sensing*. New York, NY: Springer: 147–164. https://doi.org/10.1007/978-1-4939-2602-2_10.
- Cleland, D.T.; Avers, P.E.; McNab, W.H. [and others]. 1997. National hierarchical framework of ecological units. In: Boyce, M.S.; Haney, A., eds. *Ecosystem management applications for sustainable forest and wildlife resources*. New Haven, CT: Yale University Press: 181–200.
- Cleland, D.T.; Freeouf, J.A.; Keys, J.E. [and others]. 2007. Ecological subregions: sections and subsections for the conterminous United States. Gen. Tech. Rep. WO-76D. Washington, DC: U.S. Department of Agriculture Forest Service. Map; Sloan, A.M., cartographer; presentation scale 1:3,500,000; colored. Also on CD-ROM as a GIS coverage in ArcINFO format or at <http://data.fs.usda.gov/geodata/edw/datasets.php>. [Date accessed: July 20, 2015].
- Conkling, B.L., ed. 2011. Forest Health Monitoring 2007 national technical report. Gen. Tech. Rep. SRS-147. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 159 p.
- Conkling, B.L.; Coulston, J.W.; Ambrose, M.J., eds. 2005. Forest Health Monitoring 2001 national technical report. Gen. Tech. Rep. SRS-81. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 204 p.
- Costanza, R. 1992. Toward an operational definition of ecosystem health. In: Costanza, R.; Norton, B.G.; Haskell, B.D., eds. *Ecosystem health: new goals for environmental management*. Washington, DC: Island Press: 239–256.
- Coulston, J.W.; Riitters, K.H.; Conkling, B.L., eds. 2005a. Forest Health Monitoring 2002 national technical report. Gen. Tech. Rep. SRS-84. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 97 p.

- Coulston, J.W.; Ambrose, M.J.; Riitters, K.H. [and others], eds. 2005b. Forest Health Monitoring 2003 national technical report. Gen. Tech. Rep. SRS-85. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 97 p.
- Coulston, J.W.; Ambrose, M.J.; Riitters, K.H.; Conkling, B.L., eds. 2005c. Forest Health Monitoring 2004 national technical report. Gen. Tech. Rep. SRS-90. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 81 p.
- Coulston, J.W.; Moisen, G.G.; Wilson, B.T. [and others]. 2012. Modeling percent tree canopy cover: a pilot study. *Photogrammetric Engineering and Remote Sensing*. 78(7): 715–727. <https://doi.org/10.14358/PERS.78.7.715>.
- Cumming, A.B.; Nowak, D.J.; Twardus, D.B. [and others]. 2007. Urban forests of Wisconsin: pilot monitoring project 2002. NA-FR-05-07. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northeastern Area State and Private Forestry. 33 p. Low resolution: http://www.na.fs.fed.us/pubs/fhm/pilot/pilot_study_wisconsin_02_lr.pdf. High resolution: http://www.na.fs.fed.us/pubs/fhm/pilot/pilot_study_wisconsin2_02_hr.pdf. [Date accessed: October 19, 2007].
- Cumming, A.B.; Twardus, D.B.; Smith, W.D. 2006. National Forest Health Monitoring program, Maryland and Massachusetts street tree monitoring pilot projects. NA-FR-01-06. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northeastern Area State and Private Forestry. 23 p.
- Edmonds, R.L.; Agee, J.K.; Gara, R.I. 2011. Forest health and protection. Long Grove, IL: Waveland Press, Inc. 667 p.
- Fei, S.; Desprez, J.M.; Potter, K.M. [and others]. 2017. Divergence of species responses to climate change. *Science Advances*. 3(5): e1603055. <https://doi.org/10.1126/sciadv.1603055>.
- Forest Health Monitoring (FHM). 2003. Forest Health Monitoring: a national strategic plan. 7 p. https://www.fs.fed.us/foresthealth/fhm/annc/strategic_plan03.pdf. [Date accessed: September 15, 2020].
- Forest Health Monitoring (FHM). 2019. Program description. Forest Health Monitoring fact sheet series. <https://www.fs.fed.us/foresthealth/protecting-forest/forest-health-monitoring/>. [Date accessed: August 23, 2019].
- Forest Health Protection (FHP). 2020. Insect and Disease Detection Survey (IDS) data downloads. Fort Collins, CO: U.S. Department of Agriculture Forest Service, Forest Health Technology Enterprise Team. <https://www.fs.fed.us/foresthealth/applied-sciences/mapping-reporting/detection-surveys.shtml>. [Date accessed: July 6, 2020].
- Guo, Q.; Fei, S.; Dukes, J.S. [and others]. 2015. A unified approach to quantify invasibility and degree of invasion. *Ecology*. 95(10): 2613–2621. <https://doi.org/10.1890/14-2172.1>.
- Guo, Q.; Fei, S.; Potter, K.M. [and others]. 2019. Tree diversity regulates forest pest invasion. *Proceedings of the National Academy of Sciences*. 116(15): 7382–7386. <https://doi.org/10.1073/pnas.1821039116>.
- Guo, Q.; Iannone, B.V.; Nunez-Mir, G.C. [and others]. 2017. Species pool, human population, and global vs. regional invasion patterns. *Landscape Ecology*. 32(2): 229–238. <https://doi.org/10.1007/s10980-016-0475-6>.
- Guo, Q.; Riitters, K.H.; Potter, K.M. [and others]. 2018. A subcontinental analysis of forest fragmentation and pest invasion. *Forests*. 9(12): 744. <https://doi.org/10.3390/f9120744>.
- Harris, J.L., comp.; Region 2 Forest Health Protection staff. 2011. Forest health conditions, 2009–2010: Rocky Mountain Region (R2). R2-11-RO-31. Golden, CO: U.S. Department of Agriculture Forest Service, Renewable Resources, Forest Health Protection, Rocky Mountain Region. 108 p.
- Heath, L.S.; Anderson, S.; Emery, M.R. [and others]. 2015. Indicators for climate change impacts for forests: National Climate Assessment indicators. Gen. Tech. Rep. NRS-155. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northern Research Station. 143 p.
- Homer, C.G.; Dewitz, J.A.; Yang, L. [and others]. 2015. Completion of the 2011 National Land Cover Database for the conterminous United States: representing a decade of land cover change information. *Photogrammetric Engineering and Remote Sensing*. 81(5): 345–354.
- Iannone, B.V.; Oswalt, C.M.; Liebhold, A.M. [and others]. 2015. Region-specific patterns and drivers of macroscale forest plant invasions. *Diversity and Distributions*. 21: 1181–1192. <https://doi.org/10.1111/ddi.12354>.

- Iannone, B.V.; Potter, K.M.; Guo, Q. [and others]. 2016a. Biological invasion hotspots: a trait-based perspective reveals new sub-continental patterns. *Ecography*. 39: 961–969. <https://doi.org/10.1111/ecog.01973>.
- Iannone, B.V.; Potter, K.M.; Guo, Q. [and others]. 2018. Environmental harshness drives spatial heterogeneity in biotic resistance. *NeoBiota*. 40: 87–105. <https://doi.org/10.3897/neobiota.40.28558>.
- Iannone, B.V.; Potter, K.M.; Hamil, K.-A.D. [and others]. 2016b. Evidence of biotic resistance to invasions in forests of the Eastern USA. *Landscape Ecology*. 31: 85–99. <https://doi.org/10.1007/s10980-015-0280-7>.
- Jo, I.; Potter, K.M.; Domke, G.; Fei, S. 2018. Dominant forest tree mycorrhizal type mediates understory plant invasions. *Ecology Letters*. 21: 217–224. <https://doi.org/10.1111/ele.12884>.
- Jovan, S.; Riddell, J.; Padgett, P.E.; Nash, T.H., III. 2012. Eutrophic lichens respond to multiple forms of N: implications for critical levels and critical loads research. *Ecological Applications*. 22(7): 1910–1922. <https://doi.org/10.1890/11-2075.1>.
- Koch, F.H.; Yemshanov, D.; Colunga-Garcia, M. [and others]. 2011. Potential establishment of alien-invasive forest insect species in the United States: where and how many? *Biological Invasions*. 13: 969–985.
- Koch, F.H.; Yemshanov, D.; Haack, R.A.; Magarey, R.D. 2014. Using a network model to assess risk of forest pest spread via recreational travel. *PLOS ONE*. 9(7): e102105. <https://doi.org/10.1371/journal.pone.0102105>.
- Kolb, T.E.; Wagner, M.R.; Covington, W.W. 1994. Concepts of forest health: utilitarian and ecosystem perspectives. *Journal of Forestry*. 92: 10–15.
- Krist, F.J., Jr.; Ellenwood, J.R.; Woods, M.E. [and others]. 2014. 2012–2027 national insect and disease forest risk assessment. FHTET-14-01. U.S. Department of Agriculture Forest Service, Forest Health Technology Enterprise Team. 199 p. http://www.fs.fed.us/foresthealth/technology/pdfs/2012_RiskMap_Report_web.pdf. [Date accessed: July 24, 2014].
- Lake, M.; Marshall, P.; Mielke, M. [and others]. 2006. National Forest Health Monitoring program monitoring urban forests in Indiana: pilot study 2002, part 1. Analysis of field methods and data collection. NA-FR-06-06. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northeastern Area State and Private Forestry. <http://www.fhm.fs.fed.us/pubs/ufhm/indianaforests02/indianaforests02.html>. [Date accessed: November 6, 2007].
- McWilliams, W.H.; Westfall, J.A.; Brose, P.H. [and others]. 2015. A regeneration indicator for Forest Inventory and Analysis: history, sampling, estimation, analytics, and potential use in the Midwest and Northeast United States. Gen. Tech. Rep. NRS-148. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northern Research Station. 74 p. <https://doi.org/10.2737/NRS-GTR-148>.
- Montréal Process Working Group. 1995. Criteria and indicators for the conservation and sustainable management of temperate and boreal forests. <http://www.montrealprocess.org/>. [Date accessed: March 4, 2015].
- Morin, R.S.; Liebhold, A.M.; Gottschalk, K.W. [and others]. 2006. Analysis of Forest Health Monitoring surveys on the Allegheny National Forest (1998–2001). Gen. Tech. Rep. NE-339. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northeastern Research Station. 102 p. <https://doi.org/10.2737/NE-GTR-339>.
- Morin, R.S.; Randolph, K.C.; Steinman, J. 2015. Mortality rates associated with crown health for eastern forest tree species. *Environmental Monitoring and Assessment*. 187(3): 87. <https://doi.org/10.1007/s10661-015-4332-x>.
- O'Neill, K.P.; Amacher, M.C.; Perry, C.H. 2005. Soils as an indicator of forest health: a guide to the collection, analysis, and interpretation of soil indicator data in the Forest Inventory and Analysis program. Gen. Tech. Rep. NC-258. St. Paul, MN: U.S. Department of Agriculture Forest Service, North Central Research Station. 53 p. <https://doi.org/10.2737/NC-GTR-258>.
- Oswalt, C.M.; Fei, S.; Guo, Q. [and others]. 2015. A subcontinental view of forest plant invasions. *NeoBiota*. 24: 49–54. <https://doi.org/10.3897/neobiota.24.4526>.

- Oswalt, S.N.; Smith, W.B.; Miles, P.D.; Pugh, S.A. 2019. Forest resources of the United States, 2017: a technical document supporting the Forest Service 2020 RPA Assessment. Gen. Tech. Rep. WO-97. Washington, DC: U.S. Department of Agriculture Forest Service, Washington Office. 223 p. <https://doi.org/10.2737/WO-GTR-97>.
- Palmer, C.J.; Riitters, K.H.; Strickland, T. [and others]. 1991. Monitoring and research strategy for forests—Environmental Monitoring and Assessment Program (EMAP). EPA/600/4-91/012. Washington, DC: U.S. Environmental Protection Agency. 189 p.
- Potter, K.M. 2018. Do United States protected areas effectively conserve forest tree rarity and evolutionary distinctiveness? *Biological Conservation*. 224: 34–46. <https://doi.org/10.1016/j.biocon.2018.05.007>.
- Potter, K.M. 2020. Introduction. In: Potter, K.M.; Conkling, B.L., eds. Forest Health Monitoring: national status, trends, and analysis 2019. Gen. Tech. Rep. SRS-250. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station: 5–24.
- Potter, K.M.; Conkling, B.L., eds. 2012a. Forest Health Monitoring 2008 national technical report. Gen. Tech. Rep. SRS-158. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 179 p.
- Potter, K.M.; Conkling, B.L., eds. 2012b. Forest Health Monitoring 2009 national technical report. Gen. Tech. Rep. SRS-167. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 252 p.
- Potter, K.M.; Conkling, B.L., eds. 2013a. Forest Health Monitoring: national status, trends, and analysis 2010. Gen. Tech. Rep. SRS-176. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 162 p.
- Potter, K.M.; Conkling, B.L., eds. 2013b. Forest Health Monitoring: national status, trends, and analysis 2011. Gen. Tech. Rep. SRS-185. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 149 p.
- Potter, K.M.; Conkling, B.L., eds. 2014. Forest Health Monitoring: national status, trends, and analysis 2012. Gen. Tech. Rep. SRS-198. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 192 p.
- Potter, K.M.; Conkling, B.L., eds. 2015a. Forest Health Monitoring: national status, trends, and analysis 2013. Gen. Tech. Rep. SRS-207. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 199 p.
- Potter, K.M.; Conkling, B.L., eds. 2015b. Forest Health Monitoring: national status, trends, and analysis 2014. Gen. Tech. Rep. SRS-209. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 190 p.
- Potter, K.M.; Conkling, B.L., eds. 2016. Forest Health Monitoring: national status, trends, and analysis 2015. Gen. Tech. Rep. SRS-213. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 199 p.
- Potter, K.M.; Conkling, B.L., eds. 2017. Forest Health Monitoring: national status, trends, and analysis 2016. Gen. Tech. Rep. SRS-222. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 195 p.
- Potter, K.M.; Conkling, B.L., eds. 2018. Forest Health Monitoring: national status, trends, and analysis 2017. Gen. Tech. Rep. SRS-233. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 190 p.
- Potter, K.M.; Conkling, B.L., eds. 2019. Forest Health Monitoring: national status, trends, and analysis 2018. Gen. Tech. Rep. SRS-239. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 168 p.
- Potter, K.M.; Conkling, B.L., eds. 2020. Forest Health Monitoring: national status, trends, and analysis 2019. Gen. Tech. Rep. SRS-250. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 189 p.

- Potter, K.M.; Escanferla, M.E.; Jetton, R.M.; Man, G. 2019a. Important insect and disease threats to United States tree species and geographic patterns of their potential impacts. *Forests*. 10(4): 304. <https://doi.org/10.3390/f10040304>.
- Potter, K.M.; Escanferla, M.E.; Jetton, R.M. [and others]. 2019b. Prioritizing the conservation needs of United States tree species: evaluating vulnerability to forest insect and disease threats. *Global Ecology and Conservation*. 18: e00622. <https://doi.org/10.1016/j.gecco.2019.e00622>.
- Potter, K.M.; Hargrove, W.W. 2013. Quantitative metrics for assessing predicted climate change pressure on North American tree species. *Mathematical and Computational Forestry and Natural Resources Sciences*. 5(2): 151–169.
- Potter, K.M.; Koch, F.H. 2014. Phylogenetic community structure of forests across the conterminous United States: regional ecological patterns and forest health implications. *Forest Science*. 60(5): 851–861. <https://doi.org/10.5849/forsci.13-115>.
- Potter, K.M.; Koch, F.H.; Oswalt, C.M.; Iannone, B.V. 2016. Data, data everywhere: detecting spatial patterns in fine-scale ecological information collected across a continent. *Landscape Ecology*. 31: 67–84. <https://doi.org/10.1007/s10980-015-0295-0>.
- Potter, K.M.; Woodall, C.W. 2012. Trends over time in tree and seedling phylogenetic diversity indicate regional differences in forest biodiversity change. *Ecological Applications*. 22(2): 517–531. <https://doi.org/10.1890/10-2137.1>.
- Potter, K.M.; Woodall, C.W. 2014. Does biodiversity make a difference? Relationships between species richness, evolutionary diversity, and aboveground live tree biomass across U.S. forests. *Forest Ecology and Management*. 321: 117–129. <https://doi.org/10.1016/j.foreco.2013.06.026>.
- PRISM Climate Group. 2020. 2.5-arcmin (4 km) gridded monthly climate data. <http://www.prism.oregonstate.edu>. [Date accessed: July 13, 2020].
- Raffa, K.F.; Aukema, B.; Bentz, B.J. [and others]. 2009. A literal use of “forest health” safeguards against misuse and misapplication. *Journal of Forestry*. 107: 276–277.
- Randolph, K.C. 2010a. Equations relating compacted and uncompacted live crown ratio for common tree species in the South. *Southern Journal of Applied Forestry*. 34(3): 118–123. <https://doi.org/10.1093/sjaf/34.3.118>.
- Randolph, K.C. 2010b. Comparison of the arithmetic and geometric means in estimating crown diameter and crown cross-sectional area. *Southern Journal of Applied Forestry*. 34(4): 186–189. <https://doi.org/10.1093/sjaf/34.4.186>.
- Randolph, K.C. 2013. Development history and bibliography of the U.S. Forest Service crown-condition indicator for forest health monitoring. *Environmental Monitoring and Assessment*. 185(6): 4977–4993. <https://doi.org/10.1007/s10661-012-2919-z>.
- Randolph, K.C.; Moser, W.K. 2009. Tree crown condition in Missouri, 2000–2003. Gen. Tech. Rep. SRS-113. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 11 p.
- Reams, G.A.; Smith, W.D.; Hansen, M.H. [and others]. 2005. The Forest Inventory and Analysis sampling frame. In: Bechtold, W.A.; Patterson, P.L., eds. The enhanced Forest Inventory and Analysis program—national sampling design and estimation procedures. Gen. Tech. Rep. SRS-80. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station: 11–26.
- Rebbeck, J.; Kloss, A.; Bowden, M. [and others]. 2015. Aerial detection of seed-bearing female *Ailanthus altissima*: a cost-effective method to map an invasive tree in forested landscapes. *Forest Science*. 61: 1068–1078. <https://doi.org/10.5849/forsci.14-223>.
- Riitters, K.H. 2011. Spatial patterns of land cover in the United States: a technical document supporting the Forest Service 2010 RPA assessment. Gen. Tech. Rep. SRS-136. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 64 p.
- Riitters, K.; Costanza, J. 2019. The landscape context of family forests in the United States: anthropogenic interfaces and forest fragmentation from 2001 to 2011. *Landscape and Urban Planning*. 188: 64–71. <https://doi.org/10.1016/j.landurbplan.2018.04.001>.
- Riitters, K.H.; Costanza, J.K.; Buma, B. 2017. Interpreting multiscale domains of tree cover disturbance patterns in North America. *Ecological Indicators*. 80: 147–152. <https://doi.org/10.1016/j.ecolind.2017.05.022>.
- Riitters, K.H.; Coulston, J.W.; Wickham, J.D. 2012. Fragmentation of forest communities in the Eastern United States. *Forest Ecology and Management*. 263: 85–93. <https://doi.org/10.1016/j.foreco.2011.09.022>.

- Riitters, K.; Potter, K.M.; Iannone, B.V. [and others]. 2018a. Exposure of protected forest to plant invasions in the Eastern United States. *Forests*. 9(11): 723. <https://doi.org/10.3390/f9110723>.
- Riitters, K.; Potter, K.M.; Iannone, B.V. [and others]. 2018b. Landscape correlates of forest plant invasions: a high-resolution analysis across the Eastern United States. *Diversity and Distributions*. 24: 274–284. <https://doi.org/10.1111/ddi.12680>.
- Riitters, K.H.; Tkacz, B. 2004. The U.S. Forest Health Monitoring program. In: Wiersma, G.B., ed. *Environmental monitoring*. Boca Raton, FL: CRC Press: 669–683. <https://doi.org/10.1201/9780203495476.ch30>.
- Riitters, K.H.; Wickham, J.D. 2012. Decline of forest interior conditions in the conterminous United States. *Scientific Reports*. 2: 653. <https://doi.org/10.1038/srep00653>.
- Riitters, K.H.; Wickham, J.D.; Costanza, J.K.; Vogt, P. 2016. A global evaluation of forest interior area dynamics using tree cover data from 2000 to 2012. *Landscape Ecology*. 31: 137–148. <https://doi.org/10.1007/s10980-015-0270-9>.
- Root, H.T.; McCune, B.; Jovan, S. 2014. Lichen communities and species indicate climate thresholds in southeast and south-central Alaska, USA. *The Bryologist*. 117(3): 241–252. <https://doi.org/10.1639/0007-2745-117.3.241>.
- Rose, A.K.; Coulston, J.W. 2009. Ozone injury across the Southern United States, 2002–06. *Gen. Tech. Rep. SRS-118*. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 25 p.
- Russell, M.B.; Woodall, C.W.; Potter, K.M. [and others]. 2017. Interactions between white-tailed deer density and the composition of forest understories in the Northern United States. *Forest Ecology and Management*. 384: 26–33. <https://doi.org/10.1016/j.foreco.2016.10.038>.
- Schomaker, M.E.; Zarnoch, S.J.; Bechtold, W.A. [and others]. 2007. Crown-condition classification: a guide to data collection and analysis. *Gen. Tech. Rep. SRS-102*. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 78 p. <https://doi.org/10.2737/SRS-GTR-102>.
- Schulz, B.K.; Bechtold, W.A.; Zarnoch, S.J. 2009. Sampling and estimation procedures for the vegetation diversity and structure indicator. *Gen. Tech. Rep. PNW-781*. Portland, OR: U.S. Department of Agriculture Forest Service, Pacific Northwest Research Station. 53 p. <https://doi.org/10.2737/PNW-GTR-781>.
- Schulz, B.K.; Gray, A.N. 2013. The new flora of Northeastern USA: quantifying introduced plant species occupancy in forest ecosystems. *Environmental Monitoring and Assessment*. 185: 3931–3957. <https://doi.org/10.1007/s10661-012-2841-4>.
- Simkin, S.M.; Allen, E.B.; Bowman, W.D. [and others]. 2016. Conditional vulnerability of plant diversity to atmospheric nitrogen deposition across the United States. *Proceedings of the National Academy of Sciences*. 113: 4086–4091. <https://doi.org/10.1073/pnas.1515241113>.
- Siry, J.; Cubbage, F.W.; Potter, K.M.; McGinley, K. 2018. Current perspectives on sustainable forest management: North America. *Current Forestry Reports*. 4(3): 138–149. <https://doi.org/10.1007/s40725-018-0079-2>.
- Smith, W.D.; Conkling, B.L. 2004. Analyzing forest health data. *Gen. Tech. Rep. SRS-077*. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 33 p. http://www.srs.fs.usda.gov/pubs/gtr/gtr_srs077.pdf. [Date accessed: November 6, 2007].
- Spencer, P.; Nowacki, G.; Fleming, M. [and others]. 2002. Home is where the habitat is: an ecosystem foundation for wildlife distribution and behavior. *Arctic Research of the United States*. 16: 6–17.
- Steinman, J. 2004. Forest Health Monitoring in the Northeastern United States: disturbances and conditions during 1993–2002. *NA-TP-01-04*. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northeastern Area State and Private Forestry. 46 p. http://fhm.fs.fed.us/pubs/tp/dist_cond/dc.shtml. [Date accessed: December 8, 2009].
- Teale, S.A.; Castello, J.D. 2011. The past as key to the future: a new perspective on forest health. In: Castello, J.D.; Teale, S.A., eds. *Forest health: an integrated perspective*. New York: Cambridge University Press: 3–16. <https://doi.org/10.1017/CBO9780511974977.002>.

- U.S. Department of Agriculture (USDA) Forest Service. 2004. National report on sustainable forests—2003. FS-766. Washington, DC: U.S. Department of Agriculture Forest Service. 139 p.
- U.S. Department of Agriculture (USDA) Forest Service. 2011. National report on sustainable forests—2010. FS-979. Washington, DC: U.S. Department of Agriculture Forest Service. 134 p.
- U.S. Department of Agriculture (USDA) Forest Service. 2020. Fire detection GIS data. <https://fsapps.nwcg.gov/afm/gisdata.php>. [Date accessed: June 24, 2020].
- Vogt, J.T.; Koch, F.H. 2016. The evolving role of Forest Inventory and Analysis data in invasive insect research. *American Entomologist*. 62(1): 46–58. <https://doi.org/10.1093/ae/tmv072>.
- Vose, J.M.; Clark, J.S.; Luce, C.H.; Patel-Weynand, T., eds. 2016. Effects of drought on forests and rangelands in the United States: a comprehensive science synthesis. Gen. Tech. Rep. WO-93b. Washington, DC: U.S. Department of Agriculture Forest Service, Washington Office. 289 p. <https://doi.org/10.2737/WO-GTR-93b>.
- Woodall, C.W.; Amacher, M.C.; Bechtold, W.A. [and others]. 2011. Status and future of the forest health indicators program of the USA. *Environmental Monitoring and Assessment*. 177: 419–436. <https://doi.org/10.1007/s10661-010-1644-8>.
- Woodall, C.W.; Conkling, B.L.; Amacher, M.C. [and others]. 2010. The Forest Inventory and Analysis database version 4.0: database description and users manual for Phase 3. Gen. Tech. Rep. NRS-61. Newtown Square, PA: U.S. Department of Agriculture Forest Service, Northern Research Station. 180 p. <https://doi.org/10.2737/NRS-GTR-61>.
- Woodall, C.W.; Walters, B.F.; Oswalt, S.N. [and others]. 2013. Biomass and carbon attributes of downed woody materials in forests of the United States. *Forest Ecology and Management*. 305: 48–59. <https://doi.org/10.1016/j.foreco.2013.05.030>.
- Woodall, C.W.; Walters, B.F.; Westfall, J.A. 2012. Tracking downed dead wood in forests over time: development of a piece matching algorithm for line intercept sampling. *Forest Ecology and Management*. 277: 196–204. <https://doi.org/10.1016/j.foreco.2012.04.031>.
- Yemshanov, D.; Koch, F.H.; Lu, B. [and others]. 2014. There is no silver bullet: the value of diversification in planning invasive species surveillance. *Ecological Economics*. 104: 61–72. <https://doi.org/10.1016/j.ecolecon.2014.04.024>.