

Effects of urbanization on the occurrence of *Batrachochytrium dendrobatidis*: do urban environments provide refuge from the amphibian chytrid fungus?

Daniel Saenz · Taylor L. Hall · Matthew A. Kwiatkowski

Published online: 3 August 2014
© Springer Science+Business Media New York 2014

Abstract *Batrachochytrium dendrobatidis* (*Bd*) is a widespread pathogenic fungus that is known to cause the disease, chytridiomycosis, which can be lethal to many amphibians. We compared occurrence rates on spring peepers (*Pseudacris crucifer*) in urban and forested breeding sites in eastern Texas, USA. All study sites were at approximately the same latitude and altitude, and samples were collected at the same time of year to isolate differences in *Bd* infection rates between habitat types. We found significant differences ($p < 0.001$) in the occurrence of *Bd* between habitats; with dramatically lower rates of occurrence at urban sites (19.5 %), compared to forested sites (62.9 %). The exact reason for the observed differences in the occurrence of *Bd* is not known, however, we suspect that warmer temperatures or lower population densities and lower species richness at urban sites all could play a role in our results. Our findings suggest that urban environments may provide a refuge for some amphibians from the pathogen.

Keywords Amphibians · Chytrid fungus · Urbanization · Habitat

Introduction

Most ecosystems are currently experiencing a loss of biodiversity, much of which is associated with human activity (Soule 1991; Naeem et al. 1994). Land-use change can facilitate local and regional extinction of populations by killing organisms, removing suitable habitat, and interfering

D. Saenz (✉)
Southern Research Station, Forest Service, U. S. Department of Agriculture, 506 Hayter Street,
Nacogdoches, TX 75965-3556, USA
e-mail: dsaenz@fs.fed.us

T. L. Hall
Department of Science and Mathematics, Angelina College, Lufkin, TX, USA

M. A. Kwiatkowski
Department of Biology, Stephen F. Austin State University, 1936 North Street, Nacogdoches,
TX 75962-3050, USA

with breeding activities (Collins and Storfer 2003). However, depending on the level of urbanization, some taxa may actually increase in richness due to factors such as spatial heterogeneity, intermediate disturbance, and scale (McKinney 2008).

Amphibian population declines, a global pattern that has garnered much interest in the past few decades, are certainly a major component of the overall loss of diversity (Wake 1991; Pechmann and Wilbur 1994; Houlahan et al. 2000, 2010). Along with population declines, amphibians have the highest global extinction rate among vertebrates (Stuart et al. 2004). Although habitat loss and urbanization are major threats to amphibians (Collins and Storfer 2003; Hamer and McDonnell 2008; McKinney 2008), it has become increasingly clear that many factors and combinations of these factors play important roles in the world-wide decline of amphibian populations (Alford and Richards 1999; Houlahan et al. 2000).

In addition to habitat loss and urbanization, the pathogenic fungus, *Batrachochytrium dendrobatidis* (*Bd*) is a significant threat to amphibians by causing the lethal skin disease, chytridiomycosis (Berger et al. 1998; Longcore et al. 1999; Lips et al. 2006). The pathogen is known to occur on every continent except Antarctica (Fisher et al. 2009) and has caused the decline or extinction of about 200 species of frog worldwide (Daszak et al. 2003; Skerratt et al. 2007). *Batrachochytrium dendrobatidis* may be the most serious infectious disease threat to biodiversity (Kilpatrick et al. 2010), and is now listed as an internationally notifiable disease by the World Organization for Animal Health as an epidemiologically important pathogen (Schloegel et al. 2010).

Occurrence and infectiousness of *Bd* are variable and known to be influenced by several factors. Higher altitude (Brem and Lips 2008), higher latitude (Kriger et al. 2007; Rodder et al. 2008), wetter habitats (Murphy et al. 2009, 2011), and cool seasons (Berger et al. 2004; Kriger and Hero 2008; Lannoo et al. 2011) all increase the incidence of *Bd* and decrease survival of infected amphibians. Temperature seems to be a primary factor that influences occurrence and infectiousness (Muths et al. 2008; Olson et al. 2013) but there is evidence that greater species richness increases the likelihood of infection and transmission of *Bd* (Olson et al. 2013).

Recent studies are beginning to shed light on the relationships between human impacts on the environment and *Bd*. Pauza et al. (2010) suggested that anthropomorphic activities promote the spread of *Bd*, but there is little evidence linking habitat loss with increases in the incidence of the pathogen. However, Van Sluys and Hero (2009) found higher densities of stony creek frogs (*Litoria wilcoxii*), and lower prevalence of *Bd* in open farmland compared to nearby forested habitat, suggesting that habitat strongly influenced *Bd* prevalence. Becker and Zamudio (2011) provided further direct evidence that the prevalence and infection intensity of *Bd* was inversely correlated to habitat loss in Costa Rica, Brazil, and Australia, possibly caused by increased temperatures resulting from habitat loss. Similarly, Lane and Burgin (2008) observed greater frog diversity and abundance in urban sites compared to non-urban sites in Australia but were only able to speculate that chemicals in urban wastewater provided protection from diseases, such as chytridiomycosis.

Given that habitat loss and chytridiomycosis are two major threats to biodiversity, and that current literature (Lane and Burgin 2008; Van Sluys and Hero 2009; Becker and Zamudio 2011) suggests likely interactions between the two factors, further explorations of these relationships is warranted. To date, no definitive evidence exists pertaining to the effects of urbanization on the relationship between the fungal pathogen and amphibian populations. Herein, we compare the incidence of *Bd* between amphibian populations from urban and forested habitats. Our findings provide insights for how habitat loss may affect host-pathogen interactions.

Methods

We chose the spring peeper (*Pseudacris crucifer*) as our focal species because it is known to occur in urban and forested wetlands and actively breeds in the cooler months of the year (Saenz et al. 2006) when the prevalence of *Bd* should be highest (Berger et al. 2004; Woodhams and Alford 2005; Kinney et al. 2011). In addition, spring peepers are abundant and known to have a relatively high incidence of *Bd* in our study area (Saenz et al. 2010).

We sampled frogs for *Bd* at six urban sites within the city of Nacogdoches, Texas, USA (31.6089° N, 94.6508° W); two forested sites in the Stephen F. Austin Experimental Forest (SFA EF) in Nacogdoches County, Texas (31.4667° N, 94.7833° W); and six forested sites in the Davy Crockett National Forest (DCNF) in Houston County, Texas (31.3333° N, 94.0833° W). The urban sites consisted of a variety of habitats including, drainage ditches, associated overflow areas, and a flooded soccer field (Fig. 1). These sites tended to be more exposed with less vegetative cover than forested sites. Roads, housing, and human structures were in close proximity to all urban sites (Fig. 1). The forested sites were all naturally occurring ephemeral wetlands in habitat dominated by secondary growth loblolly (*Pinus taeda*), and shortleaf (*P. echinata*) pines with a mixed deciduous hardwood component (Fig. 2).

We sampled individual spring peepers for *Bd* at 14 different sites on seven nights from 24 January to 16 March 2010. We collected as many spring peepers, by hand, as possible on a single trip to each site, to avoid resampling of individuals. Individual frogs were typically located by listening for calling activity then homing in on individual frogs. This technique was quite effective and we feel that we were able to capture over 95 % of the frogs that we observed calling in both habitat types. We sampled for *Bd* by rubbing a sterile cotton swab on the dorsum, ventral surfaces and feet of each frog for approximately 30 s. After swabbing, we immediately released the frogs at their place of capture. We used sterile techniques throughout the swabbing process; including using a new pair of gloves for handling each individual frog and autoclaving the cotton swabs prior to sampling, to prevent cross-contamination among samples. Swabs were placed into microcentrifuge tubes containing 1 ml of 70 % ethanol, stored in a freezer, and later analyzed for *Bd* DNA by polymerase chain reaction (PCR)



Fig. 1 Typical urban swabbing site, a drainage ditch behind a shopping center in Nacogdoches, Texas. (Site Nacogdoches-2)



Fig. 2 Typical forested swabbing site, an ephemeral wetland in the Davy Crockett National Forest, Houston County, Texas. (Site DCFN-3)

analysis (Pisces Molecular Laboratory, Boulder, Colorado, USA). Percent occurrence of *Bd* among frogs sampled was calculated per location, and Chi-square analyses were used to determine if *Bd* occurrence differed between the habitat types ($p < 0.05$).

Results

Spring peeper population densities appeared to be higher in forested habitats than in urban sites, as evidenced by the number of samples taken from each habitat type, and the comparable sampling efforts per habitat. We collected 41 spring peepers from six urban wetlands and 89 individuals from eight sites in forested habitats. PCR results indicated that spring peepers from forested habitats had significantly higher incidence of *Bd* than urban frogs ($\chi^2 = 21.16$, $df = 1$, $p < 0.001$). Of the individuals collected in the city of Nacogdoches, 19.5 % tested *Bd*-positive (Table 1), whereas 56.5 and 77.8 % tested positive from the DCFN and the SFA EF (Table 2), respectively. All frogs collected in this study appeared to be healthy. No dead or dying frogs were observed at our study sites.

Table 1 *Batrachochytrium dendrobatidis* PCR results from swabs taken from spring peepers collected in and around urban wetlands in the city of Nacogdoches in Nacogdoches County, Texas

Urban sites	Date	+/Total	% Positive
Nacogdoches-1	24 Jan 2010	0/6	0 %
Nacogdoches-2	28 Jan 2010	0/4	0 %
Nacogdoches-3	28 Jan 2010	0/12	0 %
Nacogdoches-4	9 Mar 2010	2/6	33.3 %
Nacogdoches-5	9 Mar 2010	1/2	50 %
Nacogdoches-6	10 Mar 2010	5/11	45.45 %
Total urban		8/41	19.51 %

Table 2 *Batrachochytrium dendrobatidis* PCR results from swabs taken from spring peepers collected in the Stephen F. Austin Experimental Forest in Nacogdoches County, Texas and the Davy Crockett National Forest in Houston County, Texas

Forested sites	Date	+/Total	% Positive
SFA EF-1	21 Feb 2010	8/10	80 %
SFA EF-2	21 Feb 2010	13/17	76.47 %
	SFA EF total	21/27	77.77 %
DCNF-1	15 Mar 2010	18/21	85.71 %
DCNF-2	15 Mar 2010	0/2	0 %
DCNF-3	15 Mar 2010	4/7	57.14 %
DCNF-4	16 Mar 2010	11/21	52.38 %
DCNF-5	16 Mar 2010	0/1	0 %
DCNF-6	16 Mar 2010	2/10	20 %
	DCNF total	35/62	56.45 %
Forested total		56/89	62.92 %

Discussion

Unlike studies that sampled and compared forested and deforested non-urban sites, we compared forested sites with urban sites to provide new insights into the relationship between human activity and *Bd*. Our findings that frogs in forested sites were significantly more likely to test positive for *Bd* than frogs in urban sites seem congruent with recent studies concluding that the risk of *Bd* contraction is higher in natural habitats (Van Sluys and Hero 2009; Becker and Zamudio 2011). Van Sluys and Hero (2009) found the incidence of *Bd* in forested sites to be 56.7 % compared to 29.8 % at open sites. In our study, we found a similar incidence of *Bd* in forested sites (62.9 %) but an even lower rate in urban sites (19.5 %). In effect, the difference in the occurrence of *Bd* in their study was just below 2:1 while we found a greater than three-fold difference between urban and forested sites. We suggest that the addition of urbanization to a deforested habitat may augment the previously established effects of deforestation on *Bd* infectiousness; whereas deforested non-urban sites may be intermediate by comparison (between urban and forested). However, further investigation is warranted to determine if this trend continues across multiple urban settings.

Although the pattern of differences in the occurrence of *Bd* between habitats appears to be consistent among recent studies, the proximate causes for these differences are still undetermined. Urban pollution may be a factor that can negatively affect *Bd*, as suggested by Lane and Burgin (2008). There are several chemical treatments that can clear *Bd* infections (Johnson et al. 2003; Berger et al. 2009; Woodhams et al. 2012); therefore, it is plausible that urban pollution may act in a similar manner. Depressional wetlands in urban areas can be sinks for sediments, nutrients, metals, greater nitrogen mineralization and nitrification (Faulkner 2004). In a recent study, Parris and Baud (2004) suggest that the synergistic effects of copper and *Bd* on amphibians may be negative, which would contradict the hypothesis that urban pollution is providing protection from *Bd*; however, sodium chloride has been shown to inhibit the growth of *Bd*, increase host survival, and has been suggested as a potentially useful antifungal agent for the prevention of chytridiomycosis (Stockwell et al. 2012). Runoff pollution from deicing salt could be a factor affecting wetlands in some regions (Sanso and Hecnar 2006; Karraker and Ruthig 2009), however, in our study region in the southern United States, deicing salt is not commonly used, therefore, it likely had no effect on the results of this study. Urban pollution may be a contributing factor in reducing the incidence of *Bd* in some cases but, to date, we have no evidence to support this assumption.

Density dependence and species richness could be other factors influencing the occurrence of *Bd* (Rachowics and Briggs 2007; Olson et al. 2013). We had a much higher incidence of *Bd* in forested sites and possibly higher densities or frogs compared to urban sites. In addition, species richness is likely to be higher in forested habitats as well (Saenz, personal observations). While our study was not designed to account for effect of host density and community complexity on the frequency of *Bd*, we cannot rule these out as factors and should be considered in future research.

While pollution and host density dependence are factors that warrant more investigation, temperature and moisture are well-accepted correlates of *Bd* infection rates (Schlaepfer et al. 2007; Murphy et al. 2009). Whitfield et al. (2012) found that despite small changes in temperature, *Bd* was much more prevalent in the cooler months, suggesting that small fluctuations in temperature may significantly affect *Bd* occurrence. Chatfield and Richards-Zawacki (2011) provided further evidence for the potential importance of temperature by using heat therapy to clear *Bd* from captive frogs. Hossack et al. (2013) found that boreal toads sampled in recently burned areas had a lower incidence of *Bd*, potentially due to a drier, warmer microclimate. This supports previous work implicating not only temperature, but moisture as an important factor influencing *Bd* infection rates as well (Murphy et al. 2009, 2011). Although we did not collect temperature and moisture data directly from the urban and forested sites, based on existing literature (Kim 1992), we expect city temperatures to be higher, and moisture levels lower in urban settings compared to forested sites; which would be consistent with the lower incidence of *Bd* observed in our study.

Furthermore, the effects of natural vegetation on microclimates are likely important in the context of this study. Raffel et al. (2010) suggest that shade may play an important role in *Bd* infection by lowering temperature. Becker et al. (2012) found that canopy density, natural vegetation, and daily average water temperature were the best predictors of *Bd* and differences in microclimate. Kim (1992) suggests that urban heating is attributed to rapidly heating urban surfaces, such as bare soil and short grasses, (see Fig. 2) which are less common in forested habitats and could lead to differences in the occurrence of *Bd* (Berger et al. 2004). We suggest that differences in temperature, generally resulting from land-use change are likely driving *Bd* occurrence rates, but we cannot rule out other factors at this point.

Although our study does not address the proximate causes of the observed differences in the occurrence rates, human-caused disturbances appear to have a clear effect on the incidence of *Bd*. We can conclude that, in our study in a temperate region, an inverse relationship with *Bd* and habitat disturbance does occur, as was observed in the tropics in previous studies (Van Sluys and Hero 2009; Becker and Zamudio 2011). Among human activities that cause habitat loss, urbanization produces some of the greatest extinction rates and eliminates a large majority of native species (McKinney 2002). However, our study adds to the paradox that disturbed habitat, including urban areas, may be acting as refugia from diseases, such as *Bd*, when species are able to tolerate urban environments and deforestation, as was suggested by Becker and Zamudio (2011).

Acknowledgments We thank Nancy Koerth, Dede Olson, and Cory Adams for constructive comments on an earlier draft of this manuscript.

References

Alford RA, Richards SJ (1999) Global amphibian declines: a problem in applied ecology. *Ann Rev Ecol Syst* 30: 133–165

- Becker CG, Zamudio KR (2011) Tropical amphibian populations experience higher disease risk in natural habitats. *Proc Natl Acad Sci U S A* 108:9893–9898
- Becker CG, Rodriguez D, Longo AV, Talaba AL, Zamudio KR (2012) Disease risk in temperate amphibian populations is higher at closed-canopy sites. *PLoS ONE* 7(10):e48205. doi:10.1371/journal.pone.0048205
- Berger L, Speare R, Daszak PD, Green ED, Cunningham AA, Goggin CL, Slocombe R, Ragan MA, Hyatt AD, McDonald KR, Hines HB, Lips KR, Marantelli G, Parkes H (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proc Natl Acad Sci U S A* 95:9031–9036
- Berger L, Speare R, Hines HB, Marantelli G, Hyatt AD, McDonald KR, Skerratt LF, Olsen V, Clarke JM, Gillespie G, Mahony M, Sheppard N, Williams C, Tyler MJ (2004) Effect of season and temperature on mortality in amphibians due to chytridiomycosis. *Aust Vet J* 82:31–36
- Berger L, Speare R, Marantelli G, Skerratt LF (2009) A zoospore inhibition technique to evaluate the activity of antifungal compounds against *Batrachochytrium dendrobatidis* and successful treatment of experimentally infected green tree frogs (*Litoria caerulea*) by fluconazole and benzalkonium chloride. *Res Vet Sci* 87:106–110
- Brem FMR, Lips KR (2008) *Batrachochytrium dendrobatidis* infection patterns among Panamanian amphibian species, habitats, and elevations during epizootic and enzootic stages. *Dis Aquat Organ* 81:189–202
- Chatfield MWH, Richards-Zawacki CL (2011) Elevated temperature as a treatment for *Batrachochytrium dendrobatidis* infection in captive frogs. *Dis Aquat Organ* 94:235–238
- Collins JP, Storfer A (2003) Global amphibian decline: sorting the hypotheses. *Divers Distrib* 9:89–98
- Daszak P, Cunningham AA, Hyatt AD (2003) Infectious disease and amphibian population declines. *Divers Distrib* 9:141–150
- Faulkner S (2004) Urbanization impacts on the structure and function of forested wetlands. *Urban Ecosyst* 7:89–106
- Fisher MC, Garner TWJ, Walker SF (2009) Global emergence of *Batrachochytrium dendrobatidis* and chytridiomycosis in space, time, and host. *Annu Rev Microbiol* 63:291–310
- Hamer AJ, McDonnell MJ (2008) Amphibian ecology and conservation in the urbanizing world: a review. *Biol Conserv* 141:2432–2449
- Hoffmann M et al (2010) The impact of conservation on the status of the World's vertebrates. *Science* 330:1503–1509
- Hossack BR, Lowe HW, Ware JL, Corn PS (2013) Disease in a dynamic landscape: host behavior and wildfire reduce amphibian chytrid infection. *Biol Conserv* 157:293–299
- Houlahan JE, Findlay CS, Schmidt BR, Meyer AH, Kuzmin SL (2000) Quantitative evidence for global amphibian population declines. *Nature* 404:752–755
- Johnson ML, Berger L, Philips L, Speare R (2003) Fungicidal effects of chemical disinfectants, UV light, desiccation and heat on the amphibian chytrid *Batrachochytrium dendrobatidis*. *Dis Aquat Organ* 57:255–260
- Karraker NE, Ruthig GR (2009) Effect of road deicing salt on the susceptibility of amphibian embryos to infection to water molds. *Environ Res* 109:40–45
- Kilpatrick AM, Briggs CJ, Daszak P (2010) The ecology and impact of chytridiomycosis: an emerging disease of amphibians. *Trends Ecol Evol* 25:109–118
- Kim HH (1992) Urban heat island. *Int J Remote Sens* 13:2319–2336
- Kinney VC, Heemeyer JL, Pessier AP, Lannoo MJ (2011) Seasonal pattern of *Batrachochytrium dendrobatidis* infection and mortality in *Lithobates areolatus*: affirmation of Vredenburg's "10,000 zoospore rule". *PLoS ONE* 6(3):e16708. doi:10.1371/journal.pone.0016708
- Kruger KM, Hero JM (2008) Altitudinal distribution of chytrid (*Batrachochytrium dendrobatidis*) infection in subtropical Australian frogs. *Aust Ecol* 33:1022–1032
- Kruger KM, Peregoglou F, Hero JM (2007) Latitudinal variation in the prevalence and intensity of chytrid (*Batrachochytrium dendrobatidis*) infection in eastern Australia. *Conserv Biol* 5:1280–1290
- Lane A, Burgin S (2008) Comparison of frog assemblages between urban and non-urban habitats in the upper Blue Mountains of Australia. *Freshw Biol* 53:2484–2493
- Lannoo MJ, Petersen C, Lovich RE, Nanjappa P, Phillips C, Mitchell JC, Macallister I (2011) Do frogs get their kicks on Route 66? Continental U.S. transect reveals spatial and temporal patterns of *Batrachochytrium dendrobatidis* infection. *PLoS ONE* 6(7):e22211
- Lips KR, Brem F, Brenes R, Reeve JD, Alford RA, Voyles J, Carey C, Livo L, Pessier AP, Collins JP (2006) Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proc Natl Acad Sci U S A* 103:3165–3170
- Longcore JE, Pessier AP, Nichols DK (1999) *Batrachochytrium dendrobatidis* gen et sp nov, a chytrid pathogenic to amphibians. *Mycologia* 91:219–227
- McKinney ML (2002) Urbanization, biodiversity, and conservation. *Bioscience* 52:883–890
- McKinney ML (2008) Effects of urbanization on species richness: a review of plants and animals. *Urban Ecosyst* 11:161–176

- Murphy PJ, St-Hilaire S, Bruer S, Corn PS, Peterson CR (2009) Distribution and pathogenicity of *Batrachochytrium dendrobatidis* in boreal toads from the Grand Teton area of western Wyoming. *EcoHealth* 6:109–120
- Murphy PJ, St-Hilaire S, Corn PS (2011) Temperature, hydric environment, and prior pathogen exposure alter the experimental severity of chytridiomycosis in boreal toads. *Dis Aquat Organ* 95:31–42
- Muths E, Pilliod DS, Livo LJ (2008) Distribution and environmental limitations of an amphibian pathogen in the Rocky Mountains, USA. *Biol Conserv* 141:1484–1492
- Naeem S, Thompson LJ, Lawler SP, Lawton JH, Woodfin RM (1994) Declining biodiversity can alter the performance of ecosystems. *Nature* 368:734–737
- Olson DH, Aanensen DM, Ronnenberg KL, Powell CI, Walker SF, Bielby J, Garner TWJ, Weaver G, Group TBM, Fisher MC (2013) Mapping the global emergence of *Batrachochytrium dendrobatidis*, the amphibian chytrid fungus. *PLoS ONE* 8(2):e56802. doi:10.1371/journal.pone.0056802
- Parris MJ, Baud DR (2004) Interactive effects of a heavy metal and chytridiomycosis on gray treefrog larvae (*Hyla chrysoscelis*). *Copeia* 2004:344–350
- Pauza MD, Driessen MM, Skerratt LF (2010) Distribution and risk factors for spread of amphibian chytrid fungus *Batrachochytrium dendrobatidis* in the Tasmanian Wilderness World Heritage Area, Australia. *Dis Aquat Organ* 92:193–199
- Pechmann JHK, Wilbur HM (1994) Putting declining amphibian populations in perspective: natural fluctuations and human impacts. *Herpetologica* 50:65–84
- Rachowics LJ, Briggs CJ (2007) Quantifying the disease transmission function: effects of density on *Batrachochytrium dendrobatidis* transmission in the mountain yellow-legged frog, *Rana mucosa*. *J Anim Ecol* 76:711–721
- Raffel TR, Michel PJ, Sites EW, Rhor JR (2010) What drives chytrid infections in newt populations? Association with substrate, temperature, and shade. *EcoHealth* 7:526–536
- Rodder D, Veith M, Lotters S (2008) Environmental gradients explaining the prevalence and intensity of infection with the amphibian chytrid fungus: the host's perspective. *Anim Conserv* 11:513–517
- Saenz D, Fitzgerald LA, Baum KA, Conner RN (2006) Abiotic correlates of anuran calling phenology: the importance of rain, temperature, and season. *Herpetol Monogr* 20:64–82
- Saenz D, Kavanaugh BT, Kwiatkowski MA (2010) *Batrachochytrium dendrobatidis* detected in amphibians from National Forests in Eastern Texas, USA. *Herpetol Rev* 41:47–49
- Sanso D, Hecnar SJ (2006) Effects of de-icing salt (NaCl) on larval wood frogs (*Rana sylvatica*). *Environ Pollut* 140:247–256
- Schlaepfer MA, Sredl MJ, Rosen PC, Ryan MJ (2007) High prevalence of *Batrachochytrium dendrobatidis* in wild populations of lowland leopard frogs *Rana yavapaiensis* in Arizona. *EcoHealth* 4:421–427
- Schloegel LM, Daszak P, Cunningham AA, Speare R, Hill B (2010) Two amphibian diseases, chytridiomycosis and ranaviral disease, are now globally notifiable to the World Organization for Animal Health (OIE): an assessment. *Dis Aquat Organ*. doi:10.3354/dao02140
- Skerratt LF, Berger L, Speare R, Cashins S, McDonald KR, Phillott DA, Hines HB, Kenyon N (2007) Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* 4:125–136
- Soule ME (1991) Conservation: tactics for a constant crisis. *Science* 253:744–750
- Stockwell MP, Clulow J, Mahony MJ (2012) Sodium chloride inhibits the growth and infective capacity of the amphibian chytrid fungus and increases host survival rates. *PLoS ONE* 7(5):e36942. doi:10.1371/journal.pone.0036942
- Stuart SN, Chanson JS, Cox NA, Young BE, Rodrigues ASL, Fischman DL, Waller RW (2004) Status and trends of amphibian declines and extinctions worldwide. *Science* 306:1783–1786
- Van Sluys M, Hero J-M (2009) How does chytrid infection vary among habitats? The case of *Litoria wilcoxii* (Anura, Hylidae) in SE Queensland, Australia. *EcoHealth* 6:576–583
- Wake DB (1991) Declining amphibian populations. *Science* 253:860
- Whitfield SM, Kerby J, Gentry LR, Donnelly MA (2012) Temporal variation in infection prevalence by the amphibian chytrid fungus in three species of frogs at La Selva, Costa Rica. *Biotropica* 44:779–784
- Woodhams DC, Alford RA (2005) Ecology of chytridiomycosis in rainforest stream frog assemblages of tropical Queensland. *Conserv Biol* 19:1449–1459
- Woodhams DC, Geiger CC, Reinert LK, Rollins-Smith LA, Lam B, Harris RN, Briggs CJ, Vredenburg VT, Voyles J (2012) Treatment of amphibians infected with chytrid fungus: learning from failed trials with itraconazole, antimicrobial peptides, bacteria, and heat therapy. *Dis Aquat Organ* 98:11–25