



Water yield following forest–grass–forest transitions

Katherine J. Elliott¹, Peter V. Caldwell¹, Steven T. Brantley², Chelcy F. Miniati¹, James M. Vose³, and Wayne T. Swank¹

¹USDA Forest Service, Southern Research Station, Coweeta Hydrologic Lab, Otto, NC 28763, USA

²Joseph W. Jones Ecological Research Center, Ichauway, Newton, GA 31770, USA

³USDA Forest Service, Southern Research Station, Center for Integrated Forest Science, Raleigh, NC 27695, USA

Correspondence to: Katherine J. Elliott (kelliott@fs.fed.us)

Received: 18 October 2016 – Discussion started: 24 October 2016

Accepted: 31 January 2017 – Published: 16 February 2017

Abstract. Many currently forested areas in the southern Appalachians were harvested in the early 1900s and cleared for agriculture or pasture, but have since been abandoned and reverted to forest (old-field succession). Land-use and land-cover changes such as these may have altered the timing and quantity of water yield (Q). We examined 80 years of streamflow and vegetation data in an experimental watershed that underwent forest–grass–forest conversion (i.e., old-field succession treatment). We hypothesized that changes in forest species composition and water use would largely explain long-term changes in Q . Aboveground biomass was comparable among watersheds before the treatment (208.3 Mg ha^{-1}), and again after 45 years of forest regeneration (217.9 Mg ha^{-1}). However, management practices in the treatment watershed altered resulting species composition compared to the reference watershed. Evapotranspiration (ET) and Q in the treatment watershed recovered to pre-treatment levels after 9 years of abandonment, then Q became less (averaging 5.4 % less) and ET more (averaging 4.5 % more) than expected after the 10th year up to the present day. We demonstrate that the decline in Q and corresponding increase in ET could be explained by the shift in major forest species from predominantly *Quercus* and *Carya* before treatment to predominantly *Liriodendron* and *Acer* through old-field succession. The annual change in Q can be attributed to changes in seasonal Q . The greatest management effect on monthly Q occurred during the wettest (i.e., above median Q) growing-season months, when Q was significantly lower than expected. In the dormant season, monthly Q was higher than expected during the wettest months.

1 Introduction

Forests play a critical role in regulating hydrological processes in headwater catchments by moderating the timing and magnitude of streamflow (Burt and Swank, 2002; Chang, 2003; Ice and Stednick, 2004; Ford et al., 2011b; Vose et al., 2011). Hydrological processes in forests are particularly sensitive to disturbances that reduce tree vigor or leaf area and thus decrease evapotranspiration (ET) (Aranda et al., 2012; Edburg et al., 2012; Brantley et al., 2013). Most efforts at studying the effects of disturbance on watershed hydrology have focused on quantifying the effects of forest harvesting practices on water yield (Q) (Bosch and Hewlett, 1982; Stednick, 1996; Burton, 1997; Brown et al., 2005; Wei and Zhang, 2010; Ford et al., 2011a; Zhang and Wei, 2012; Liu et al., 2015). Reviews have shown that, in general, harvesting < 20 % of the basal area shows no detectable increase in annual Q ; but Q increases thereafter as the percentage of basal area harvested increases (Bosch and Hewlett, 1982; Andréassian, 2004; Brown et al., 2005). However, recent work that aims to merge ecology of the resulting forest and species composition with traditional hydrology approaches (i.e., ecohydrology) has advanced our understanding greatly. For example, Brantley et al. (2013, 2015) showed that lasting changes in annual Q (lower) and persistent changes in peak flow (> 20 %, after the most intense storms) were observed with only about a 5 % basal area loss of eastern hemlock (*Tsuga canadensis* (L.) Carrière). Brantley et al. (2013) also suggested that a change in forest composition with less evergreen hemlock relative to deciduous trees could result in an increase in Q in winter months.

Most of the eastern US forests have been harvested at least once since the late 1800s (Yarnell, 1998; Foster et al., 2003; Thompson et al., 2013; Martinuzzi et al., 2015), and many forested areas have undergone forest-to-agriculture land-use changes, and then been abandoned to revert back to forest (i.e., abandoned agriculture or old-field succession) (Otto, 1983; Trimble et al., 1987; Wear and Bolstad, 1998; Belle-mare et al., 2002; Alvarez, 2007; Thiemann et al., 2009; Ramankutty et al., 2010; Kirk et al., 2012). Land abandonment has also been prevalent and ongoing since the early 20th century in other countries (Cramer et al., 2008; García-Ruiz and Lana-Renault, 2011). Land-use and land-cover (LULC) changes, such as forest–grass–forest transitions, may alter the timing and quantity of Q . Because land-use conversion from forests to agriculture often includes a combination of changes in vegetation composition and soil physical attributes, it is difficult to separate the effects of vegetation changes from soil changes (see reviews by Neary et al., 2009; Zimmermann et al., 2010; Houlbrooke and Laurenson, 2013; Morris and Jackson, 2016). Land-cover conversion that requires heavy machinery or includes livestock grazing decreases soil infiltration and saturated hydraulic conductivity (e.g., Hassler et al., 2011; Price et al., 2011; Morris and Jackson, 2016), and can thus increase peak flow during storms, and flood frequency and severity (Reinhart, 1964; Hornbeck, 1973; Burt and Swank, 2002; Alila et al., 2009; Green and Alila, 2012). Without soil compaction and alteration of water flow pathways, forest trees typically use more water and extract water from deeper soil than shallower-rooted grasses (Zhang et al., 2001; Kulmatiski and Beard, 2013), which could result in higher ET and lower Q at the catchment scale.

Several studies have compared Q and ET of forests and pastures. Analyzing 250 catchments worldwide, Zhang et al. (2001) found that forested catchments had higher ET than grass pastures, with few exceptions. Replacing trees with grass cover generally increases Q by decreasing ET (Hibbert, 1969; Bosch and Hewlett, 1982; Farley et al., 2005), although not always (Brauman et al., 2012; Amatya and Harrison, 2016). In some basins, when agricultural land use is reduced and forest cover increased, Q is unchanged, and can be explained in part by the species-specific traits in water use (e.g., deciduous versus evergreen, and/or late season versus early season perennial grass) (Cruise et al., 2010), and the geomorphological differences among biomes (Zhou et al., 2015).

Large differences among tree species in their leaf and canopy conductance, transpiration per unit leaf area, and whole-tree water use for any given diameter exist in eastern US deciduous forests (Wullschleger et al., 2001; Ford and Vose, 2007; Ford et al., 2011a). This is especially true when comparing hardwoods within diffuse-porous and ring-porous xylem functional groups (Taneda and Sperry, 2008; Ford et al., 2011a; Von Allmen et al., 2015). *Liriodendron tulipifera*, a diffuse-porous species common to the eastern temperate deciduous forest biome, has among the highest transpiration

rates of forest trees; while rates of *Acer rubrum* L. and *Betula lenta* L., also common diffuse-porous species, are lower than *L. tulipifera*. However, they have relatively high transpiration rates compared to common ring-porous *Quercus* species (Ford et al., 2011a).

Few studies have examined long-term changes in catchment hydrology through a forest–grass–forest transition, with specific attention focused on species compositional changes and their effect on ET and Q . A treated watershed within the Coweeta Hydrologic Laboratory, western North Carolina, experienced this LULC transition, and reported similar Q between forest and grass when the grass cover was well fertilized (Hewlett and Hibbert, 1961; Hibbert, 1969; Bosch and Hewlett, 1982; Burt and Swank, 1992). However, they did not investigate why Q was lower than expected after grass-cover abandonment and through the early successional development of the deciduous forest. Road construction could be a contributing factor because installing temporary roads to facilitate timber harvesting can affect hydrology (Harr et al., 1975; Alila et al., 2009), but only 3.3 % of the watershed area was in temporary roads (inactive for the least 50 years). In addition, roads comprising less than 6 % of the watershed area appear not to change storm hydrographs significantly (Harr et al., 1975; Swank et al., 2001; Alila et al., 2009). In a more recent study, Ford et al. (2011b) suggested that the decline in Q over time could be due to a shift in the dominant tree species in the treated, old-field succession watershed.

Working in the same experimental watershed as authors above, we compared the long-term changes (1934–2015) in the following: (1) aboveground biomass, leaf area index (LAI) and species and functional (xylem anatomy) group composition; (2) estimated growing-season mean daily water use (DWU); (3) annual water-balance derived ET; and (4) daily, monthly, and annual Q between the treated, old-field succession, watershed (WS6) and nearby reference (WS14, WS18) watersheds with an emphasis on the period of reforestation. We hypothesized that (1) a shift in species composition and their attendant DWU will largely explain long-term changes in Q , (2) annual Q would be lower in the treated WS6 through forest succession concurrent with greater DWU with additional changes in timing of Q due to altered species composition, and (3) monthly Q would be greater in the treated WS6 for wet periods (high or peak flows), and this effect would be greatest in the dormant season.

2 Materials and methods

2.1 Study area

The study was conducted at the USDA Forest Service Coweeta Hydrologic Laboratory, a 2185 ha forested basin in the Nantahala National Forest in North Carolina, USA




Figure 1. Map of the Coweeta Basin with treated WS6 and reference (WS14, WS18) watersheds with rain gauges, weirs, and vegetation plots.

(35°06' N, 83°43' W). Climate in the Coweeta Basin is classified as marine, humid temperate (Swift et al., 1988), with mild temperature (average 12.8 °C) and ample precipitation (average 1795 mm yr⁻¹). Three watersheds (WS6, WS14 and WS18) within 1 km of one another, similar in elevation, slope, and aspect were used in this study (Table 1). WS14 and WS18 are untreated reference watersheds (Fig. 1). WS6 experienced a disturbance regime similar to forest conversion to pasture and subsequent abandonment common across the region (see below). Soils in all watersheds are moderately permeable, well-drained, moderately deep to very deep, and with a saprolite layer up to 6 m deep (Thomas, 1996).

2.2 History of disturbance

Before 1842, the Coweeta Basin was burned semiannually (Douglass and Hoover, 1988). Between 1842 and 1900, light semiannual burning and grazing continued. From 1912 to 1923 heavy logging occurred (Douglass and Hoover, 1988). Loss of American chestnut (*Castanea dentata* (Marshall) Borkh.) in the 1930s (Woods and Shanks, 1959; Elliott and Swank, 2008) was followed by loss of *Tsuga canadensis* (L.) Carrière in the 2000s (Elliott and Vose, 2011).

The disturbance regime in WS6, the treated watershed, was extensive (Table S1 in the Supplement). In July 1941, 12% of the catchment (1.06 ha area) along the stream was cut to determine how riparian vegetation affects *Q* (Dunford and Fletcher, 1947). In 1958, the entire watershed was clear-cut, merchantable timber was removed, and the residue was piled and burned. In 1959, surface soil was scarified and seeded to *Festuca octoflora* grass. In 1960, the watershed was treated with a one-time application of 1100 kg ha⁻¹ lime, 110 kg ha⁻¹ 30-10-10 NPK, and

Table 1. Characteristics of treated (WS6) and reference (WS14, WS18) watersheds at the Coweeta Hydrologic Laboratory in Otto, NC, USA. Mean annual precipitation (*P*) and water yield (*Q*) based on data collected over water years (WY, May–April) from 1934 to 2015.

Characteristic	Units	Watersheds		
		6	14	18
Area	ha	9.0	61.03	12.46
Mean elevation	m	824	878	823
Mean basin slope	%	50	50	55
Aspect		NW	NW	NW
Year of first complete flow record	WY	1939	1938	1938
Nearest rain gauge		SRG41	SRG41	SRG96
Mean precipitation (<i>P</i>)	mm yr ⁻¹	1843	1843	2031
Mean water yield (<i>Q</i>)	mm yr ⁻¹	866	997	1021
Mean evapotranspiration (ET = <i>P</i> – <i>Q</i>)	mm yr ⁻¹	978	845	1010
<i>Q/P</i>		0.47	0.54	0.50

18.4 kg ha⁻¹ granular 60% potash. Between 1960 and 1965, *Kalmia latifolia* L., *Rhododendron maximum* L., and other hardwood sprouts were suppressed with spot applications of 2,4D [(2,4dichlorophenoxy) acetic acid] to maintain the watershed in grass cover (Hibbert, 1969). In 1965, the watershed was fertilized again, as above. In 1967, the grass was treated with the herbicides atrazine [2-chloro-(4-ethylamino)-6-9-isopropylamino]-Strizine], paraquat [1,1 dimethyl 4,4 bipyridinium ion (dichloride salt)], and 2,4D [(2,4dichlorophenoxy) acetic acid] (Douglass et al., 1969), and then left undisturbed (hereafter, old-field succession). Although the grass was not cut or grazed, the lime and fer-

tilizer amendments with attendant high productivity and nutrient uptake by the grass make these applications somewhat similar to agricultural practices. The original objectives of the conversion from forest-to-grass were to compare water use of grass versus hardwoods (Hibbert, 1969; Swank and Crossley, 1988) and to determine how conversion to grass affects discharge characteristics (Burt and Swank, 1992).

Two adjacent forested watersheds (WS14, WS18) were selected as references to provide an adequate number of sample plots (described below) for analysis of changes in vegetation. These reference watersheds with similar physiography (Table 1) are characteristic of mature, second-growth hardwood forests, and have remained unmanaged since 1923 (Swank and Crossley, 1988). We considered forest age for WS14 and WS18 since the loss of *C. dentata* presently to be > 75 years old.

2.3 Measurements

2.3.1 Vegetation

The relative importance of woody species over time was characterized with repeated tree surveys. In treated WS6, surveys were conducted in 1934, 1958, 1982, 1995, and 2012. In 1934, only five 0.08 ha permanent plots were measured along the east side of the watershed; in all subsequent surveys, plots were placed across the entire watershed. In 1958, a pretreatment strip inventory sampled 25 % of the watershed area with 10 m wide strips approximately 40 m apart extending along transects from the ridge-top to the stream channel. This sampling method resulted in a total of 37 unequally sized plots (ranging from 0.02 to 0.14 ha), including the riparian corridor. In 1982, 34 0.02 ha plots were permanently marked continuously along five transects from ridge-top to near stream; these 34 plots were re-measured in 1995 and 2012.

In reference WS14, 31 0.08 ha permanent plots were surveyed in 1934, 1969, 1988–1992 (hereafter, 1992), and 2009–2010 (hereafter, 2009). In reference WS18, eight 0.08 ha permanent plots were surveyed in 1934, 1953, 1969, 1992, and 2009.

In all watersheds and for all survey periods, diameter of woody stems ≥ 2.54 cm at breast height (DBH, 1.37 m above ground) was measured by species and recorded into 2.54 cm DBH classes. In 1934, only percent cover was recorded for the two evergreen shrubs, *Rhododendron maximum* and *Kalmia latifolia*; for this reason, we do not estimate biomass and LAI (square meters of leaf area per square meter of ground area) for these species in 1934. In all other years, stem diameters of these evergreen shrubs were measured in the same manner as the tree diameters. Median DBH values were used to calculate basal area, aboveground biomass, leaf biomass, and LAI. We used species-specific allometric equations developed on-site to estimate the aboveground biomass, leaf biomass, and LAI contribution of each species in each

watershed (McGinty, 1972; Santee and Monk, 1981; Martin et al., 1998; Ford and Vose, 2007; B. D. Kloeppel, unpublished data; C. F. Miniati, unpublished data). Species nomenclature follows Kirkman et al. (2007).

2.3.2 Water yield (Q) and evapotranspiration (ET)

We used both chronological-pairing (i.e., corresponding to the same meteorological input) and frequency-pairing (described below) analyses to detect potential hydrologic responses of Q and ET to LULC change. Both analyses used the paired watershed approach (Wilm, 1944, 1949). The chronological pairing approach allowed us to create a time series of estimated change in annual Q and ET over the period of record and to relate these changes to both the treatment and climate. In addition, this analysis allowed us to determine when a consistent change in Q began, enabling us to establish the time period of interest for the frequency pairing. The frequency-pairing approach allowed us to compare the post-treatment distribution of monthly and annual Q to that of the pretreatment period. We used WS18 and WS6 as the reference and treatment watersheds, respectively. We did not compare WS6 to WS14 because there were gaps in the WS14 flow record in the years immediately following the grass conversion and herbicide application. For both watersheds, a 5 min stream stage was used to estimate Q (Reinhard and Pierce, 1964; Swift et al., 1988).

We modeled WS6 annual Q and ET as a function of WS18, incorporating the effect of grass conversion and reforestation treatments over time. Annual Q was computed on a May–April water year to minimize the effects of year-to-year changes in storage, as soils are generally at their wettest by the beginning of May. The empirical chronological-pairing model was fit using PROC NLIN (SASv9.4, SAS Institute, Cary, NC) and had the following form:

$$\hat{Q}_T = a + bQ_R + eM_1t_1 + \left[M_2c \left(h - \frac{1}{1 + \exp^{-t_2}} \right) \right], \quad (1)$$

where \hat{Q}_T = predicted Q from treated watershed WS6 (mm yr^{-1}), Q_R = measured Q from reference watershed WS18 (mm yr^{-1}), M_1 = management representing grass conversion, ($M_1 = 1$ for water years between and including 1960 and 1966; $M_1 = 0$ otherwise), t_1 = time since grass fertilization ($t_1 = \text{water year} - \text{fertilization year}$ for water years between and including 1960 and 1966 where fertilization years include water years 1959, 1961, and 1966; $t_1 = 0$ otherwise), M_2 = management representing reforestation after grass conversion ($M_2 = 1$ for water years greater than or equal to 1967; $M_2 = 0$ otherwise), and t_2 = time since reforestation after grass conversion ($t_2 = \text{water year} - 1967$ for water years greater than or equal to 1967; $t_2 = 0$ otherwise); a , b , c , e , and h are fitted parameters.

This overall modeling approach has been used in prior studies to assess the impact of forest management on Q (Ford et al., 2011b; Kelly et al., 2016). The $a + bQ_R$ term

in EQ1 reflects the relationship between reference and treatment watersheds assuming no treatment. The increasing linear Q response (eM_1t_1 term in EQ1) accounts for the decline in annual grass production after fertilization and water use, as noted by Hibbert (1969). The $M_2c(h - \frac{1}{1+\exp^{-t_2}})$ term in EQ1 accounts for the exponential decline in Q as the forest regenerates, which has been observed in numerous paired watershed experiments (Swank et al., 1988).

As in Ford et al. (2011b), we define the Q treatment response, D_Q , as the difference between the observed Q in the treated watershed (Q_T) and that predicted by the model assuming no treatments had taken place (\hat{Q}_T):

$$D_Q = Q_T - \left(\hat{Q}_T; M_1, M_2 = 0\right). \quad (2)$$

The proportion of the variability explained by the model was quantified using the ratios of the error-to-total sum of squares and the total-to-error degrees of freedom as follows:

$$R_{\text{adjusted}}^2 = 1 - \frac{\text{SSE}}{\text{SST}} \times \frac{\text{df}_T}{\text{df}_E}. \quad (3)$$

Parameter estimates were interpreted as statistically significant at $\alpha = 0.05$. Observed annual ET was computed as precipitation (P) $- Q_T$ while expected ET with no treatment was computed as $P - \hat{Q}_T$, both assuming the largely impermeable bedrock underlying the basin that results in negligible deep groundwater losses (Douglass and Swank, 1972). Watershed P was estimated using a nearby 8 in. (20.3 cm) National Weather Service standard rain gauge, SRG 96 (Laseter et al., 2012). The ET treatment response, D_{ET} , is then as follows:

$$D_{\text{ET}} = [P - Q_T] - \left([P - \hat{Q}_T]; M_1, M_2 = 0\right). \quad (4)$$

2.3.3 Frequency-pairing flow distributions

We used the frequency-pairing method (Alila et al., 2009; Brantley et al., 2015) to detect differences in frequency between observed and predicted annual and monthly Q after treatment. Briefly, frequency pairing is an analytical method that quantifies differences in observed and predicted Q parameters based on the probability of occurrence of a given Q (or flow at a given probability) rather than based on occurrence at a discrete time (i.e., chronological pairing). This accounts for rainfall amount and antecedent soil conditions. We used pre-treatment Q during water years 1939–1941, 1948–1951, and 1956–1958, to estimate the expected cumulative distribution functions (CDFs, F_Y) for observed and predicted Q in the treatment watershed using the following linear regression equation:

$$\hat{Y}_i = b_0 + b_1 X_i, \quad (5)$$

where X_i is the observed Q in the reference watershed for period i (day of year) and \hat{Y}_i is the expected Q for the

treatment watershed under undisturbed conditions for the same period. We used PROCMODEL (SAS v9.3, SAS Institute, Cary, NC) to predict monthly post-treatment Q in the treatment watershed from May 1979 to April 2015 and annual post-treatment Q for water years 1980–2015. To model monthly Q , we separated the data by calendar month and created twelve separate regression equations. Using separate regression equations for each month helped account for variations in paired watershed Q relationships among months and helped to distinguish differences in effects among seasons.

Observed and predicted Q values were then plotted as an estimate of the probability of occurrence for ranked event $Y_{(i)}$ during any time period i . The exceedance probability, $1 - p$, was estimated for each period using the following equation:

$$1 - F_Y \left[\hat{Y}_i\right] = \frac{m - 0.40}{n + 0.20}, \quad (6)$$

where m is the rank for a given flow and n is the total number of flow periods in the distribution. This function provided an empirical estimate of the quantile for a given flow value (Cunnane, 1978; Stedinger et al., 1993). Confidence limits for each predicted flow at each probability of occurrence were estimated as follows:

$$Y_m \pm z_{1-\frac{\alpha}{2}} \sqrt{(\text{Var}_1[Y_m] + \text{Var}_2[Y_m])}. \quad (7)$$

We used a pair of Monte Carlo simulations to estimate the variability associated with the predictive uncertainty in Eq. (7) (Var_1), and the uncertainty associated with the sampling variability at each rank (Var_2). For these analyses, we used 1000 iterations for each simulation. We used the raw, expected post-treatment values from Eq. (5) to correct for the loss of variability in the upper tails of the distribution (Alila et al., 2009). The CDFs were then used to construct flow duration curves to assess changes in untransformed Q at monthly and annual intervals by comparing the change in magnitude for a given probability or the change in probability for a given magnitude (Alila et al., 2009; Green and Alila, 2012; Brantley et al., 2015).

2.3.4 Growing-season daily water use (DWU)

Plant water loss was estimated by scaling up sap flux measurements of numerous species and stem diameter sizes at Coweeta Hydrologic Laboratory (Ford and Vose, 2007; Ford et al., 2011b; Brantley et al., 2013; Miniati, unpublished) using methods outlined in Ford et al. (2011a). We fitted the observed growing-season mean DWU (kg day^{-1}) to stem DBH (cm) using a power function of the following form:

$$\text{DWU} = b_0 \cdot \text{DBH}^{b_1}. \quad (8)$$

Species were grouped into xylem functional types (diffuse-porous, ring-porous, semi-ring porous, evergreen shrub, or tracheid) and growing-season DWU models were developed for each xylem functional type. For example, *Carya*

Table 2. Summary of growing-season daily water use (DWU, kg day⁻¹) models for each xylem functional group as a function of stem diameter at breast height (DBH, cm); $DWU = b_0 \cdot DBH_1^b$.

Xylem group	<i>N</i>	Min DBH (cm)	Max DBH (cm)	<i>b</i> ₀	<i>b</i> ₁	Adjusted <i>R</i> ²	SE of estimate
Diffuse-porous	95	7.4	61.8	0.1428	1.7676	0.70	30.3
Evergreen Shrub	16	5.3	16.3	0.6445	0.7002	0.00	2.5
Ring-porous	38	23.9	86.7	0.2392	1.1488	0.55	9.1
Semi-ring-porous	18	20.2	55.7	0.0009	2.8557	0.88	8.8
Tracheid	116	9.5	67.5	0.0005	2.8411	0.73	8.6

spp. have semi-ring porous xylem; *Quercus* spp. and *Oxydendrum arboreum* have ring-porous xylem; and *Betula lenta*, *Liriodendron tulipifera*, and *Acer rubrum* have diffuse-porous xylem (Table S2). Because *R. pseudoacacia* behaves more like a diffuse-porous species, its measured values of DWU and DBH were combined with the diffuse-porous model. Even though *Robinia pseudoacacia* has ring-porous xylem, it is isohydric (i.e., maintaining stable leaf water potentials as soil water potentials drop, Klein, 2014) and has higher DWU than *Quercus* or *Carya* (Miniat and Hubbard, unpublished). For the two understory evergreen species, *Kalmia latifolia* and *Rhododendron maximum*, we applied the mean DWU value from 16 instrumented shrubs because DWU models based on DBH alone provided limited predictive power (Table 2). We estimated growing-season mean-plot DWU by modeling DWU by functional type and vegetation surveys by diameter for all watersheds. We did not estimate DWU for the 1934 survey, when *C. dentata* was most abundant, because most of the trees had been affected by chestnut blight, compromising their functional xylem.

3 Results

3.1 Vegetation dynamics

Prior to treatment, species composition and aboveground biomass among the watershed were similar (Fig. 2). In 1934, aboveground biomass was comparable among the treated WS6 and references WS14 and WS18, averaging 200 Mg ha⁻¹ ($p = 0.706$) (Fig. 2a, Table S3). Biomass declined in WS6 (99.51 Mg ha⁻¹) from 1934 to 1958 prior to conversion to grass, and in WS18 (148.42 Mg ha⁻¹) from 1934 to 1953 (Table S3). The decline in biomass and LAI between 1934 and the 1950s was primarily due to the loss of *Castanea dentata* (Fig. 2a–c). In 1934, *C. dentata* occupied from 40 to 54 % of the biomass and 29–43 % of the LAI across the three watersheds (Fig. S1 in the Supplement).

The grass cover in the treated watershed was highly productive, but following the herbicide treatment (i.e., old-field succession), early successional vegetation rapidly established (Fig. S1a). During the 5 years when WS6 was main-




Figure 2. Mean (\pm SE bars) (a) aboveground biomass, (b) leaf biomass, and (c) leaf area index (LAI) for the treated WS6 and reference (WS14, WS18) watersheds over time.

tained in grass, biomass ranged from 5.67 to 7.30 Mg ha⁻¹. In 1968, 1 year after cessation of treatment, the aboveground biomass was 3.92 Mg ha⁻¹ in WS6. At that time, the 1-year-old field was dominated by *Erechtites hieracifolia* (L.) Raf., *Phytolacca americana* L., *Eupatorium* spp., and *Equisetum arvense* L. and had remnants of *Festuca octoflora*. In the years between 1968 and 1982, WS6 was rapidly colo-

nized by *Robinia pseudoacacia* and *Liriodendron tulipifera* (Fig. S1a), whereas the most abundant species in the reference watersheds in the years following the loss of *C. dentata* (1969 to 2010s) were *Quercus* spp. and *Acer rubrum* (Fig. S1b–c; Tables S4–S6).

Forest composition following grass cover was biased towards tree species with deep functional sapwood and diffuse-porous xylem. In 1934, all watersheds were dominated by species with semi-ring-porous (*C. dentata* and *Carya*) or ring-porous (*Quercus*) xylem, accounting for more than 80 % of the aboveground biomass (Fig. 3a–c) and 80 % of the LAI (Tables S4–S6). Although species with semi-ring-porous xylem declined in all watersheds over time, the increase in species with diffuse-porous xylem was greater in the treated watershed compared to reference watersheds (Fig. 3a–c). As the young forest developed following grass herbicide and abandonment, species with diffuse-porous xylem and *R. pseudoacacia* dominated forest biomass, while species with ring-porous xylem were only 2.7 %. By 2012, 93 % of vegetation in the treatment watershed was comprised of species with diffuse-porous xylem (Fig. 3a), while the reference watersheds were about half of the species with ring-porous xylem (Fig. 3b and c).

3.2 Water yield (Q) and evapotranspiration (ET)

The forest–grass–forest treatment of WS6 resulted in significant effects on Q over time. Models of annual Q explained more than 98 % of the variability in Q over the period of record. Initial harvesting increased Q by 99 mm (10.5 % above the expected Q) in 1960 (Fig. 4), and Q remained higher than expected during the grass conversion period except in 1959, 1961, and 1966 when grass production was highest due to fertilizer application. The largest treatment effect occurred in 1967 when herbicide was applied to the watershed, resulting in a Q increase of 259 mm (31 % above the expected Q) (Fig. 4). Q remained higher than expected for approximately 9 years after the herbicide treatment as the vegetation re-established. Beginning in 1977 and continuing through 2015, Q was less than expected in 32 of 35 years (Fig. 4), suggesting that the new forest used more water (i.e., had higher ET) than expected had it not undergone treatment. Since 1980, on average, annual Q decreased by 6.1 %, ranging from a Q increase in 1981 of 30 mm (+5.5 %) to a decrease of 142 mm (16 %) in 2003. ET increased by 4.5 % on average, relative to what was expected in the absence of management (Fig. S2).

3.3 Changes in flow distribution

In addition to the forest–grass–forest treatment changing the amount of Q , it also fundamentally changed the distribution of Q , with the most pronounced changes at the height of the growing and dormant seasons. The annual and monthly Q relationships between the reference and treatment




Figure 3. Percentage (\pm SE bars) of aboveground biomass for the xylem functional groups (diffuse-porous, ring-porous, semi-ring porous, tracheid, and evergreen shrub) in the (a) treated WS6, (b) reference WS14, and (c) reference WS18 over time.

watersheds for the pre-treatment period were highly significant (annual: $n = 10$, $r^2 = 0.97$, $p < 0.001$; monthly: $n = 10$, $r^2 = 0.94$, $p < 0.001$) using the frequency-pairing method. Annual Q was unchanged at low and high probabilities of non-exceedance ($< 10\%$), but was lower in some years between the 30 and 60 % probability of non-exceedance (Fig. 5a). Monthly Q was higher than expected at high probability of non-exceedance in February (Fig. 5b), whereas monthly Q was lower than expected at the high probability of non-exceedance in July (Fig. 5c). Median monthly Q was lower than expected for only January (−14.8 %) and May (−13.4 %) (Table 3). At wetter periods (above median Q), monthly Q was lower than expected for several months and immediately following the growing season




Figure 4. Changes in water yield (Q , $D = Q_T - (\hat{Q}_T; M_1, M_2 = 0)$) for the treated WS6 over time (bars). Solid lines are the standard errors of the mean prediction. We used the paired-watershed approach with WS18 as the reference. The year of harvest, conversion to Kentucky-31 fescue grass (*Festuca octoflora*) cover, fertilize, herbicide, and abandonment to allow forest regeneration are denoted by dashed lines.

(June–October, December; Table 3); whereas, for February–April, monthly Q was higher than expected. At drier periods (below median Q), February, March, and September had lower-than-expected monthly Q (Table 3). No significant changes in monthly Q distributions were observed in November.

3.4 Daily water use (DWU)

Growing-season DWU differed among species for any given DBH largely dependent on xylem anatomy (Table 2, Fig. 6). For example, DWU for a tree with 50 cm DBH could be 6.5 times higher with diffuse-porous xylem compared to ring-porous xylem (Fig. 5). *Liriodendron tulipifera*, *Betula lenta* and *Nyssa sylvatica* had the highest DWU; *Acer rubrum* and *Carya* were intermediate; and *Quercus alba*, *Q. montana*, and *Q. rubra* had the lowest estimated DWU compared to all other species (Ford et al., 2011b). *Robinia pseudoacacia* had higher DWU than *Quercus* or *Carya* (Miniat and Hubbard, unpublished). Models based on DBH and xylem anatomy explained 55–88 % of the variability in DWU among tree species (Table 2). For the evergreen understory species, however, DBH explained little variation in DWU, even though the standard errors were quite low.

Mean growing-season DWU for each catchment increased over time, but the treated watershed showed the greatest increase (Fig. 7a). In the 2010s, the 45-year-old forest in WS6 had 25–43 % higher DWU than the > 75-year-old reference forests (Fig. 7a–d), despite lower leaf area than the refer-

Table 3. Relative changes in monthly water yield (Q) for different parts of the cumulative distribution function for the period May 1979 to April 2015 for the treated WS6 using the frequency-pairing method. “Lower” and “Higher” denote direction and significance ($p < 0.05$) of change; NS = not significant.

Month	Change in median monthly Q (%)	Change in Q below the median Q	Change in Q above the median Q
Jan	Lower (−14.8)	NS	NS
Feb	NS	Lower	Higher
Mar	NS	Lower	Higher
Apr	NS	NS	Higher
May	Lower (−13.4)	NS	NS
Jun	NS	NS	Lower
Jul	NS	NS	Lower
Aug	NS	NS	Lower
Sep	NS	Lower	Lower
Oct	NS	NS	Lower
Nov	NS	NS	NS
Dec	NS	NS	Lower

ence watersheds at that time (Fig. 2c). In reference WS14, tree species with diffuse-porous xylem contributed 48–63 % of the total water use between 1969 (age 35) and 2009 (age 75+), while evergreen shrubs contributed 20–23 %, and tree species with ring-porous xylem contributed 13 % or less to the total water use (Fig. 7c). Since the grass cover was




Figure 5. Changes in the cumulative distribution function (CDF) expressed as historic probability of non-exceedance for (a) annual water yield (Q), and monthly Q for (b) February and (c) July. * ($p < 0.05$) and ** ($p < 0.01$) denote years in the distribution functions when Q was significantly lower or higher than predicted.

abandoned in WS6, tree species with diffuse-porous xylem alone have contributed more than 90 % of the total daily water use in that watershed (Fig. 7b).

4 Discussion

We hypothesized that a shift in species composition and the resulting shift in DWU would largely explain long-term changes in Q in the treated watershed as the forest regenerated following grass abandonment. We found that forest species composition in the treated watershed shifted from dominance by species with ring-porous xylem prior to grass conversion to species with diffuse-porous xylem through old-field succession. With this major shift in species composition, DWU increased from 1982 to 2012 in the treated watershed, and it was much higher than that in the older refer-




Figure 6. Growing-season daily water use of tree species by xylem functional group (diffuse-porous, ring-porous, semi-ring porous, evergreen shrub and tracheid) and DBH (diameter at 1.37 m above ground).

ence watersheds. These changes in species composition and DWU correspond with the long-term trend in lower-than-expected Q over that time period. Seasonal variation in Q helped to explain this long-term pattern.

4.1 Vegetation dynamics

Species composition has changed dramatically in the treated watershed through old-field succession following the forest–grass–forest transition. Prior to conversion to grass (1958), the forest was dominated by *Quercus montana* and *Q. coccinea*, similar to the reference watersheds at that time. After the grass was treated with herbicide, and the forest was allowed to reestablish, the forest shifted to dominance by *Liriodendron tulipifera* and *Robinia pseudoacacia*. Other studies have found that shade-intolerant *R. pseudoacacia* and *L. tulipifera* respond and grow rapidly following clear-cutting or other disturbances that create large canopy gaps (Elliott and Swank, 1994; Elliott et al., 1997, 1998; Shure et al., 2006; Boring et al., 2014). During grass dominance all woody species were eliminated with spot herbicide application. This treatment killed stump sprouts, and during forest succession recruitment favored small, wind-dispersed seeds, and discriminated against large-seeded and slow growing species such as *Quercus*, *Carya*, *Tilia*, and *Aesculus* (Elliott et al., 1997, 2002). Aboveground biomass approached pretreatment levels after 45 years of forest growth; however, LAI remained lower than that of the pretreatment or reference watershed conditions. The lower LAI could be attributed to the differences among species in the ratio of leaf area per total aboveground biomass and crown structure, where shade-intolerant *R. pseudoacacia* and *L. tulipifera* have lower ratios and concentrate foliage to the uppermost crown more than intermediate shade-tolerant *Quercus* (Kato et al., 2009).




Figure 7. (a) Mean (\pm SE bars) growing-season daily water use (DWU) versus forest age in the treated WS6 and reference (WS14, WS18) watersheds; (b) DWU versus forest age in treated WS6 by xylem functional group (diffuse-porous, ring-porous, semi-ring porous, evergreen shrub, and tracheid); (c) DWU versus forest age in reference WS14 by xylem functional group; and (d) DWU versus forest age in reference WS18 by xylem functional group.

Many studies have investigated forest growth following harvesting (e.g., Palik et al., 2012; Boring et al., 2014; Loftis et al., 2014; Stanturf et al., 2014; Boggs et al., 2016), and the hardwood species composition that succeeds following harvest depends largely on the severity of disturbance, i.e., partial harvest, retention harvest, or clear-cutting, as well as the geographical region (Halpin and Lorimer, 2016). In northern Appalachian forests, *Prunus pensylvanica* and *Betula papyrifera* are common pioneer species that assume early dominance following clear-cutting (Hornbeck et al., 2014). In central Appalachian forests, *Prunus serotina*, *Acer rubrum*, *Betula lenta*, and *Fagus grandifolia* dominate following ex-

tensive harvests (Kochenderfer, 2006; D’Amato et al., 2015). *Robinia pseudoacacia* and *L. tulipifera*, two species that recruit and grow rapidly after clear-cutting, are much more abundant in the southern Appalachians (Elliott and Vose, 2011; Boring et al., 2014) than in the central Appalachians (Kochenderfer, 2006), and are absent in the northern Appalachians (Campbell et al., 2007; Hornbeck et al., 2014).

4.2 Species effects on water yield (Q) and evapotranspiration (ET)

We found that annual Q declined and ET increased through old-field succession relative to the time prior to the grass conversion. After 1980, 13 years following herbicide application, Q was consistently lower than expected for the next 35 years. Q was reduced by 6.5 % averaged over this time period; however, in 16 of those years, Q was greatly reduced (> 50 mm, 9.2 %). In 2003 and 2015, Q was reduced by 142 mm (16 %) and 113 mm (17 %), respectively. This supports our hypothesis that changes in ET and Q have occurred as a result of a shift in species composition. We also found that species effects were seasonal and influenced certain parts of the flow regime.

The range of changes in Q after treatment suggests that species composition affects storage and use of water under a wide range of precipitation conditions that play out over monthly and annual scales. For example, in 2003, when the decrease in Q was greatest (-142 mm; 16 %), P was 6 % greater than the long-term (1939–2015) average but this followed 4 years of below-average P . Average P for 1999–2002 was 23 % below the long-term average. In this case, the vegetation in the old-field succession watershed may have used more of the available water in 2003, following the dry period, than the vegetation in reference watersheds. As a result, less of the available water served to refill soil storage in the treated watershed compared to the reference watershed, resulting in a larger predicted decrease in Q in 2003. In 1981 when Q was higher than expected ($+30$ mm; 5.5 %), P was 29 % lower than the long-term average but this followed the second highest annual P in 1980 ($+27$ % greater than the 1939–2015 average). Much of the excess rainfall occurred at the end of the 1980 water year and the beginning of water year 1981. Precipitation during March and April of water year 1980, and May of water year 1981 was 123, 35, and 39 % greater than the long-term (1939–2015) average for those months, respectively. Given that the Q for the treated watershed was higher than expected in wetter months (those above median Q) of the dormant season, these wet months resulted in a higher-than-expected annual Q .

Our monthly analysis showed that changes in ET and Q varied seasonally. First, changes in monthly distribution of Q suggest that old-field succession and the consequent species changes have lowered streamflow during the growing season during wetter months. We observed that Q was lower than expected in September during both drier (below me-

dian Q) and wetter periods suggesting that changes in soil storage at the end of the growing season highly influences base flow. Others have found that forest clear-cutting had a longer-lasting influence on streamflow distribution, even when annual Q returned to baseline conditions within a few years (Burt et al., 2015; Kelly et al., 2016).

Second, changes in monthly distribution of Q suggest that there is a potential for increased frequency and severity of high flows in dormant season months under wet conditions. This could be particularly concerning during severe tropical storms. However, for the Appalachian region most tropical storms occur later in the year (September–December) (Holland and Webster, 2007; Smith et al., 2011). Interestingly, we found lower-than-expected Q during wetter periods for September–December months. If timing of large storms remains unchanged, then shifting species composition from those that have conservative water use (i.e., ring-porous xylem) to those that are less conservative (i.e., diffuse-porous xylem) could mitigate the effects of high flows during large storms.

The observed changes in monthly Q during the dormant season indicate a likelihood of soil saturation during the wettest periods. Higher-than-expected Q in the dormant season is likely a result of lower ET and higher soil moisture at that time of year (Berghuijs et al., 2014; Burt et al., 2015), rather than reduced infiltration capacity. For example, in an earlier study, Burt and Swank (1992) reported that the dead grass was not removed following herbicide application on the treated watershed and so the infiltration capacity remained high throughout 1967 and 1968. More likely the higher-than-expected Q in the dormant season is due to the lack of evergreen species in the treated watershed. Where evergreen species are a component of forested watersheds, they can affect ET and Q in the dormant season (Brantley et al., 2013, 2015); they transpire during dormant season months as long as environmental conditions are suitable (Ford and Vose, 2007; Ford et al., 2011a; Brantley, unpublished data) and they intercept precipitation during the dormant season because they retain their foliage. Even though evergreens (shrubs + tracheids) were a relatively small component (13.8 % of total aboveground biomass) of the old-field succession watershed before treatment, after treatment there were no evergreen shrubs due to the severity of the treatment. Yet, they remain a component (6.0 and 15.9 % for WS14 and WS18, respectively) of the reference watersheds. Thus, evergreen species reduce soil moisture storage and have the potential to mitigate spring flooding because of their contribution to ET and their location within riparian zones (Brantley et al., 2015).

Our results demonstrate that species changes largely explain the decreasing trend in Q following old-field abandonment based on modeled growing-season DWU over time, and enable us to assess the effects of forest structure and species composition on Q . For example, the estimates of DWU (Fig. 6) are consistent with the differences in tem-

poral patterns of Q between the old-field succession WS6 and reference watersheds (Fig. 3). The mean DWU in WS6 was greater in 1995 than DWU in the reference watersheds in 1969 or 1992, suggesting that Q in WS6 became less than expected between these years due to altered DWU. Similarly, mean DWU in the 45-year-old old-field succession WS6 was greater still in 2012 than the > 75-year-old reference watersheds, WS14 or WS18, in 2010. Indeed, Q was consistently less than expected during this period, and was significantly less in 32 of the 35 years (including 1995, 2010, and 2012).

Few studies have examined the consequence of shifts in hardwood species composition on the hydrologic cycle (Swank et al., 2014; Caldwell et al., 2016). Changes in forest composition, structure and age as well as climate will interact to induce long-term changes in Q from forested mountain watersheds (Von Allmen et al., 2015; Caldwell et al., 2016). We found stronger and longer-lasting decreases in annual and monthly Q through old-field succession, than found by clear-cutting alone followed by forest succession (Reinhart, 1964; Hornbeck, 1973; Swank et al., 2001, 2014; Troendle et al., 2001; Adams and Kochenderfer, 2014; Hornbeck et al., 2014). For example, researchers at the Fernow Experimental Forest in West Virginia examined changes in annual Q following clear-cutting (Adams and Kochenderfer, 2014); there, the initial increase in Q returned to pretreatment levels within 3–4 years. In another treated watershed (WS7) in Coweeta that was allowed to regenerate naturally after the clear-cut, there was only 1 year when observed Q was significantly lower than predicted (Swank et al., 2014).

The observed changes in monthly and annual Q for the treated WS6 were largely a result of a rapid response of co-dominant species with less conservative transpiration rates (Wullschlegel et al., 2001; Ford et al., 2012; Boggs et al., 2015; Brantley et al., 2015). Under similar environmental conditions, both *L. tulipifera* and *R. pseudoacacia* have much higher daily water use than species with ring-porous and semi-ring-porous xylem, such as *Quercus* and *Carya* (Ford et al., 2011b; Vose et al., 2016a, b). Overall, we estimated that growing-season daily water use increased significantly following old-field abandonment, and it was much higher in the 45-year-old treated watershed than the > 75-year-old reference watersheds.

5 Conclusions

Our long-term results are relevant to land areas that are currently in pasture and those that have reverted back to forests. In many parts of the world, pasture land and cropland area have increased since the 1990s as natural landscapes have been converted to agricultural ecosystems (e.g., Scanlon et al., 2007; Rodriguez et al., 2010); in other areas agricultural land has been abandoned (see review Rey Benayas et al., 2007). In general, grass pastures transpire less water and have lower interception loss than forests, resulting in greater Q for

this LULC type (e.g., Wang et al., 2008; Holdo and Nippert, 2015). In the forest–grass–forest watershed, for 2 of the 5 years when the watershed was in grass cover, Q was equivalent to the pre-conversion hardwood forest, while for the other 3 years Q was greater under grass cover. Fertilizer application in 2 of the 5 years resulted in high grass productivity (Hewlett, 1961; Hewlett and Hibbert, 1966; Burt and Swank, 1992) such that LAI was maximized allowing for ET similar to that of the reference forested watershed. Q increased initially once herbicide was applied to the grass, quickly returned to expected levels, and then declined relative to expected levels as the abandoned old-field was allowed to regenerate to forest.

We found that within a deciduous forest, species identity matters in terms of how much precipitation leaves the watershed as ET versus Q . Through old-field succession, the treated watershed was dominated by water-demanding species with higher DWU than the pretreatment forest. We demonstrate that a shift in tree species composition from dominance by species with ring-porous xylem to dominance by species with diffuse-porous xylem can increase DWU, and in turn, produce a long-term reduction in Q .

Even within unmanaged watersheds, hydrologic parameters are not stationary (*sensu* Milly et al., 2008; Burt et al., 2015) and subtle changes in species composition can influence Q , particularly in dry years (Caldwell et al., 2016). Species-specific ecohydrological models (e.g., Novick et al., 2016) are increasingly vital in predicting long-term changes in ET and Q (Sun et al., 2016; Vose et al., 2016a, b). If drought frequency and severity increase as expected (Allen et al., 2010; Ayres et al., 2014; Peters et al., 2015; Swain and Hayhoe, 2015), then understanding the interaction of land use, species, and climate change on water resources will become even more important in the future (Grant et al., 2013; Clark et al., 2016; Kelly et al., 2016; Vose et al., 2016b). As previously outlined as a critical research need (Vose et al., 2016b), our results provide an example of scaling DWU from tree level, plots, and small watersheds in order to understand the species-specific influences on water balance and stream-flow dynamics in diverse eastern US deciduous forests.

6 Data availability

All data in this manuscript are archived at USDA Forest Service, Southern Research Station, Coweeta Hydrologic Laboratory, Otto, NC, 28763.

The Supplement related to this article is available online at doi:10.5194/hess-21-981-2017-supplement.

Author contributions. Wayne T. Swank and Katherine J. Elliott conceived the study and designed the experiment. Katherine J. Elliott and Peter V. Caldwell wrote the paper. Katherine J. Elliott, Peter V. Caldwell, Chelcy F. Miniati, and Steven T. Brantley performed

the analyses. Wayne T. Swank, James M. Vose, Chelcy F. Miniati, and Katherine J. Elliott performed the research. Katherine J. Elliott, Peter V. Caldwell, Steven T. Brantley, Chelcy F. Miniati, James M. Vose, and Wayne T. Swank contributed to discussions and editing.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was supported by the USDA Forest Service, Southern Research Station, and by NSF grants DEB0218001 and DEB0823293 to the Coweeta LTER program at the University of Georgia. Any opinions, findings, conclusions, or recommendations expressed in the material are those of the authors and do not necessarily reflect the views of the National Science Foundation or the University of Georgia. We acknowledge the support of the long-term climate and hydrologic data network at Coweeta Hydrologic Laboratory as well as many individuals, past and present, especially Patsy Clinton, Charles Marshall, and Stephanie Laseter for field and climate data collection and processing.

Edited by: K. Bishop

Reviewed by: two anonymous referees

References

- Adams, M. B. and Kochenderfer, J. N.: Recovery of Central Appalachian forested watersheds: Comparison of Fernow and Coweeta, in: Long-Term Response of a Forest Watershed Ecosystem: Clearcutting in the Southern Appalachians (The Long-term Ecological Research Network Series), edited by: Swank, W. T. and Webster, J. R., Oxford University Press, Oxford, UK, 194–212, 2014.
- Alila, Y., Kuraś, P. K., Schnorbus, M., and Hudson, R.: Forests and floods: a new paradigm sheds light on age-old controversies, *Water Resour. Res.*, 45, W08416, doi:10.1029/2008WR007207, 2009.
- Allen, C. D., Macalady, A. K., Chenchouni, H., Bachelet, D., McDowell, N., Vennetier, M., Kitzberger, T., Rigling, A., Breshears, D. D., Hogg, E. H., Gonzalez, P., Fensham, R., Zhang, Z., Castro, J., Demidova, N., Lim, J.-H., Allard, G., Running, S. W., Semerci, A., and Cobb, N.: A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests, *Forest Ecol. Manage.*, 259, 660–684, doi:10.1016/j.foreco.2009.09.001, 2010.
- Alvarez, M.: The State of America's forests, DIANE Publishing, Darby, Pennsylvania, 2007.
- Amatya, D. M. and Harrison, C. A.: Grass and forest potential evapotranspiration comparison using five methods in the Atlantic coastal plain, *J. Hydrol. Eng.*, 21, 05016007, doi:10.1061/(ASCE)HE.1943-5584.0001341, 2016.
- Andréassian, J. H.: Waters and forests: from historical controversy to scientific debate, *J. Hydrol.*, 291, 1–27, doi:10.1016/j.jhydrol.2003.12.015, 2004.

- Aranda, I., Forner, A., Cuesta, B., and Valladares, F.: Species-specific water use by forest tree species: From the tree to the stand, *Agr. Water Manage.*, 114, 67–77, doi:10.1016/j.agwat.2012.06.024, 2012.
- Ayres, M. P., Hicke, J. A., Kerns, B. K., McKenzie, D., Littell, J. S., Band, L. E., Luce, C. H., Weed, A. S., and Raymond, C. L.: Disturbance regimes and stressors, in: *Climate Change and United States Forests*, edited by: Peterson, D. L., Vose, J. M., and Patel-Weynand, T., Springer, New York, 55–92, 2014.
- Bellemare, J., Motzkin, G., and Foster, D. R.: Legacies of the agricultural past in the forested present: an assessment of historical land-use effects on rich mesic forests, *J. Biogeogr.*, 29, 1401–1420, doi:10.1046/j.1365-2699.2002.00762.x, 2002.
- Berghuijs, W. R., Sivapalan, M., Woods, R. A., and Savenije, H. H. G.: Patterns of similarity of seasonal water balances: A window into streamflow variability over a range of time scales, *Water Resour. Res.*, 50, 5638–5661, doi:10.1002/2014WR015692, 2014.
- Boggs, J., Sun, G., Domec, J.-C., McNulty, S., and Treasure, E.: Clearcutting upland forest alters transpiration of residual trees in the riparian buffer zone, *Hydrol. Process.*, 29, 4979–4992, doi:10.1002/hyp.10474, 2015.
- Boggs, J., Sun, G., and McNulty, S.: Effects of timber harvest on water quantity and quality in small watersheds in the Piedmont of North Carolina, *J. Forest.*, 114, 27–40, doi:10.5849/jof.14-102, 2016.
- Boring, L. R., Elliott, K. J., and Swank, W. T.: Successional forest dynamics 30 years following clearcutting, in: *Long-Term Response of a Forest Watershed Ecosystem: Clearcutting in the Southern Appalachians (The Long-term Ecological Research Network Series)*, edited by: Swank, W. T. and Webster, J. R., Oxford University Press, Oxford, UK, 11–35, 2014.
- Bosch, J. M. and Hewlett, J. D.: A review of catchment experiments to determine the effect of vegetation changes on water yield and evapotranspiration, *J. Hydrol.*, 33, 3–23, doi:10.1016/0022-1694(82)90117-2, 1982.
- Brantley, S., Ford, C. R., and Vose, J. M.: Future species composition will affect forest water use after loss of hemlock from southern Appalachian forests, *Ecol. Appl.*, 23, 777–790, doi:10.1890/12-0616.1, 2013.
- Brantley, S. T., Miniati, C. F., Elliott, K. J., Laseter, S. H., and Vose, J. M.: Changes to southern Appalachian water yield and stormflow after loss of a foundation species, *Ecohydrology*, 8, 518–528, doi:10.1002/eco.1521, 2015.
- Brauman, K. A., Freyberg, D. L., and Daily, G. C.: Potential evapotranspiration from forest and pasture in the tropics: A case study in Kona, Hawaii, *J. Hydrol.*, 440–441, 52–61, doi:10.1016/j.jhydrol.2012.03.014, 2012.
- Brown, A. E., Zhang, L., McMahon, T. A., Western, A. W., and Vertessy, R. A.: A review of paired catchment studies for determining changes in water yield resulting from alterations in vegetation, *J. Hydrol.*, 310, 28–61, doi:10.1016/j.jhydrol.2004.12.010, 2005.
- Burt, T. and Swank W. T.: Forests or floods?, *Geogr. Rev.*, 15, 37–41, 2002.
- Burt, T. P. and Swank, W. T.: Flow frequency responses to hardwood-to-grass conversion and subsequent succession, *Hydrol. Process.*, 6, 179–188, doi:10.1002/hyp.3360060206, 1992.
- Burt, T. P., Howden, N. J. K., McDonnell, J. J., Jones, J. A., and Hancock, G. R.: Seeing the climate through the trees: observing climate and forestry impacts on streamflow using a 60-year record, *Hydrol. Process.*, 29, 473–480, doi:10.1002/hyp.10406, 2015.
- Burton, T. A.: Effects of basin-scale timber harvest on water yield and peak streamflow, *J. Am. Water Resour. Assoc.*, 33, 1187–1194, doi:10.1111/j.1752-1688.1997.tb03545.x, 1997.
- Caldwell, P. V., Miniati, C. F., Elliott, K. J., Swank, W. T., Brantley, S. T., and Laseter, S. H.: Declining water yield from forested mountain watersheds in response to climate change and forest mesophication, *Global Change Biol.*, 22, 2997–3012, doi:10.1111/gcb.13309, 2016.
- Campbell, J. L., Driscoll, C. T., Eagar, C., Likens, G. E., Siccama, T. G., Johnson, C. E., Fahey, T. J., Hamburg, S. P., Holmes, R. T., Bailey, A. S., and Buso, D. C.: Long-term trends from ecosystem research at the Hubbard Brook Experimental Forest, GTR-NRS-17, USDA Forest Service, Northern Research Station, Newtown Square, Pennsylvania, 2007.
- Chang, M.: *Forest Hydrology: An Introduction to Forests and Water*, CRC Press, Boca Raton, Florida, 2003.
- Clark, J. S., Iverson, L., Woodall, C. W., Allen, C. D., Bell, D. M., Bragg, D. C., D’Amato, A. W., Davis, F. W., Hersh, M. H., Ibanez, I., Jackson, S. T., Matthews, S., Pederson, N., Peters, M., Schwartz, M. W., Waring, K. M., and Zimmermann, N. E.: The impacts of increasing drought on forest dynamics, structure, and biodiversity in the United States, *Global Change Biol.*, 22, 2329–2352, doi:10.1111/gcb.13160, 2016.
- Cramer, V. A., Hobbs, R. J., and Standish, R. J.: What’s new about old fields: Land abandonment and ecosystem assembly, *Trends Ecol. Evol.*, 23, 104–112, doi:10.1016/j.tree.2007.10.005, 2008.
- Cruise, J. F., Laymon, C. A., and Al-Hamdan, O. A.: Impact of 20 years of land-cover change on the hydrology of streams in the southeastern United States, *J. Am. Water Resour. Assoc.*, 46, 1159–1170, doi:10.1111/j.1752-1688.2010.00483.x, 2010.
- Cunnane, C.: Unbiased plotting positions – a review, *J. Hydrol.*, 37, 205–222, 1978.
- D’Amato, A. W., Catanzaro, P. F., and Fletcher L. S.: Early regeneration and structural response to patch selection and structural retention in second-growth northern hardwoods, *Forest Sci.*, 61, 183–189, doi:10.5849/forsci.13-180, 2015.
- Douglass, J. E. and Hoover, M. D.: History of Coweeta, in: *Forest Hydrology and Ecology at Coweeta, Ecological Studies*, edited by: Swank, W. T. and Crossley Jr., D. A., Springer-Verlag, New York, 17–31, 1988.
- Douglass, J. E. and Swank, W. T.: Streamflow modification through management of eastern forests, Res. Paper SE-94, USDA Forest Service, Southeastern Forest Experiment Station, Asheville, North Carolina, 1972.
- Douglass, J. E., Cochrane, D. R., Bailey, G. W., Teasley, J. I., and Hill, D. W.: Low herbicide concentration found in streamflow after a grass cover is killed, Res. Note SE-108, USDA Forest Service, Southeastern Forest Experiment Station, Asheville, North Carolina, 1969.
- Dunford, E. G. and Fletcher, P. W.: Effect of removal of streambank vegetation upon water yield, *T. Am. Geophys. Un.*, 28, 105–110, 1947.
- Edburg, S. L., Hicke, J. A., Brooks, P. D., Pendall, E. G., Ewers, B. E., Norton, U., Gochis, D., Gutmann, E. D., and Meddens, A. J. H.: Cascading impacts of bark beetle-caused tree mortality

- on coupled biogeophysical and biogeochemical processes, *Front. Ecol. Environ.*, 10, 416–424, doi:10.1890/110173, 2012.
- Elliott, K. J. and Swank, W. T.: Changes in tree species diversity after successive clearcuts in the southern Appalachians, *Vegetatio*, 115, 11–18, doi:10.1007/BF00119382, 1994.
- Elliott, K. J. and Swank, W. T.: Long-term changes in forest composition and diversity following early logging (1919–1923) and the decline of American chestnut (*Castanea dentata* (Marshall) Borkh.), *Plant Ecol.*, 197, 155–172, doi:10.1007/s11258-007-9352-3, 2008.
- Elliott, K. J. and Vose, J. M.: The contribution of the Coweeta Hydrologic Laboratory to developing an understanding of long-term (1934–2008) changes in managed and unmanaged forests, *Forest Ecol. Manage.*, 261, 900–910, doi:10.1016/j.foreco.2010.03.010, 2011.
- Elliott, K. J., Boring, L. R., Swank, W. T., and Haines, B. R.: Successional changes in diversity and composition in a clearcut watershed in Coweeta Basin, North Carolina, *Forest Ecol. Manage.*, 92, 67–85, doi:10.1016/S0378-1127(96)03947-3, 1997.
- Elliott, K. J., Boring, L. R., and Swank, W. T.: Changes in vegetation structure and diversity following grass-to-forest succession, *Am. Midl. Nat.*, 140, 219–232, doi:10.1674/0003-0031(1998)140[0219:CIVSAD]2.0.CO;2, 1998.
- Elliott, K. J., Boring, L. R., and Swank, W. T.: Aboveground biomass and nutrient accumulation 20 years after clear-cutting a southern Appalachian watershed, *Can. J. Forest Res.*, 32, 667–683, doi:10.1139/x02-009, 2002.
- Farley, K. A., Jobbagy, E. G., and Jackson, R. B.: Effects of afforestation on water yield: A global synthesis with implications for policy, *Global Change Biol.*, 11, 1565–1576, doi:10.1111/j.1365-2486.2005.01011.x, 2005.
- Ford, C. R. and Vose, J. M.: *Tsuga canadensis* (L.) Carr. mortality will impact hydrologic processes in southern Appalachian forest ecosystems, *Ecol. Appl.*, 17, 1156–1167, doi:10.1890/06-0027, 2007.
- Ford, C. R., Hubbard, R. M., and Vose, J. M.: Quantifying structural and physiological controls on canopy transpiration of planted pine and hardwood stands in the southern Appalachians, *Ecohydrology*, 4, 183–195, doi:10.1002/eco.136, 2011a.
- Ford, C. R., Laseter, S. H., Swank, W. T., and Vose, J. M.: Can forest management be used to sustain water-based ecosystem services in the face of climate change?, *Ecol. Appl.*, 21, 2049–2067, doi:10.1890/10-2246.1, 2011b.
- Ford, C. R., Elliott, K. J., Clinton, B. D., Kloeppe, B. D., and Vose, J. M.: Forest dynamics following hemlock mortality in the southern Appalachians, *Oikos*, 121, 523–536, doi:10.1111/j.1600-0706.2011.19622.x, 2012.
- Foster, D., Swanson, F., Aber, J., Burke, I., Brokaw, N., Tilman, D., and Knapp, A.: The importance of land-use legacies to ecology and conservation, *BioScience*, 53, 77–88, doi:10.1641/0006-3568(2003)053[0077:TLOLUL]2.0.CO;2, 2003.
- García-Ruiz, J. and Lana-Renault, N.: Hydrological and erosive consequences of farmland abandonment in Europe, with special reference to the Mediterranean region – A review, *Agr. Ecosyst. Environ.*, 140, 317–338, doi:10.1016/j.agee.2011.01.003, 2011.
- Grant, G. E., Tague, C. L., and Allen, C. D.: Watering the forest for the trees: an emerging priority for managing water in forest landscapes, *Front. Ecol. Environ.*, 11, 314–321, doi:10.1890/120209, 2013.
- Green, K. C. and Alila, Y.: A paradigm shift in understanding and quantifying the effects of forest harvesting on floods in snow environments, *Water Resour. Res.*, 48, W10503, doi:10.1029/2012WR012449, 2012.
- Halpin, C. R. and Lorimer, C. G.: Trajectories and resilience of stand structure in response to variable disturbance severities in northern hardwoods, *Forest Ecol. Manage.*, 365, 69–82, doi:10.1016/j.foreco.2016.01.016, 2016.
- Harr, R. D., Harper, W. C., Krygier, J. T., and Hsieh, F. S.: Changes in storm hydrographs after road building and clear-cutting in the Oregon Coast Range, *Water Resour. Res.*, 11, 436–444, doi:10.1029/WR011i003p00436, 1975.
- Hassler, S. K., Zimmermann, B., van Breugel, M., Hall, J. S., and Elsenbeer, H.: Recovery of saturated hydraulic conductivity under secondary succession on former pasture in the humid tropics, *Forest Ecol. Manage.*, 261, 1634–1642, doi:10.1016/j.foreco.2010.06.031, 2011.
- Hewlett, J. D.: Response of fescue to natural moisture gradient on an artificial slope, Res. Note SE-152, USDA Forest Service, Southeastern Forest Experiment Station, Asheville, North Carolina, 1961.
- Hewlett, J. D. and Hibbert, A. R.: Increases in water yield after several types of forest cutting, *Q. Bull. Int. Assoc. Sci. Hydrol.*, 6, 5–17, 1961.
- Hewlett, J. D. and Hibbert, A. R.: Factors affecting the response of small watersheds to precipitation in humid areas, in: *Forest Hydrology*, edited by: Soppe, W. E. and Lull, H. W., Pergamon Press, Oxford, UK, 275–290, 1966.
- Hibbert, A. R.: Water yield changes after converting a forested catchment to grass, *Water Resour. Res.*, 5, 634–640, 1969.
- Holdo, R. M. and Nippert, J. B.: Transpiration dynamics support resource partitioning in African savanna trees and grasses, *Ecology*, 96, 1466–1472, doi:10.1890/14-1986.1, 2015.
- Holland, G. J. and Webster, P. J.: Heightened tropical cyclone activity in the North Atlantic: natural variability of climate trend?, *Philos. T. Roy. Soc. A*, 365, 2695–2716, doi:10.1098/rsta.2007.2083, 2007.
- Hornbeck, J. W.: Storm flow from hardwood-forested and cleared watersheds in New Hampshire, *Water Resour. Res.*, 9, 346–354, 1973.
- Hornbeck, J. W., Bailey, A. S., Eagar, C., and Campbell, J. L.: Comparisons with results from the Hubbard Brook Experimental Forest in the northern Appalachians, in: *Long-Term Response of a Forest Watershed Ecosystem: Clearcutting in the Southern Appalachians* (The Long-term Ecological Research Network Series), edited by: Swank, W. T. and Webster, J. R., Oxford University Press, Oxford, UK, 213–228, 2014.
- Houlbrooke, D. J. and Laurenson, S.: Effect of sheep and cattle treading damage on soil microporosity and soil water holding capacity, *Agr. Water Manage.*, 121, 81–84, doi:10.1016/j.agwat.2013.01.010, 2013.
- Ice, G. G. and Stednick, J. D.: *A Century of Forest and Wildland Watershed Lessons*, Society of American Foresters, Bethesda, Maryland, 2004.
- Kato, A., Moskal, L. M., Schiess, P., Swanson, M. E., Calhoun, D., and Stuetzle, W.: Capturing tree crown formation through implicit surface reconstruction using airborne lidar data, *Remote Sens. Environ.*, 113, 1148–1162, doi:10.1016/j.rse.2009.02.010, 2009.

- Kelly, C. N., McGuire, K. J., Miniati, C. F., and Vose, J. M.: Streamflow response to increasing precipitation extremes altered by forest management, *Geophys. Res. Lett.*, 43, 3727–3736, doi:10.1002/2016GL068058, 2016.
- Kirk, R. W., Bolstad, P. V., and Manson, S. M.: Spatio-temporal trend analysis of long-term development patterns (1900–2030) in a Southern Appalachian county, *Landsc. Urban Plan.*, 104, 47–58, doi:10.1016/j.landurbplan.2011.09.008, 2012.
- Kirkman, L. K., Brown, C. L., and Leopold, D. J.: *Native Trees of the Southeast*, Timber Press, Portland, Oregon, 2007.
- Klein, T.: The variability of stomatal sensitivity to leaf water potential across tree species indicates a continuum between isohydric and anisohydric behaviours, *Funct. Ecol.*, 28, 1313–1320, doi:10.1111/1365-2435.12289, 2014.
- Kochenderfer, J. N.: Fernow and the Appalachian hardwood region, in: *The Fernow Watershed Acidification Study*, edited by: Adams, M. B., DeWalle, D. R., and Hom, H. L., Springer, Dordrecht, the Netherlands, 17–39, 2006.
- Kulmatiski, A. and Beard, K. H.: Woody plant encroachment facilitated by increased precipitation intensity, *Nat. Clim. Change*, 3, 833–837, doi:10.1038/nclimate1904, 2013.
- Laseter, S. H., Ford, C. R., Vose, J. M., and Swift, L. W. Jr.: Long-term temperature and precipitation trends at the Coweeta Hydrologic Laboratory, Otto, North Carolina, USA, *Hydrol. Res.*, 43, 890–901, doi:10.2166/nh.2012.067, 2012.
- Liu, W., Wei, X., Liu, S., Liu, Y., Fan, H., Zhang, M., Yin, J., and Zhan, M.: How do climate and forest changes affect long-term streamflow dynamics? A case study in the upper reach of Poyang River basin, *Ecohydrology*, 8, 46–57, doi:10.1002/eco.1486, 2015.
- Loftis, D. L., Schweitzer, C. J., and Keyser, T. L.: Structure and species composition of upland hardwood communities after regeneration treatments across environmental gradients, in: *Sustaining Young Forest Communities: Ecology and management of early successional habitats in the central hardwood region, USA*, edited by: Greenberg, C., Collins, B., and Thompson III, F., Springer, New York, 59–71, 2014.
- Martin, J. G., Klooppel, B. D., Schaefer, T. L., Kimbler, D. L., and McNulty, S. G.: Aboveground biomass and nitrogen allocation of ten deciduous southern Appalachian tree species, *Can. J. Forest Res.*, 28, 1648–1659, doi:10.1139/x98-149, 1998.
- Martinuzzi, S., Bavier-Pizarro, G. I., Lugo, A. E., and Radeloff, V. C.: Future land-use changes and the potential for novelty in ecosystems of the United States, *Ecosystems*, 18, 1332–1342, doi:10.1007/s10021-015-9901-x, 2015.
- McGinty, D. T.: *The ecological roles of Kalmia latifolia L. and Rhododendron maximum L. in the hardwood forest at Coweeta*, MS thesis, University of Georgia, Athens, Georgia, 1972.
- Milly, P. C. D., Betancourt, J., Falkenmark, M., Hirsch, R. M., Kundzewicz, Z. W., Lettenmaier, D. P., and Stouffer, R. J.: Climate change – Stationarity is dead: Whither water management?, *Science*, 319, 573–574, doi:10.1126/science.1151915, 2008.
- Morris, L. A. and Jackson, C. R.: Interactions of soils and land uses with water quantity and quality, in: *Soil Ecosystems Services*, edited by: Stromberger, M., Comerford, N., and Lindbo, D., Soil Sci. Soc. Am., Madison, WI, doi:10.2136/2015.soilecosystemsservices.2014.0077, 2016.
- Neary, D. G., Ice, G. G., and Jackson, C. R.: Linkages between forest soils and water quality and quantity, *Forest Ecol. Manage.*, 258, 2269–2281, doi:10.1016/j.foreco.2009.05.027, 2009.
- Novick, K. A., Miniati, C. F., and Vose, J. M.: Drought limitations to leaf-level gas exchange: results from a model linking stomatal optimization and cohesion-tension theory, *Plant Cell Environ.*, 39, 583–596, doi:10.1111/pce.12657, 2016.
- Otto, J. S.: The decline of forest farming in southern Appalachia, *J. Forest.*, 27, 18–27, 1983.
- Palik, B., Martin, M., Zenner, E., Blinn, C., and Kolka, R.: Overstory and regeneration dynamics in riparian management zones of northern Minnesota forested watersheds, *Forest Ecol. Manage.*, 271, 1–9, doi:10.1016/j.foreco.2012.01.045, 2012.
- Peters, M. P., Iverson, L. R., and Matthews, S. N.: Long-term droughtiness and drought tolerance of eastern US forests over five decades, *Forest Ecol. Manage.*, 345, 56–64, doi:10.1016/j.foreco.2015.02.022, 2015.
- Price, K., Jackson, C. R., Parker, A. J., Reitan, T., Dowd, J., and Cyterski, M.: Effects of watershed land use and geomorphology on stream low flows during severe drought conditions in the southern Blue Ridge Mountains, Georgia and North Carolina, United States, *Water Resour. Res.*, 47, W02516, doi:10.1029/2010WR009340, 2011.
- Ramankutty, N., Heller, E., and Rhemtulla, J.: Prevailing myths about agricultural abandonment and forest regrowth in the United States, *Ann. Assoc. Am. Geogr.*, 100, 502–512, doi:10.1080/00045601003788876, 2010.
- Reinhart, K. G.: Effect of a commercial clearcutting in West Virginia on overland flow and storm runoff, *J. Forest.*, 62, 167–171, 1964.
- Reinhart, K. G. and Pierce, R. S.: *Stream-gaging stations for research on small watersheds*, USDA Forest Service, Washington, D.C., 1964.
- Rey Benayas, J. M., Martins, A., Nicolau, J. M., and Schulz, J. J.: Abandonment of agricultural land: an overview of drivers and consequences, *Perspec. Agric. Vet. Sci. Nutr. Nat. Resour. No. 057*, CABI Publishing, Oxfordshire, UK, doi:10.1079/PAVSNNR20072057, 2007.
- Rodriguez, D. A., Tomasella, J., and Linhares, C.: Is the forest conversion to pasture affecting the hydrological response of Amazonian catchments: Signals in the Ji-Paraná Basin, *Hydrol. Process.*, 24, 1254–1269, doi:10.1002/hyp.7586, 2010.
- Santee, W. R. and Monk, C. D.: Stem diameter and dry weight relationships in *Tsuga canadensis* (L.) Carr., *Bull. Torrey Bot. Club*, 108, 320–323, 1981.
- Scanlon, B. R., Jolly, I., Sophocleous, M., and Zhang, L.: Global impacts of conversions from natural to agricultural ecosystems on water resources: Quantity versus quality, *Water Resour. Res.*, 43, W03437, doi:10.1029/2006WR005486, 2007.
- Shure, D. J., Phillips, D. L., and Bostick, P. E.: Gap size and succession in cutover southern Appalachian forests: an 18 year study of vegetation dynamics, *Plant Ecol.*, 185, 299–318, doi:10.1007/s11258-006-9105-8, 2006.
- Smith, J. A., Villarini, G., and Baeck, M. L.: Mixture distributions and the hydroclimatology of extreme rainfall and flooding in the Eastern United States, *J. Hydrometeorol.*, 12, 294–309, doi:10.1175/2010JHM1242.1, 2011.
- Stanturf, J. A., Palik, B. J., and Dumroese, R. K.: Contemporary forest restoration: A review emphasizing function, *Forest*

- Ecol. Manage., 331, 292–323, doi:10.1016/j.foreco.2014.07.029, 2014.
- Stedinger, J. R., Vogel, R. M., and Foufoula-Georgiou, E.: Frequency analysis of extreme events, in: Handbook of Hydrology, edited by: Maidment, D. R., McGraw-Hill, New York, 18.10–18.66, 1993.
- Stednick, J. D.: Monitoring the effects of timber harvest on annual water yield, *J. Hydrol.*, 176, 79–95, doi:10.1016/0022-1694(95)02780-7, 1996.
- Sun, S., Sun, G., Cohen, E., McNulty, S. G., Caldwell, P. V., Duan, K., and Zhang, Y.: Projecting water yield and ecosystem productivity across the United States by linking an ecohydrological model to WRF dynamically downscaled climate data, *Hydrol. Earth Syst. Sci.*, 20, 935–952, doi:10.5194/hess-20-935-2016, 2016.
- Swain, S. and Hayhoe, K.: CMIP5 projected changes in spring and summer drought and wet conditions over North America, *Clim. Dynam.*, 44, 2737–2750, doi:10.1007/s00382-014-2255-9, 2015.
- Swank, W. T. and Crossley, D. A. Jr.: Introduction and site description, in: Forest Hydrology and Ecology at Coweeta, Ecological Studies, edited by: Swank, W. T. and Crossley Jr., D. A., Springer-Verlag, New York, 3–16, 1988.
- Swank, W. T., Vose, J. M., and Elliott, K. J.: Long-term hydrologic and water quality responses following commercial clearcutting of mixed hardwoods on a southern Appalachian catchment, *Forest Ecol. Manage.*, 143, 163–178, doi:10.1016/S0378-1127(00)00515-6, 2001.
- Swank, W. T., Knoepp, J. D., Vose, J. M., Laseter, S. N., and Webster, J. R.: Response and recovery of water yield and timing, stream sediment, abiotic parameters, and stream chemistry following logging, in: Long-Term Response of a Forest Watershed Ecosystem: Clearcutting in the Southern Appalachians (The Long-term Ecological Research Network Series), edited by: Swank, W. T. and Webster, J. R., Oxford University Press, Oxford, UK, 36–56, 2014.
- Swift Jr., L. W., Cunningham, G. B., and Douglass, J. E.: Climate and hydrology, in: Forest Hydrology and Ecology at Coweeta, Ecological Studies, edited by: Swank, W. T. and Crossley Jr., D. A., Springer-Verlag, New York, 35–55, 1988.
- Taneda, H. and Sperry, J. S.: A case-study of water transport in co-occurring ring- versus diffuse-porous trees: contrasts in water status, conducting capacity, cavitation and vessel refilling, *Tree Physiol.*, 28, 1641–1652, doi:10.1093/treephys/28.11.1641, 2008.
- Thiemann, J. A., Webster, C. R., Jenkins, M. A., Hurley, P. M., Rock, J. H., and White, P. S.: Herbaceous-layer impoverishment in a post-agricultural Southern Appalachian landscape, *Am. Midl. Nat.*, 162, 148–168, doi:10.1674/0003-0031-162.1.148, 2009.
- Thomas, D. J.: Soil survey of Macon County, North Carolina, USDA Natural Resource Conservation Service, US Government Printing Office, Washington, D.C., 1996.
- Thompson, J. R., Carpenter, D. N., Cogbill, C. V., and Foster, D. R.: Four centuries of change in northeastern United States forests, *PLoS One*, 8, e72540, doi:10.1371/journal.pone.0072540, 2013.
- Trimble, S. W., Weirich, F. H., and Hoag, B. L.: Reforestation and the reduction of water yield on the southern piedmont since circa 1940, *Water Resour. Res.*, 23, 425–437, doi:10.1029/WR023i003p00425, 1987.
- Troendle, C. A., Wilcox, M. S., Bevenger, G. S., and Porth, L. S.: The Coon Creek water yield augmentation project: Implementation of timber harvesting technology to increase streamflow, *Forest Ecol. Manage.*, 143, 179–187, doi:10.1016/S0378-1127(00)00516-8, 2001.
- Von Allmen, E. I., Sperry, J. S., and Bush, S. E.: Contrasting whole-tree water use, hydraulics, and growth in a co-dominant diffuse-porous vs. ring-porous species pair, *Trees*, 29, 717–728, doi:10.1007/s00468-014-1149-0, 2015.
- Vose, J. M., Sun, G., Ford, C. R., Bredemeier, M., Ostsuki, K., Wei, A., Zhang, Z., and Zhang, L.: Forest ecohydrological research in the 21st century: what are the critical needs?, *Ecohydrology*, 4, 146–158, doi:10.1002/eco.193, 2011.
- Vose, J. M., Miniati, C. F., Luce, C. H., Asbjornsen, H., Caldwell, P. V., Campbell, J. L., Grant, G. E., Isaak, D. J., Loheide, S. P., and Sun, G.: Ecohydrological implications of drought, in: Effects of Drought on Forests and Rangelands in the United States: A comprehensive science synthesis, edited by: Vose, J. M., Clark, J. S., Luce, C. H., and Patel-Weyand, T., GTR-WO-93b, USDA Forest Service, Washington Office, Washington, D.C., 231–251, 2016a.
- Vose, J. M., Miniati, C. F., Luce, C. H., Asbjornsen, H., Caldwell, P. V., Campbell, J. L., Grant, G. E., Isaak, D. J., Loheide, S. P., and Sun, G.: Ecohydrological implications of drought for forests in the United States, *Forest Ecol. Manage.*, 380, 335–345, doi:10.1016/j.foreco.2016.03.025, 2016b.
- Wang, E., Cresswell, H., Paydar, Z., and Gallant, H.: Opportunities for manipulating catchment water balance by changing vegetation type on a topographic sequence: a simulation study, *Hydrol. Process.*, 22, 736–749, doi:10.1002/hyp.6655, 2008.
- Wear, D. N. and Bolstad, P.: Land-use changes in Southern Appalachian landscapes: spatial analysis and forecast evaluation, *Ecosystems*, 1, 575–594, doi:10.1007/s100219900052, 1998.
- Wei, X. and Zhang, M.: Quantifying streamflow change caused by forest disturbance at a large spatial scale: A single watershed study, *Water Resour. Res.*, 46, W12525, doi:10.1029/2010WR009250, 2010.
- Wilm, H. G.: Statistical control of hydrologic data from experimental watersheds, *T. Am. Geophys. Un.*, 2, 618–622, 1944.
- Wilm, H. G.: How long should experimental watersheds be calibrated?, *T. Am. Geophys. Un.*, 30, 272–278, 1949.
- Woods, F. W. and Shanks, R. E.: Natural replacement of chestnut by other species in the Great Smoky Mountains National Park, *Ecology*, 40, 349–361, 1959.
- Wullschlegel, S. D., Hanson, P. J., and Todd, D. E.: Transpiration from a multi-species deciduous forest as estimated by xylem sap flow techniques, *Forest Ecol. Manage.*, 143, 205–213, doi:10.1016/S0378-1127(00)00518-1, 2001.
- Yarnell, S. L.: The southern Appalachians: a history of the landscape, GTR-SRS-18, USDA Forest Service, Southern Research Station, Asheville, North Carolina, 1998.
- Zhang, L., Dawes, W. R., and Walker, G. R.: Response of mean annual evapotranspiration to vegetation changes at catchment scale, *Water Resour. Res.*, 37, 701–708, doi:10.1029/2000WR90032509.00, 2001.
- Zhang, M. and Wei, X.: The effects of cumulative forest disturbance on streamflow in a large watershed in the central interior of British Columbia, Canada, *Hydrol. Earth Syst. Sci.*, 16, 2021–2034, doi:10.5194/hess-16-2021-2012, 2012.

Zhou, G., Wei, X., Chen, X., Zhou, P., Liu, X., Xiao, Y., Sun, G., Scott, D. F., Zhou, S., Han, L., and Su, Y.: Global pattern for the effect of climate and land cover on water yield, *Nat. Commun.*, 6, 5918, doi:10.1038/ncomms6918, 2015.

Zimmermann, B., Papritz, A., and Elsenbeer, H.: Asymmetric response to disturbance and recovery: Changes of soil permeability under forest-pasture-forest transitions, *Geoderma*, 159, 209–215, doi:10.1016/j.geoderma.2010.07.013, 2010.