

# Areas of residential development in the southern Appalachian Mountains are characterized by low riparian zone nitrogen cycling and no increase in soil greenhouse gas emissions

Peter Baas  · Jennifer D. Knoepp · Daniel Markewitz · Jacqueline E. Mohan

Received: 24 February 2016 / Accepted: 27 February 2017 / Published online: 10 March 2017  
© Springer International Publishing Switzerland 2017

**Abstract** The critical role streamside riparian zones play in mitigating the movement of nitrogen (N) and other elements from terrestrial to aquatic ecosystems could be threatened by residential development in the southern Appalachian Mountains. Many studies have investigated the influence of agriculture on N loading to streams but less is known about the impacts of residential development. Here we consider the dynamics of changing riparian land use in the southern Appalachians that includes increased residential development at the expense of both forests and agriculture. We hypothesized that increased inputs of inorganic N from residential development will

increase nitrogen cycling rates relative to forests, thereby preventing terrestrial N retention and increasing soil nitrate losses through leaching. In addition, we hypothesized that such development will increase emissions of N<sub>2</sub>O, CO<sub>2</sub>, and CH<sub>4</sub>, all potent greenhouse gases. We found riparian soil potential N cycling rates as well as N<sub>2</sub>O and CO<sub>2</sub> efflux to be much greater with agricultural land use compared to either forested or residential land use. Our data suggest that residential development of forested riparian ecosystems does not increase N cycling or removal and, thus, might allow for greater potential N leaching into streams. Both agricultural and residential land use exhibited CH<sub>4</sub> efflux while forested ecosystems were responsible for CH<sub>4</sub> uptake. Overall, regional greenhouse gas emissions are projected to decline as high N<sub>2</sub>O and CO<sub>2</sub> emitting agricultural land is converted to residential use.

---

Responsible Editor: Jack Brookshire.

**Electronic supplementary material** The online version of this article (doi:10.1007/s10533-017-0318-9) contains supplementary material, which is available to authorized users.

---

P. Baas (✉) · J. E. Mohan  
Odum School of Ecology, University of Georgia, 140 E  
Green St, Athens, GA 30602, USA  
e-mail: pbaas83@gmail.com

J. D. Knoepp  
JD Knoepp, USDA Forest Service, Southern Research  
Station, Coweeta Hydrologic Laboratory, 3160 Coweeta  
Lab Road, Otto, NC 28763, USA

D. Markewitz  
D Markewitz, Warnell School of Forestry and Natural  
Resources, University of Georgia, 180 E Green St,  
Athens, GA 30602, USA

**Keywords** Land use change · Nitrogen · Riparian · Carbon dioxide · Methane · Nitrous oxide · Denitrification

## Introduction

Residential land use in the southern Appalachian Mountains has intensified over the last few decades and is expected to continue over subsequent decades (Gragson and Bolstad 2006; Kirk et al. 2012). Traditionally, human development in this region has

consisted of small scale agriculture and low density residential development (Gragson and Bolstad 2006). However, current predictions expect 75% of the new development in the region to be of a suburban nature (Kirk et al. 2012), representing higher density residential areas. By 2030 it is expected that 67% of this development will be on previously forested land (Kirk et al. 2012).

Forested riparian zones (land adjacent to streams) retain nitrogen (N) moving toward the stream (Groffman et al. 2002) and therefore are critical for maintaining water quality. In the southern Appalachian Mountains (Webster et al. 2012) riparian zones are relatively narrow (Clinton et al. 2010) and sensitive to changes in land use (Turner et al. 2003). Indeed, agricultural and suburban land use has been shown to increase stream nitrate ( $\text{NO}_3^-$ ) concentrations substantially compared to forested streams (Kaushal et al. 2011; Webster et al. 2012). In southern Appalachia, streams and rivers were found to be particularly sensitive to conversion of forests resulting in detrimental effects on the water purification services they provide (Rosemond et al. 2015). Under forested conditions, the main source of stream nitrate is microbial mineralization of terrestrial organic N. In contrast, waste water leaching is the dominant source of  $\text{NO}_3^-$  to streams in developed areas (Kaushal et al. 2011). Riparian zone degradation is one of the major causes of decreasing water quality in the US (Faustini et al. 2009).

Riparian ecosystems can retain up to 89% of the N produced from upland anthropogenic activities (Dosskey 2001; Vought et al. 1994) and up to 98% of forest upslope inputs (Fisher and Likens 1973; Pinay et al. 1989). This N retention results in the formation of hotspots for N and C accumulation which can lead to increased ecosystem productivity (Harms and Grimm 2008). Forested riparian zones in the southern Appalachian Mountains with high rates of N mineralization and nitrification have been shown to effectively mitigate nutrient influx associated with upslope disturbances (Knoepp and Clinton 2009). Additionally, redox conditions in the riparian zone are generally favorable for high rates of denitrification, the process responsible for converting inorganic nitrogen into gaseous forms (Davidson and Swank 1986).

Nitrogen loading from upslope positions results in increased rates of riparian N cycling (Groffman et al. 2002), which often coincides with greater emissions of

nitrous oxide ( $\text{N}_2\text{O}$ ) (Groffman et al. 1998; Hefting et al. 2003). Approximately 6% of global terrestrial  $\text{N}_2\text{O}$  emissions come from riparian zones (Bouwman et al. 2013; Groffman et al. 2000). Soils are capable of reducing, and thus consuming,  $\text{N}_2\text{O}$  but uptake <2% of emissions; as such decreasing atmospheric  $\text{N}_2\text{O}$  will likely not result from increased uptake but rather through emission reductions (Schlesinger 2013). In southern Appalachian Mountain riparian zones, pastures used for cattle grazing had emissions as high as  $24.5 \text{ kg N}_2\text{O-N ha}^{-1} \text{ yr}^{-1}$  (Walker et al. 2002); restoration of a vegetative riparian zone through cattle removal and tree planting decreased emissions by 75% (Walker et al. 2009). Further, this restoration method shifted the source of  $\text{N}_2\text{O}$  from nitrification to denitrification (Walker et al. 2009) suggesting that a more complete reduction of  $\text{NO}_3^-$  by denitrification reduced  $\text{N}_2\text{O}$  emissions.

In addition to  $\text{N}_2\text{O}$  efflux,  $\text{CO}_2$  and  $\text{CH}_4$  efflux can be high in riparian zones; particularly as C cycling is coupled with N cycling.  $\text{CO}_2$  emissions are well studied, with soil temperature and moisture being identified as the main controlling factors (Davidson et al. 2002). Processes of  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions, although also responsive to temperature and moisture, are complicated due to their sensitivity to oxygen concentrations and because net efflux rates are the result of both reduction and oxidation reactions (Tiedje et al. 1984). Soil moisture generally stimulates  $\text{CH}_4$  emissions (Schlesinger and Bernhardt 2013), while  $\text{CH}_4$  uptake rates have been found to be both stimulated and inhibited by soil moisture. Maximum  $\text{CH}_4$  uptake rates often occur around 15% volumetric water content (Boeckx and Van Cleemput 1996). In general, soil aeration, regulated by texture and bulk density, controls long-term patterns of  $\text{N}_2\text{O}$  and  $\text{CH}_4$  fluxes, while soil temperature, moisture and substrate availability (e.g. N supply and throughput) determine short-term responses (Werner et al. 2007).

The changes in land use in the southern Appalachians have the potential to alter regional soil nitrogen cycling and emission of greenhouse gases through effects on N cycling processes. For example, conversion of forest to grass dominated riparian systems (i.e. pasture or lawn) could result in increased rates of net soil  $\text{CO}_2$  efflux because grasslands often have a greater  $\text{CO}_2$  soil efflux than forests (Raich and Tufekciogul 2000). Rates of soil  $\text{CH}_4$  uptake from the atmosphere often decreases when forest land is converted to

agricultural use (Powlson et al. 1997). Additionally, while fertilizer additions in agricultural or residential land use generally increase rates of C sequestration (Tian et al. 2014) it also reduces soil CH<sub>4</sub> uptake potential (Stuedler et al. 1989) resulting in net positive CH<sub>4</sub> emissions from agricultural pasture or residential lawns (Law et al. 2004).

The objective of this study was to assess rates of N cycling processes and greenhouse gas emissions in riparian areas under different land uses and elucidate relationships with edaphic drivers. We hypothesized that higher N cycling rates in riparian areas of agricultural and residential development would result in greater N<sub>2</sub>O efflux due to increased rates of nitrification and denitrification relative to upland forested ecosystems. We further proposed that high rates of N cycling in agricultural and residential ecosystems would result in decreased CH<sub>4</sub> uptake and potentially greater CH<sub>4</sub> efflux, as well as greater CO<sub>2</sub> efflux compared to forested riparian areas. We examined these rates of greenhouse gas emissions in riparian zones to estimate the impact of land use change at a regional scale.

## Materials and methods

### Site description

The study was conducted in Macon County, North Carolina, in the Blue Ridge physiographic province in the southern Appalachian Mountains (Fenneman 1917). This region receives an average of 1700 mm of precipitation a year (Coweeta Long Term Ecological Research Database 1936–2013). The highest temperatures are between May and September (20 °C) and the lowest temperatures are between December and February (5 °C) (Laseter et al. 2012). The growing season starts in May and ends in September (Swift and Cunningham 1988). Historically, this region has been dominated by logging and agricultural activities with the majority of the region clear-cut in the early 1900s (Gragson and Bolstad 2006). Small scale agricultural activities occupied 22% of the landscape in the early 1900s and have steadily decreased to 4% as of 2010 (Bolstad and Vose 2005; Kirk et al. 2012). Both logging activities and agricultural land use have been significantly reduced since the 1960s (Gragson and Bolstad 2006).

Currently, forested ecosystems represent about 86% of the riparian cover, however, residential land use (currently about 10%) is becoming more predominant and details of historic, current and projected residential land use have been extensively described in Kirk et al. (2012) and Gragson and Bolstad (2006).

We selected 8 riparian sites that included forest cover (N = 3), undisturbed since the 1920s; agricultural use (N = 3), comprised of managed pasture with and without grazing, and low density residential development (N = 2). All sites were adjacent to a first or second order headwater stream. At each site we established a 20 m wide and 80 m long riparian zone sampling area perpendicular to the stream. Two of the forested sites were on soils predominantly classified as a fine loamy, parasesquic, mesic Typic Hapludult (Evard-Cowee soil series; altitude: 818 and 815 m) while the third was mainly a fine-loamy, mixed, superactive, mesic Humic Hapludult (Saunook soil series; altitude 811 m). The agricultural sites included one Saunook soil series (altitude: 664 m; hay production only), a second with a mix of Evard-Cowee and Saunook soil series (53/47%, respectively, altitude: 661 m; limited cattle grazing with no access to the stream), and a third site classified as a mix of a coarse-loamy, mixed, superactive, mesic Oxyaquic Humudept (60%; Reddies fine sandy loam soil series) and Saunook soil series (27%; altitude: 761 m; cattle grazing with access to the stream). The residential sites were mainly classified as Saunook soil series; altitude: 661 m; tree dominated vegetation) and a mix of fine-loamy, isotic, mesic Typic Humudept (59%; Tuckasegee-Cullasaja soil series) and Evard-Cowee soil series (41%; altitude: 720 m; mix of lawn and tree vegetation). All soil series data were retrieved from the USDA Soil Survey (NRCS Soil Survey Staff 2014).

### Field sampling

Measurements for greenhouse gas emissions were taken eight times between May 2012 and May 2013 (2012: May, July, September, November and December; 2013: February, March and May) at five locations in each site (CH<sub>4</sub> fluxes were only collected from November 2012 through May 2013). At each date, three locations were randomly selected within 3 m from the stream and two additional locations were placed up to 20 meters from the stream and at least 10

m apart. No differences were found based on sampling location along the hillslope and the sampling locations were, therefore, treated as replicates. In May, July, and November of 2012, and March and May of 2013, at each greenhouse gas measurement location, soil cores were collected for laboratory measurement of potential N cycling rates and soil abiotic characteristics, as described below. Soil bulk density was determined in April 2013 for all sites. Using precipitation data retrieved from the NOAA climate database for Macon County, NC (NOAA 1950–2013), we calculated cumulative 24 h and 48 h precipitation before sampling.

At every gas flux sampling date and location, we measured soil moisture and temperature at 5-cm depth in the mineral soil using a Hydrosense sensor (Campbell Scientific Inc., Logan, UT, USA) and soil thermometer, respectively. Soil samples (0–15 cm) were collected for nitrogen (N) cycling assays and edaphic characteristics using a 5-cm diameter stainless steel soil push tube from three randomly determined locations per site; soil samples were stored separately for analyses. We divided core samples into O-horizon (if present), 0–5, 5–10, and 10–15 cm mineral soil and stored samples in sealed plastic bags. Each soil sample was sieved to <2 mm and homogenized. Gravimetric soil moisture was determined by oven drying ~2 g of soil to constant weight at 105 °C. Assessments of available inorganic nitrogen ( $\text{NO}_3^-$  and  $\text{NH}_4^+$ ) were determined using 2 M KCl extractions in March 2013 on freshly sieved soils for the top 10 cm (USEPA 1983a, b). Soil pH in water (2:1) was determined on air-dried soils as described in Robertson et al. (1999) in July 2012, November 2012, March and May 2013. Soil bulk density ( $\text{g soil cm}^{-3}$ ) of the surface 0–15 cm was determined on air dried soils collected in April 2013 using a 4.3 cm diameter PVC pipe. These soils were also used for C and N contents using a Flash EA 1112 NC analyzer (Thermo Scientific).

### Nitrogen cycling assays

We determined potential nitrification (pNTR) and potential denitrification (pDNF) on fresh soil samples using oxic and anoxic incubation techniques, respectively. pNTR was determined within 48 h of soil collection on field moist soils stored at 4 °C and pDNF was conducted on soils stored (4 °C) for less 7 days.

To assess pNTR we placed five grams of sieved soil (<2 mm) from each sample location near the stream ( $N = 8$  plots (3 forest + 3 agriculture + 2 residential)  $\times 10$  plots per site = 80 samples) in separate 37 mL serum vials with 15 mL of media ( $0.33 \text{ g L}^{-1}$   $(\text{NH}_4)_2\text{SO}_4$  and buffered with  $0.14 \text{ g L}^{-1}$   $\text{K}_2\text{HPO}_4$  and  $0.027 \text{ g L}^{-1}$   $\text{KH}_2\text{PO}_4$  in DI water). Each serum vial was wrapped and capped with aluminum foil to prevent evaporation and UV light inhibition of ammonia-oxidizing bacteria. After addition of the media, vials were gently shaken at 10 relative centrifugal force at 25 °C; 2 mL sub-samples were collected after 0.5, 2, ~7 (between 6 and 8 h) and 24 h of incubation using a cut-off pipette tip (to facilitate pipetting a slurry). Samples were centrifuged for 10 min at 11,000 rcf and the supernatant was immediately frozen at -20 °C until thawing for  $\text{NO}_2^- + \text{NO}_3^-$  analysis using colorimetric methods (Bendschneider and Robinson 1952). pNTR rates were determined by linear regression analysis of changes in KCl solution  $\text{NO}_3^-$  concentration over time and are presented per soil dry weight. All regressions were significant with  $r^2 \geq 0.8$ . Based on the depth of the horizon sampling and its bulk density we determined total rate per gram of soil for each core.

Anoxic pDNF rates were determined using the acetylene block method (Groffman et al. 1999). Five grams of sieved soil (just 1 g for O-horizon samples due to limited sample) was added to 37 mL serum vials. Serum vials were purged with He for 1 min to displace oxygen from the vial, and then 5 mL of incubation media was added to the serum vials. Media consisted of dextrose (1 mM) and sodium nitrate (1 mM) in DI water purged for 30 min with He. Assays were initiated by replacing 4 mL of headspace with 99% pure acetylene (10% v/v). Samples were incubated at 20 °C while shaking (150 rpm) for 3 h. Gas subsamples were taken after 0.5 and 3 h, and stored in 3.5 mL vacutainers (Labco, Lampeter, UK) until analysis for  $\text{N}_2\text{O}$  on a GC-ECD with a 10-port Valco valve to prevent acetylene from saturating the detector. Rates of pDNF were determined via regression analysis of changes in headspace  $\text{N}_2\text{O}$  concentrations over time and calculated on a soil dry weight basis. All regressions were significant with  $r^2 \geq 0.8$ . Based on the depth of the horizon sampling and its bulk density we determined total rate per gram of soil for each core.

## Greenhouse gas emissions

We measured net soil CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes between 09:00 and 16:00 h using static PVC flux chambers with an inner diameter of 15.1 cm and a height of 8 cm. The PVC collars were installed 1 h before flux determination to a soil depth of 10 cm. PVC lids were adapted with septa for gas sampling; 9 mL gas samples were taken using a ten cc plastic syringe at 1, 5, 10 and 30 min after placing the lid on the collar. Gas samples were analyzed for CO<sub>2</sub> using a LICOR-7000 (LICOR Inc., Lincoln, NE, USA) and for N<sub>2</sub>O and CH<sub>4</sub> using a Shimadzu GC-ECD and GC-FID (Shimadzu Co., Kyoto, Japan), respectively. Fluxes were determined using linear regression analysis. Rates are presented as g C m<sup>-2</sup> d<sup>-1</sup> (CO<sub>2</sub>), mg N m<sup>-2</sup> d<sup>-1</sup> (N<sub>2</sub>O), and mg C m<sup>-2</sup> d<sup>-1</sup> (CH<sub>4</sub>). We assumed flux measurements to be representative for the whole day as was found by Bremer et al. (1998). We used a trapezoidal integration approach to estimate a cumulative flux rate from July 2012–May 2013 (CO<sub>2</sub> and N<sub>2</sub>O) and November 2012 through May 2013 (CH<sub>4</sub>). Total riparian area (20 meters adjacent to a stream) was estimated using the buffer tool in ArcGIS 10.0 on the Coweeta LTER stream GIS data (Coweeta Long Term Ecological Research Database 1936–2013). We used our annual gas flux estimates to calculate changes in the contribution of the riparian area in CO<sub>2</sub> equivalents (CO<sub>2</sub>-eq) (CH<sub>4</sub> = 86-CO<sub>2</sub>-eq; N<sub>2</sub>O = 268-CO<sub>2</sub>-eq (Myhre 2013)) fluxes over a 20 year timespan for Macon County, North Carolina based on projected land use changes from 2010–2030 (Kirk 2009). To provide an assessment of variance we have propagated the error of the annual fluxes.

## Statistics

Greenhouse gas fluxes, potential N cycling rates and edaphic characteristics measured over time were tested for significant land use effects using a mixed model repeated measures approach restricted maximum likelihood (REML). Date and land use type were fixed effects, site was a random effect (Random: site, as well as interaction terms). We used Tukey post hoc mean separation test to identify significant differences among dates and land use type. If date and land use type interactions were significant, we tested land use type effect for each sample date using one-way ANOVA and Tukey post hoc mean separation


analyses. Data were either log transformed or rank transformed (Conover and Iman 1981) if needed to acquire a normal distribution for parametric analyses. All errors depicted represent the standard error of the mean. The relationship between process drivers and gas flux rates were determined using Spearman correlation analysis on non-transformed data to determine predictors for greenhouse gas fluxes and N cycling rates. All descriptions of uncertainty are in the form of standard error (SE) of the mean calculated by taking an average of the respective site and determining the SE of the 2 (Residential) or 3 sites (Agriculture and Forested). All statistical analyses were conducted in JMP 11 (SAS Institute Inc., Cary, NC) and significant differences are indicative of  $P < 0.05$  unless otherwise stated.

## Results

### Edaphic characteristics

Land use had a significant effect on soil temperature (Fig. 1; Table 1). Soil temperatures were greater in agricultural sites ( $15.4 \pm 0.55$  °C) compared to forested ( $12.4 \pm 0.45$  °C) and residential ( $13.5 \pm 0.67$  °C) sites on seven out of eight sample dates, with the greatest temperatures in July 2012 ( $24.0 \pm 0.42$  °C). Land use (Fig. 1; Table 1) showed no significant effect on soil moisture but sample dates differed significantly; land use and sample date interaction also tended to be significant ( $p = 0.062$ ). Agricultural land use had greater soil moisture in five out of the eight sampling times (Fig. 1; Table 1).

Land use had a significant effect on soil pH, total N and extractable NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> concentrations and bulk density of the surface 15 cm; land use and date interactions were not significant. Soil pH and total NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup> concentrations were greater in soils with agricultural land use compared to both residential and forested lands (Table 2); residential and forested sites were not significantly different. Soil pH was greater in agricultural land use compared to residential and forested land use for November 2012, March and May 2013 while both residential and agricultural soil pH were greater than forested in July 2012. Soil total N differed among sites and was greatest in agricultural sites, followed by residential and then forested land use types (Table 2). ( $F_{2,6} = 11.5, p = 0.01$ ; Table 2).



**Fig. 1** Volumetric soil moisture and soil temperature in riparian forests of the southern Appalachian Mountains. The points indicate the mean and the error bars indicate the standard error of the mean ( $n = 2-3$ ). Asterisk indicates Ag is significantly greater than Forest and Res ( $P < 0.05$ ). Forest forested riparian zones, Res riparian zones in residential development, Ag riparian zones under agricultural development

Soil bulk density was greater in agricultural sites than forested and residential sites ( $F_{2,5} = 12.2$ ,  $P < 0.01$ ). Sample date had a significant effect on pH; pH was greater in May 2013 compared to July 2012. Land use

effect on total C concentration was not significant (Table 2).

### Nitrogen cycling

pNTR and pDNF rates were significantly affected by both land use and sample date and ranged from  $-0.2$  to  $3.4 \mu\text{g N g}_{\text{soil}}^{-1} \text{d}^{-1}$  and  $-0.2$  to  $122 \mu\text{g N g}_{\text{soil}}^{-1} \text{d}^{-1}$ , respectively (Fig. 2; Table S1). pNTR had a significant interaction between land use and sample date (Table 1;  $F_{2,5} = 67.0$ ,  $P < 0.001$ ). pNTR rates were greater in agricultural sites compared to forested and residential sites in May 2012 ( $F_{6,14} = 5.1$ ,  $P < 0.05$ ), November 2012 ( $F_{7,16} = 13.9$ ,  $P < 0.001$ ), March 2013 ( $F_{7,16} = 7.4$ ,  $P < 0.001$ ) and May 2013 ( $F_{7,24} = 4.8$ ,  $P < 0.001$ ); post hoc mean separation tests did not identify significant differences among sample dates. pDNF rates were greatest in agricultural sites; residential sites were greater than forests. July 2012 had the greatest pDNF rates while May 2013 had the lowest. O-horizon pNTR (Table S2) was responsible for  $19 \pm 11\%$  (Forested) to  $3 \pm 2\%$  (Res) of the overall site activity and  $24 \pm 4\%$  (Forested) and  $10 \pm 21\%$  (Res) of the pDNF activity (Table S2).

### Greenhouse gas emissions and land use

The riparian soil  $\text{CO}_2$  gas fluxes exhibited significant land use, sampling date and land use by sampling date interaction effects (Fig. 3a; Table 1).  $\text{CO}_2$  flux was greater in Agricultural land use compared to


**Table 1** Results for edaphic characteristics and greenhouse gas fluxes tested with linear mixed model statistics for land use and date

Variable	Land-use			Sampling date			Land-use $\times$ sampling date		
	df, $df_{\text{error}}$	F	P	df, $df_{\text{error}}$	F	P	df, $df_{\text{error}}$	F	P
Edaphic characteristics									
Soil temperature	2, 5	8.6	0.023	7, 271	580.0	<0.001	14, 271	9.7	<0.001
Soil moisture	2, 5	3.5	0.111	7, 260	26.1	<0.001	14, 260	1.7	0.062
pH	2, 5	15.7	0.007	3, 138	4.3	0.006	6, 138	1.1	0.37
Fluxes									
$\text{CO}_2$	2, 5	30.7	0.001	7, 258	20.9	>0.001	14, 258	2.0	0.017
$\text{CH}_4$	2, 5	3.4	0.12	4, 160	1.7	0.15	8, 160	1.3	0.23
$\text{N}_2\text{O}$	2, 8	3.6	0.080	7, 124	3.1	0.005	14, 123	0.53	0.91
Potential N-cycling									
Nitrification	2, 5	6.1	0.044	4, 106	1.5	0.21	8, 105	2.5	0.016
Denitrification	2, 5	67.0	<0.001	4, 169	18.5	<0.001	8, 169	0.8	0.58

**Table 2** Soil properties (mean±SE) for bulk density (BD), pH, total carbon (C) and nitrogen (N), and extractable  $\text{NH}_4^+$  and  $\text{NO}_3^-$  in the top 15 cm of mineral soil in the southern Appalachian Mountains

	BD ( $\text{g}/\text{cm}^3$ )	pH	C ( $\text{mg}/\text{g}$ )	N ( $\text{mg}/\text{g}$ )	$\text{NH}_4^+$ ( $\text{mg N kg soil}^{-1}$ )	$\text{NO}_3^-$ ( $\text{mg N kg soil}^{-1}$ )
For	$0.65 \pm 0.06^b$	$5.0 \pm 0.40^b$	$26.1 \pm 1.3$	$1.1 \pm 0.1^c$	$2.3 \pm 0.6^b$	$0.05 \pm 0.04^b$
Res	$0.69 \pm 0.05^b$	$5.2 \pm 0.51^b$	$27.2 \pm 1.4$	$1.6 \pm 0.1^b$	$3.0 \pm 0.9^b$	$0.8 \pm 0.6^b$
Ag	$1.08 \pm 0.08^a$	$5.9 \pm 0.42^a$	$24.6 \pm 1.3$	$2.1 \pm 0.1^a$	$9.3 \pm 4.3^a$	$5.9 \pm 2.3^a$

Total C and N were determined using an Elementar Flash EA 1112 NC analyzer.  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations are shown for March 2013. For = forest (N=3); Res = residential (N=2); Ag = agricultural (N=3); BD = bulk density. The letters indicate significant differences between land use types



**Fig. 2** Potential nitrification (a) and potential denitrification (b) rates for the different sampling dates (total including O-horizon and mineral soil 0–15) in the southern Appalachian Mountains. The symbols indicate the mean and the error bars indicate the standard error of the mean ( $n = 2-3$ ). Forest


forested riparian zones, Res riparian zones in residential development; Ag riparian zones under agricultural development. Asterisk indicates Ag is significantly greater than Forest and Res ( $P < 0.05$ )

residential land use in four of the eight sampling times, and agricultural land use was greater than forested systems in six out of the eight sampling times. Residential site  $\text{CO}_2$  flux was greater than forested sites at only one sample date. Forest riparian sites showed a negative net  $\text{CH}_4$  flux, (i.e.  $\text{CH}_4$  consumption) at all sample times, while agricultural and residential sites varied between  $\text{CH}_4$  production and  $\text{CH}_4$  consumption. However, due to high variance, residential sites differed significantly from forested sites with a net  $\text{CH}_4$  efflux in only one sample date (November 2012;  $F_{7,30} = 2.3$ ,  $P = 0.050$ ) and agricultural  $\text{CH}_4$  flux rates did not differ from either forested or residential land use sites. Riparian  $\text{N}_2\text{O}$  fluxes varied greatly among sites, ranging from the least in forested sites ( $-12 \pm 11 \text{ mg N m}^{-2} \text{ y}^{-1}$ ) to the greatest in agriculture sites ( $428 \pm 316 \text{ mg N m}^{-2} \text{ y}^{-1}$ ), however, no significant land use effect was found. On the other hand, significant differences were observed between sampling dates where  $\text{N}_2\text{O}$  fluxes in

May 2012 and July 2012 were greater than March 2013. The cumulative annual riparian  $\text{CO}_2$  flux differed significantly with land use with a greater calculated annual flux in agricultural land compared to both residential and forested land use ( $F_{2,5} = 22.1$ ,  $P = 0.003$ ) (Table 3). Conversely, cumulative riparian  $\text{CH}_4$  (Table 3;  $F_{2,5} = 1.2$ ,  $P = 0.39$ ) and riparian  $\text{N}_2\text{O}$  fluxes (Table 3;  $F_{2,5} = 0.4$ ,  $P = 0.71$ ) showed no significant land use effect. Currently, emissions of  $\text{CO}_2$  account for greater than 97% of the total  $\text{CO}_2$ -eq of greenhouse gas emissions from riparian zones in this study;  $\text{N}_2\text{O}$  flux was relevant only in agricultural sites, contributing  $\sim 3\%$ , and  $\text{CH}_4$  flux was insignificant ( $<1\%$ ).

Land use in Macon County, North Carolina is predicted to shift, with agricultural use declining 0.4%, residential use increasing 2.6% and forest decreasing 2.2% between 2010 and 2030 (Fig. 4) (Kirk et al. 2012). These land use changes are used to estimate changes in annual riparian greenhouse gas

**Fig. 3** CO<sub>2</sub> (a), CH<sub>4</sub> (b) and N<sub>2</sub>O (c) fluxes. The bars indicate the mean and the error bars indicate the standard error of the mean. *Forest* forested riparian zones, *Res* riparian zones in residential development, *Ag* riparian zones under agricultural development (n = 2–3)




**Table 3** Cumulative fluxes (mean  $\pm$  SE) for CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O for the three land use types in the southern Appalachian Mountains

	CO <sub>2</sub> (kg C m <sup>-2</sup> )	CH <sub>4</sub> (mg C m <sup>-2</sup> )	N <sub>2</sub> O (mg N m <sup>-2</sup> )
For	1.2 $\pm$ 0.1 <sup>b</sup>	-346 $\pm$ 101 <sup>b</sup>	-19 $\pm$ 9
Res	1.0 $\pm$ 0.0 <sup>b</sup>	247 $\pm$ 426 <sup>a</sup>	36 $\pm$ 8
Ag	2.0 $\pm$ 0.5 <sup>a</sup>	58 $\pm$ 97 <sup>a</sup>	385 $\pm$ 283

For = forest; Res = residential; Ag = pasture. Fluxes are July 2012–May 2013 (CO<sub>2</sub> and N<sub>2</sub>O) and November 2012–May 2013 (CH<sub>4</sub>). The letters indicate significant differences between land use types ( $P < 0.05$ )

emissions. Our estimates suggest that riparian soil CH<sub>4</sub> efflux will increase while both CO<sub>2</sub> and N<sub>2</sub>O fluxes will decrease (Fig. 4). Overall changes in land use could result in an overall decrease in regional riparian CO<sub>2</sub>-eq emissions by 0.7% during the 2010–2030 time period. Reductions in riparian CO<sub>2</sub> and N<sub>2</sub>O emissions due to decreasing land area in agricultural use accounted for most of the reduction in greenhouse gas equivalents. On the other hand, county-wide riparian CH<sub>4</sub> emissions are predicted to



**Fig. 4** Projected land use changes in the southern Appalachian Mountains from 2010–2030 based on Kirk (2009) and estimated regional riparian CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O fluxes for Macon County, NC. Greenhouse gas emissions (GHG) for all of Macon County's riparian zones are presented in CO<sub>2</sub>-equivalent fluxes. The bars indicate the estimated change in area or GHG emission rate. *Forest* forested riparian zones, *Res* riparian zones in residential development, *Ag* riparian zones under agricultural development

increase due to increasing land area present in riparian residential land use, thus decreasing the CH<sub>4</sub> consuming forested ecosystems.

## Drivers of fluxes and processes

Spearman correlation analyses (Table 4) found CO<sub>2</sub> fluxes to be positively correlated to soil temperature, pH and potential denitrification in all land use types. CO<sub>2</sub> flux was negatively correlated with 24 h cumulative precipitation in the forested sites only. CH<sub>4</sub> fluxes were positively correlated to soil temperature, soil moisture and potential denitrification. Analysis by land use type found a positive correlation between CH<sub>4</sub> efflux and soil moisture in residential land use. N<sub>2</sub>O fluxes were positively correlated to soil temperature, pH, 48 h cumulative precipitation and potential nitrification and denitrification.

Correlation analysis between pNTR and pDNF and edaphic variables showed positive correlation with soil pH (pNTR, Pearson correlation coefficient ( $\rho$ ) = 0.39; pDNF,  $\rho$  = 0.58;  $P < 0.05$ ) (data not shown). Soil temperature at the time of sample collection was significantly correlated with pDNF rates ( $\rho$  = 0.33;  $P < 0.01$ ) but had no relationship with pNTR. March 2013 NO<sub>3</sub><sup>-</sup> concentrations showed a significant positive relationship with both pNTR ( $\rho$  = 0.89;  $P < 0.01$ ) and pDNF ( $\rho$  = 0.69;  $P < 0.01$ ) and NH<sub>4</sub><sup>+</sup> with pDNF ( $\rho$  = 0.46;  $P < 0.05$ ).

## Discussion

We estimated the consequences of residential development on previously agricultural or forested lands on riparian nitrogen transformations and greenhouse gas emissions in the southern Appalachian Mountains. This study suggests that the future will likely be characterized by lower nitrogen cycling rates, possibly less soil N retention and greater CH<sub>4</sub> emissions. Our data also indicate that due to a projected decrease in high greenhouse gas emitting agricultural lands, there would be no net effect of increased residential development on regional greenhouse gas forcing.

## Land use effects on edaphic characteristics

The manner in which edaphic characteristics (soil bulk density, extractable NH<sub>4</sub><sup>+</sup> and NO<sub>3</sub><sup>-</sup>, and total N) varied among different land use types in our study is similar to what has been found by others (Groffman et al. 1998; Walker et al. 2009). However, our finding of no land use effect on soil moisture dynamics contrasts with a study by Groffman et al. (2002) in an urban setting (Baltimore, MD USA) that showed significantly reduced soil moisture in sites with residential land use compared to forested sites; reductions were linked to decreased soil microbial

**Table 4** Spearman rank correlations for greenhouse gas efflux rates and soil and environmental variables

Correlation variable	CO <sub>2</sub>				CH <sub>4</sub>				N <sub>2</sub> O			
	All	For	Res	Ag	All	For	Res	Ag	All	For	Res	Ag
Temperature	<b>0.63**</b>	<b>0.60**</b>	<b>0.43**</b>	<b>0.71**</b>	<b>0.16*</b>	0.08	0.13	0.02	<b>0.19*</b>	0.01	0.12	<b>0.21*</b>
Soil moisture	0.06	<b>-0.24*</b>	-0.08	-0.06	<b>0.20**</b>	0.00	<b>0.38</b>	-0.09	0.03	0.06	-0.17	<b>-0.20†</b>
pH	<b>0.33**</b>	-0.08	0.24	<b>-0.24†</b>	0.16	0.14	-0.06	-0.06	<b>0.24**</b>	0.10	0.11	-0.14
Precip 48 h	-0.04	-0.15	-0.06	-0.14	0.10	0.17	-0.14	0.04	<b>0.14*</b>	0.02	0.10	0.09
Precip 24 h	<b>-0.14*</b>	<b>-0.18†</b>	-0.20	<b>-0.18†</b>	0.06	0.15	-0.19	0.17	0.07	0.04	0.00	0.01
pNTR	0.18	-0.09	0.13	<b>-0.31*</b>	0.18	0.11	-0.33	0.05	<b>0.22*</b>	0.01	0.25	0.04
pDNF	<b>0.48**</b>	<b>0.24†</b>	0.12	0.15	<b>0.28**</b>	-0.03	0.18	0.15	<b>0.30**</b>	-0.08	-0.14	<b>0.27*</b>

The values indicate the correlation coefficient  $\rho$  (rho). Coefficients with  $p < 0.1$  are shown in bold

For forested sites, Res residential sites, Ag agricultural sites, pNTR potential nitrification rates 0–15 cm mineral soil, pDNF potential denitrification rates 0–15 cm mineral soil

\*  $P < 0.05$

\*\*  $P < 0.01$

†  $P < 0.1$

activity (Groffman et al. 2002). On the other hand, our soil temperature data are consistent with observations by Davidson et al. (1998) where land use effects on soil temperature were significant and temperatures in agricultural lands were greater compared to forested land; residential developments were in between.

#### Land use effects on nitrogen cycling

Studies by Groffman et al. (2000) and Hefting et al. (2003) in urban/residential riparian zones were characterized by large nitrogen inputs along with high nitrification and denitrification rates. In our study system residential land use resulted in greater total soil N concentration but extractable N ( $\text{NH}_4^+$  and  $\text{NO}_3^-$ ) along with pNTR and pDNF rates did not differ from forested systems. However, agricultural riparian systems had greater extractable and total N and exhibited pNTR and pDNF rates greater than both residential and forested sites; rates that were similar to those found in other studies (Groffman et al. 2000; Hefting et al. 2003). This suggests that while southern Appalachian residential riparian ecosystems may retain some of the additional N inputs, as evidenced by increased total N, N processing rates did not increase significantly. Observed increases in stream  $\text{NO}_3^-$  with increasing residential development (Webster et al. 2012) support this limited N retention hypothesis. A lack of riparian soil N retention combined with the demonstrated sensitivity of southern Appalachian in-stream processes to nutrient (N and P) additions (Rosemond et al. 2015) indicates stream ecosystem functions could be threatened by future residential development of forested ecosystems. The O-horizon, present only in the forested systems, was responsible for a large proportion of N cycling activity, demonstrating the important role organic horizons play in N processing and, thus, preventing N movement into streams under predominantly forested conditions.

#### Greenhouse gas emissions and land use change

Riparian forest  $\text{CO}_2$  emissions in this study exceeded previous estimates for similar riparian forests in the region (Bolstad and Vose 2005). Similar rates (2.0–2.8  $\text{kg C m}^{-2} \text{y}^{-1}$ ), however, have been reported for forests generally in the region (Vose and Bolstad 2007).  $\text{CO}_2$  flux rates for agricultural systems in our

study were approximately three times greater than observed by Bolstad and Vose (2005). Although soil temperature was important in explaining  $\text{CO}_2$  flux rates, an increase in soil moisture and high root density in pastures were also important likely drivers (Kellman et al. 2007). Our data puts forward the idea that the higher soil moistures and input of labile carbon from grasses in agricultural sites could contribute to higher  $\text{CO}_2$  fluxes compared to forest sites. Previous studies have shown that  $\text{CO}_2$  emissions were correlated with both temperature and soil moisture (Davidson et al. 2012). We hypothesized that the absence of a strong moisture effect in our study might be due to the relatively high rainfall of 2100 mm during our study period (July 2012–June 2013) compared to an average of 1700 mm for the ten prior years (Coweeta Long Term Ecological Research Database 1936–2013), which may have alleviated any soil moisture limitation. The negative correlation between total precipitation 24 h prior and  $\text{CO}_2$  flux measurements is likely the result of reduced diffusion rates for both  $\text{O}_2$  into and  $\text{CO}_2$  out of the soil, reducing assessed respiration rates (Davidson et al. 2012). The reduction in  $\text{CO}_2$  flux with decreasing soil pH in this study has been found in several previous studies and has been attributed to higher maintenance costs for microbes under more acidic conditions (Anderson and Domsch 1993) and potentially a shift to a fungal dominated community characterized by lower  $\text{CO}_2$  production per unit microbial biomass (Blagodatskaya and Anderson 1998).

The forested riparian ecosystems in this study were distinguished from the other land uses in consistently consuming  $\text{CH}_4$  (i.e. a net negative flux). In the southern Appalachian Mountains these riparian forests can be considered hotspots of  $\text{CH}_4$  uptake (Le Mer and Roger 2001), which, when converted to either agricultural or residential development may become a  $\text{CH}_4$  source.  $\text{CH}_4$  consumption rates measured in the forested ecosystems in our study were on par with the high uptake rates found in tropical forests (190–700  $\text{mg C m}^{-2} \text{y}^{-1}$ ) (Verchot et al. 2000). We measured  $\text{CH}_4$  production rates in both the residential and agriculture ecosystems of  $\sim 3 \text{ mg C m}^{-2} \text{y}^{-1}$ , rates similar to those measured in loam grassland soils (Boeckx et al. 1997). Within each land use type,  $\text{CH}_4$  flux was best explained by a positive correlation with soil temperature (across all sites) and soil moisture (residential only); a finding that is consistent with

other studies (Verchot et al. 2000, Werner et al. 2007). Soil CH<sub>4</sub> consumption may also partly respond to forest density. In the urban forests of Baltimore CH<sub>4</sub> consumption was 30% less compared to rural or suburban forests (Goldman et al. 1995). Here we find a similar response in that the riparian zones of the residential areas are partly forested but consume less CH<sub>4</sub> than the undeveloped riparian forests.

We found biologically relevant N<sub>2</sub>O fluxes only in the agricultural ecosystems ( $\sim 500 \text{ mg N m}^{-2} \text{ y}^{-1}$ ). Fluxes measured in our study were comparable to emissions measured by others in riparian zones exposed to nutrient inputs (Groffman et al. 1998; Hefting et al. 2003) and within other agricultural ecosystems (Matson et al. 1998). High rates of  $2000 \text{ mg N m}^{-2} \text{ y}^{-1}$  measured in high nitrate loaded forested riparian buffers adjacent to agricultural land in the Netherlands (Hefting et al. 2003) were not equaled here. Similar to other studies, N<sub>2</sub>O production was positively correlated with soil temperature (Smith et al. 1998). N<sub>2</sub>O was also positively correlated with pH, although this result may potentially be an artifact of higher soil pH in the agricultural sites as a negative relationship between pH and N<sub>2</sub>O fluxes is expected based on greater N<sub>2</sub>O:N<sub>2</sub> ratios of gaseous production under acidic conditions (Van den Heuvel et al. 2011). On the other hand, a higher pH generally also corresponds to a greater total denitrification rate that may enhance N<sub>2</sub>O production. Higher precipitation over the previous 48 h was weakly but significantly associated with higher N<sub>2</sub>O emissions and could indicate a lag effect in the microbial response to increases in soil moisture (Sexstone et al. 1985). Over all sites and in the sites with agricultural land use, potential denitrification proved the best predictor for N<sub>2</sub>O emissions and, therefore, denitrification is the most likely pathway for N<sub>2</sub>O production in the studied ecosystems.

### Scaling-up to Macon County

Our emissions results coupled with projections of regional land use change (Kirk et al. 2012) estimate that overall CO<sub>2</sub> and N<sub>2</sub>O emissions will decrease by 2030 due to a reduction in the total riparian area with agricultural land use. However, the replacement of forests with residential development may effectively shift riparian zones from a net CH<sub>4</sub> sink into a net source. The combined effect of the CO<sub>2</sub> and N<sub>2</sub>O

decrease and the CH<sub>4</sub> increase, however, is a net reduction in the CO<sub>2</sub>-eq of greenhouse gas emissions. Additionally, the deforestation due to increasing residential development, will result in the removal of a large C pool in the form of aboveground biomass (Bolstad and Vose 2005) and reduce future CO<sub>2</sub> sequestration (Soussana et al. 2004). Similarly, the effect of forest removal on greenhouse gas emission depends on post-harvest usage (e.g. furniture, bioenergy, etc.) but these components are beyond the scope of the present study. Although the error of cumulative CO<sub>2</sub>-equivalent emissions were about half of the mean change in emissions our data suggest that, solely based on riparian zone soil greenhouse gas emissions in this study, the projected reduction of agricultural land cover is predicted to reduce overall regional gas emissions.

### Conclusions

Agricultural land use supports the largest fluxes of greenhouse gases in southern Appalachian Mountain riparian zones. Projected changes in land use suggest that conversion of agricultural land to residential development will result in decreased greenhouse gas emissions. On the other hand, increases in residential development at the expense of forested land, results in a moderate increase in emissions by transforming soils from a net CH<sub>4</sub> sink into a CH<sub>4</sub> source. Hence, overall, projected patterns of land use change would result in no net alteration of total riparian CO<sub>2</sub>-eq greenhouse gas emissions. Our results also found that riparian zone N cycling and removal was largely driven by denitrification. However, conversion of forested riparian zones to residential development with increased N inputs due to fertilization and septic system installation resulted in no increase in the N removal capacity resulting in potentially negative effects on water quality.

**Acknowledgements** We thank Megan Machmuller and Kate Helmick for assistance in the field and we thank Richard Conant for providing insightful comments on an earlier version of the manuscript. Chemical analyses were conducted at the University of Georgia Odum School of Ecology Analytical lab, at the USDA Forest Service, Coweeta Hydrologic Laboratory and Analytical Lab. This research was supported by the National Science Foundation through the Coweeta LTER (DEB-0823293) and by the USDA Forest Service, Southern Research Station, Coweeta Hydrologic Laboratory project funds.

## References

- Anderson T-H, Domsch K (1993) The metabolic quotient for CO<sub>2</sub> (qCO<sub>2</sub>) as a specific activity parameter to assess the effects of environmental conditions, such as pH, on the microbial biomass of forest soils. *Soil Biol Biochem* 25(3):393–395
- Bendschneider K, Robinson RI (1952) A new spectrophotometric method for the determination of nitrite in sea water. *J Mar Res* 11:87–96
- Blagodatskaya EV, Anderson T-H (1998) Interactive effects of pH and substrate quality on the fungal-to-bacterial ratio and qCO<sub>2</sub> of microbial communities in forest soils. *Soil Biol Biochem* 30(10):1269–1274
- Boeckx P, Van Cleemput O (1996) Methane oxidation in a neutral landfill cover soil: influence of moisture content, temperature, and nitrogen-turnover. *J Environ Qual* 25(1):178–183
- Boeckx P, Van Cleemput O, Villaralvo I (1997) Methane oxidation in soils with different textures and land use. *Nutr Cycl Agroecosyst* 49(1–3):91–95
- Bolstad PV, Vose JM (2005) Forest and pasture carbon pools and soil respiration in the southern Appalachian Mountains. *Forest Sci* 51(4):372–383
- Bouwman AF, Beusen AHW, Griffioen J, Van Groenigen JW, Hefting MM, Oenema O, Van Puijenbroek PJTM, Seitzinger S, Slomp CP, Stehfest E (2013) Global trends and uncertainties in terrestrial denitrification and N<sub>2</sub>O emissions. *Philos Trans R Soc B* 368(1621):20130112
- Bremer DJ, Ham JM, Owensby CE, Knapp AK (1998) Responses of soil respiration to clipping and grazing in a tallgrass prairie. *J Environ Qual* 27(6):1539–1548
- Clinton BDCBD, Vose JMVJM, Knoepp JDKJD, Elliott KJEKJ, Reynolds BCRBC, Zarnoch SJZSJ (2010) Can structural and functional characteristics be used to identify riparian zone width in southern Appalachian headwater catchments? *Can J For Res* 40(2):235–253
- Conover WJ, Iman RL (1981) Rank transformations as a bridge between parametric and nonparametric statistics. *Am Stat* 35(3):124–129
- Coweeta Long Term Ecological Research Database (1936–2013) <http://coweeta.uga.edu/datacatalog>. In: Stephanie Lassiter, USDA Forest Service
- Davidson E, Belk E, Boone RD (1998) Soil water content and temperature as independent or confounded factors controlling soil respiration in a temperate mixed hardwood forest. *Glob Chang Biol* 4(2):217–227
- Davidson EA, Savage K, Bolstad P, Clark DA, Curtis PS, Ellsworth DS, Hanson PJ, Law BE, Luo Y, Pregitzer KS (2002) Belowground carbon allocation in forests estimated from litterfall and IRGA-based soil respiration measurements. *Agric Meteorol* 113(1–4):39–51
- Davidson EA, Samanta S, Caramori SS, Savage K (2012) The Dual Arrhenius and Michaelis-Menten kinetics model for decomposition of soil organic matter at hourly to seasonal time scales. *Glob Chang Biol* 18(1):371–384
- Dosskey MG (2001) Toward quantifying water pollution abatement in response to installing buffers on crop land. *Environ Manag* 28(5):577–598
- Faustini JM, Kaufmann PR, Herlihy AT, Paulsen SG (2009) Assessing stream ecosystem conditions in the United States. *EOS* 90(36):309–320
- Fenneman NM (1917) Physiographic subdivision of the United States. *Proc Natl Acad Sci* 3(1):17–22
- Fisher SG, Likens GE (1973) Energy flow in Bear Brook, New Hampshire: an integrative approach to stream ecosystem metabolism. *Ecol Monogr* 43(4):421–439
- Goldman MB, Groffman PM, Pouyat RV, McDonnell MJ, Pickett ST (1995) CH<sub>4</sub> uptake and N availability in forest soils along an urban to rural gradient. *Soil Biol Biochem* 27(3):281–286
- Gragson T, Bolstad P (2006) Land use legacies and the future of southern Appalachia. *Soc Nat Resour* 19(2):175–190
- Groffman P, Gold A, Jacinthe P-A (1998) Nitrous oxide production in riparian zones and groundwater. *Nutr Cycl Agroecosyst* 52(2–3):179–186
- Groffman P, Holland E, Myrold D, Robertson G, XiaoMing Z, Coleman D, Bledsoe C, Sollins P (1999) Denitrification. In: Robertson GP, Coleman DC, Bledsoe CS, Sollins P (eds) *Standard soil methods for long-term ecological research*. Oxford University Press, New York, pp 272–288
- Groffman P, Gold A, Addy K (2000) Nitrous oxide production in riparian zones and its importance to national emission inventories. *Chemosph Glob Chang Sci* 2(3–4):291–299
- Groffman PM, Boulware NJ, Zipperer WC, Pouyat RV, Band LE, Colosimo MF (2002) Soil nitrogen cycle processes in urban riparian zones. *Environ Sci Technol* 36(21):4547–4552
- Harms T, Grimm N (2008) Hot spots and hot moments of carbon and nitrogen dynamics in a semiarid riparian zone. *J Geophys Res* 113:G101020
- Hefting M, Bobbink R, de Caluwe H (2003) Nitrous oxide emission and denitrification in chronically nitrate-loaded riparian buffer zones. *J Environ Qual* 32(4):1194
- Kaushal SS, Groffman PM, Band LE, Elliott EM, Shields CA, Kendall C (2011) Tracking nonpoint source nitrogen pollution in human-impacted watersheds. *Environ Sci Technol* 45(19):8225–8232
- Kellman L, Beltrami H, Risk D (2007) Changes in seasonal soil respiration with pasture conversion to forest in Atlantic Canada. *Biogeochemistry* 82(1):101–109
- Kirk R (2009) Land use and terrestrial carbon storage in Western North Carolina from 1850–2030: a historical reconstruction and simulation study. University of Minnesota, Minnesota
- Kirk RW, Bolstad PV, Manson SM (2012) Spatio-temporal trend analysis of long-term development patterns (1900–2030) in a Southern Appalachian County. *Landsc Urban Plan* 104(1):47–58
- Knoepp J, Clinton B (2009) Riparian zones in southern Appalachian headwater catchments: carbon and nitrogen responses to forest cutting. *For Ecol Manag* 258(10):2282–2293
- Laseter SH, Ford CR, Vose JM, Swift LW (2012) Long-term temperature and precipitation trends at the Coweeta Hydrologic Laboratory, Otto, North Carolina, USA. *Hydrol Res* 43(6):890
- Law N, Band L, Grove M (2004) Nitrogen input from residential lawn care practices in suburban watersheds in Baltimore County, MD. *J Environ Plan Manag* 47(5):737–755
- Le Mer J, Roger P (2001) Production, oxidation, emission and consumption of methane by soils: a review. *Eur J Soil Biol* 37(1):25–50

- Matson PA, Naylor R, Ortiz-Monasterio I (1998) Integration of environmental, agronomic, and economic aspects of fertilizer management. *Science* 280(5360):112–115
- NOAA (1950–2011) National Oceanic and Atmospheric Administration, National Climatic Data Center. Franklin, North Carolina climate station.
- NRCS Soil Survey Staff (2014) Natural Resources Conservation Service, United States Department of Agriculture, online at <http://websoilsurvey.nrcs.usda.gov/>. Accessed 2 Mar 2014
- Pinay G, Decamps H, Arles C, Lacassin-Seres M (1989) Topographic influence on carbon and nitrogen dynamics in riverine woods. *Arch Hydrobiol* 114(3):401–414
- Powelson D, Goulding K, Willison T, Webster C, Hütsch B (1997) The effect of agriculture on methane oxidation in soil. *Nutr Cycl Agroecosyst* 49(1–3):59–70
- Raich JW, Tufekciogul A (2000) Vegetation and soil respiration: correlations and controls. *Biogeochemistry* 48(1):71–90
- Robertson G, Coleman D, Bledsoe C, Sollins P (1999) Standard soil methods for long-term ecological research. Oxford University Press, USA
- Rosemond AD, Benstead JP, Bumpers PM, Gulis V, Kominoski JS, Manning DW, Suberkropp K, Wallace JB (2015) Experimental nutrient additions accelerate terrestrial carbon loss from stream ecosystems. *Science* 347(6226):1142–1145
- Schlesinger WH (2013) An estimate of the global sink for nitrous oxide in soils. *Glob Chang Biol* 19:2929–2931
- Schlesinger WH, Bernhardt ES (2013) *Biogeochemistry: an analysis of global change*. Academic Press, Cambridge
- Sexstone AJ, Parkin TB, Tiedje JM (1985) Temporal response of soil denitrification rates to rainfall and irrigation. *Soil Sci Soc Am J* 49(1):99–103
- Smith K, Thomson P, Clayton H, McTaggart I, Conen F (1998) Effects of temperature, water content and nitrogen fertilisation on emissions of nitrous oxide by soils. *Atmos Environ* 32(19):3301–3309
- Soussana JF, Loiseau P, Vuichard N, Ceschia E, Balesdent J, Chevallier T, Arrouays D (2004) Carbon cycling and sequestration opportunities in temperate grasslands. *Soil Use Manag* 20(2):219–230
- Studler P, Bowden R, Melillo J, Aber J (1989) Influence of nitrogen fertilization on methane uptake in temperate forest soils. *Nature* 341:314–316
- Swift JLW, Cunningham GB (1988) Climatology and hydrology. In: Swank WT, Crossley DA Jr (eds) *Forest hydrology and ecology at Coweeta*, vol 66. Springer, New York, p 469
- Tian H, Chen G, Lu C, Xu X, Hayes DJ, Ren W, Pan S, Huntzinger DN, Wofsy SC (2014) North American terrestrial CO<sub>2</sub> uptake largely offset by CH<sub>4</sub> and N<sub>2</sub>O emissions: toward a full accounting of the greenhouse gas budget. *Clim Chang* 129(3–4):413–426
- Tiedje J, Sexstone A, Parkin T, Revsbech N (1984) Anaerobic processes in soil. *Plant Soil* 76(1–3):197–212
- Turner MG, Pearson SM, Bolstad P, Wear DN (2003) Effects of land-cover change on spatial pattern of forest communities in the Southern Appalachian Mountains (USA). *Landsc Ecol* 18(5):449–464
- USEPA (1983a) Methods for chemical analysis of water and waste. Determination of nitrite/nitrate by automated cadmium reduction. In: Method 353.2, Environmental Monitoring and Support Lab., Office of Research and Development, USEPA, Cincinnati, OH
- USEPA (1983b) Methods for chemical analysis of water and waste. Determination of nitrogen as ammonia. In: Method 350.1, Environmental Monitoring and Support Lab., Office of Research and Development, USEPA, Cincinnati, OH
- Van den Heuvel RN, Bakker SE, Jetten MSM, Hefting MM (2011) Decreased N<sub>2</sub>O reduction by low soil pH causes high N<sub>2</sub>O emissions in a riparian ecosystem. *Geobiology* 9(3):294–300
- Verchot LV, Davidson EA, Cattânio JH, Ackerman IL (2000) Land-use change and biogeochemical controls of methane fluxes in soils of eastern Amazonia. *Ecosystems* 3(1):41–56
- Vose JM, Bolstad PV (2007) Biotic and abiotic factors regulating forest floor CO<sub>2</sub> flux across a range of forest age classes in the southern Appalachians. *Pedobiologia* 50(6):577–587
- Vought LBM, Dahl J, Pedersen CL, Lacoursière JO (1994) Nutrient retention in riparian ecotones. *Ambio* 23(6):342–348
- Walker J, Geron C, Vose J, Swank W (2002) Nitrogen trace gas emissions from a riparian ecosystem in southern Appalachia. *Chemosphere* 49(10):1389–1398
- Walker J, Vose J, Knoepp J, Geron C (2009) Recovery of nitrogen pools and processes in degraded riparian zones in the southern Appalachians. *J Environ Qual* 38(4):1391–1399
- Webster JR, Benfield EF, Cecala KK, Chamblee JF, Dehring CA, Gragson T, Cymerman JH, Jackson CR, Knoepp JD, Leigh DS (2012) Water quality and exurbanization in southern appalachian streams. *River Conserv Manag* pp 89–104
- Werner C, Kiese R, Butterbach-Bahl K (2007) Soil-atmosphere exchange of N<sub>2</sub>O, CH<sub>4</sub>, and CO<sub>2</sub> and controlling environmental factors for tropical rain forest sites in western Kenya. *J Geophys Res* 112(D3):D03308