

forest ecology

# The Influence of Dormancy Break Requirements on Germination and Viability Responses to Winter Submergence in Acorns of Three Bottomland Red Oak (Sect. *Lobatae*) Species

**Tracy S. Hawkins**

The objective of this study was to examine the relation of dormancy break requirements to winter submergence effects on germination and viability in cherrybark, willow, and Nuttall oak acorns. Acorns were submerged in a greentree reservoir and received 0, 21, 42, 63, or 84 days of winter submergence, followed by 8 weeks of incubation in four temperature regimes (15/6°, 20/10°, 25/15°, and 30/20° C). Winter submergence substituted for cold stratification requirements of cherrybark oak acorns, and dormancy break was achieved with 63 days of submergence. Although winter submergence exerted a positive effect on germination in Nuttall and willow oak acorns, dormancy break was not achieved in acorns of either species. Germination percentages were highest in the 30/20° C incubation temperature—cherrybark oak acorns (75–83 percent), followed by Nuttall oak (33–55 percent) and willow oak (14–52 percent). Among treatment combinations, all willow oak nongerminants were viable. Viability loss in cherrybark and Nuttall oak acorns was greatest in the 63- and 84-day treatments. The dynamics of acorn germination in response to winter hydrologic regime is influenced by submergence duration that may, or may not, satisfy dormancy break requirements in acorns of these red oak species.

**Keywords:** acorn germination, acorn submergence, bottomland forests, *Quercus*, red oak acorns

Flooding regime is one of the most important environmental factors affecting plant species distribution in bottomland forests in the southeastern United States (Hodges 1997, Gardiner 2001). Red oaks (*Quercus* spp.) of section *Lobatae* are often a major component of these forests; therefore, much research has focused on tree morphological and physiological responses to flooding frequency and duration. In turn, these research efforts have identified species-specific adaptations that direct plant distribution throughout the forest and allow for persistence in this type of habitat (Pezeshki and Anderson 1996, Gravatt and Kirby 1998, Anderson and Pezeshki 1999, Gardiner and Krauss 2001, and others). Far fewer studies have examined the effect of hydrologic regime on bottomland oak acorn germination and viability loss, and intra- and interspecific comparisons have varied dependent upon research protocol. Prior research has shown that acorn germination may be influenced by submergence duration (Briscoe 1961, Larsen 1963, Guo et al. 2002), water temperature (Guo et al. 1998), and whether submergence is static or intermittent (Walls et al. 2005).

However, the fate of acorns that fail to germinate (i.e., remain viable or are nonviable) is severely underreported in the literature (but see Guo et al. 1998), and knowledge gaps regarding causative effects for interspecific differences in acorn germination success or failure continue to exist. Additionally, post-submergence acorn fate has not been investigated relative to seed dormancy.

Seed dormancy is defined as a failure of seeds to germinate, although environmental conditions (e.g., water, temperature, light availability) are favorable for germination (Baskin and Baskin 2004, 2014). In turn, this ecological adaptation delays seed germination until conditions are conducive for seedling establishment and survival. The underlying cause of seed dormancy may be anatomical, such as an impermeable barrier (e.g., seed-coat) that prevents water uptake (imbibition; Baskin et al. 2000) or an underdeveloped embryo that must grow and develop prior to seed germination (Baskin and Baskin 2005). Alternatively, if seeds possess a developed embryo and readily imbibe water, dormancy break occurs when environmental cues, such as a period of warm or cold stratification,

Manuscript received July 24, 2018; accepted April 17, 2019; published online June 5, 2019.

**Affiliation:** Tracy S. Hawkins ([tracyhawkins@fs.fed.us](mailto:tracyhawkins@fs.fed.us)), USDA Forest Service, Center for Bottomland Hardwoods Research, Box 9681, Mississippi State, MS 39762.

**Acknowledgments:** The author thanks Rory Thornton for help with data collection, the Mississippi Forestry Commission for permitting acorn collection at Winona Seed Orchard, and Dr. Charles Sabatia for contributions to data analyses and review of an earlier draft of this manuscript.



16 cm × 16 cm × 2 cm clear plastic dishes and on moistened river sand purchased from a local hardware store. Four replicates of 25 acorns of each oak species were placed in each of the incubation temperature regimes on November 10, 2010.

Sixteen bags of acorns for each species were removed from the GTR and returned to the laboratory at 21, 42, 63, and 84 days of winter submergence. Acorns were removed from the bags, washed gently, and placed on moist sand in clear plastic dishes. Four replicates of 25 acorns for each of the species were placed in each of the four incubators.

Acorn germination was recorded at 1-week intervals for 8 weeks. Emergence of the radicle ( $\leq 2$  mm) was the criterion for germination. At 8 weeks of incubation, all acorns that did not germinate were checked for viability. Firm, white embryos were considered viable. Acorns with soft, brown to gray embryos, or those partially or fully decayed were considered nonviable.

### Statistical Analysis


Means and standard errors were calculated for germination and nonviability percentages. A binomial distribution generalized linear model with partitioned analysis of least significant means was used to test for fixed effects and interaction of winter submergence duration and incubation temperature on acorn germination and viability. An overdispersion parameter was included for fitting the model. A post-hoc Tukey's HSD ( $P = .05$ ) procedure was used to adjust  $P$ -values for simple effect comparisons of the levels of temperature at a given level of winter submergence treatment and vice versa. The SAS procedure GLIMMIX was used to perform all analyses (SAS Institute Inc. 2007).

### Results

The water temperature in the GTR at the depth of submerged acorns (45 cm) was 13.0° C at the beginning of the study (November 10, 2010). Thereafter, water temperatures steadily declined, showed minimal fluctuation over the course of winter submergence treatments, and remained below 8° C (Figure 1). Acorns did not germinate during submergence.

The interaction of winter submergence duration and incubation temperature had a significant effect on germination percentages of cherrybark oak acorns ( $df = 12$ ,  $F = 10.00$ ,  $P < .0001$ ). Germination percentages increased with increasing winter submergence duration when incubated at 15/6° ( $df = 4$ ,  $F = 137.18$ ,  $P < .0001$ ) and 20/10° C ( $df = 4$ ,  $F = 102.81$ ,  $P < .0001$ ) through the 63-day treatment, and at 25/15° C ( $df = 4$ ,  $F = 29.95$ ,  $P < .0001$ ) through the 42-day treatment (Table 1). In the 63- and 84-day treatments, mean total germination percentages across all incubation temperatures were  $\geq 75 \pm 5$  percent (Table 1), and in these treatments, maximum germination percentages were reached within 4 weeks at the 25/15° and 30/20° C incubation temperatures (data not shown).

The interaction of winter submergence duration and incubation temperature affected nonviability percentages of cherrybark oak acorns ( $df = 12$ ,  $F = 5.06$ ,  $P < .0001$ ). In the control and 21-day treatment, all cherrybark oak acorns that did not germinate were viable (Table 1). However, in the 42-day treatment,  $40 \pm 4$  percent and  $21 \pm 6$  percent of acorns were nonviable in the 15/6° and 20/10° C temperatures, respectively. All nongerminants in the 63- and 84-day treatments were nonviable in all temperature regimes (Table 1).



**Figure 1.** Water temperatures (° C; depth = 45 cm) received by cherrybark (*Quercus pagoda*), Nuttall (*Q. texana*), and willow oak (*Q. phellos*) acorns during a gradient of winter submergence treatments in a greentree reservoir at Sam D. Hamilton Noxubee Wildlife Refuge, Noxubee County, Mississippi. Arrows indicate time (date) of acorn removal from the GTR, and numbers represent the total number of days of the winter submergence treatment.

Mean total germination in willow oak acorns was affected independently by winter submergence duration ( $df = 4$ ,  $F = 10.22$ ,  $P < .0001$ ) and incubation temperature ( $df = 3$ ,  $F = 10.91$ ,  $P < .0001$ ). At 30/20° C, total germination was  $\geq 46$  percent among winter submergence treatments, with the exception of the 42-day treatment, where mean total germination was  $14 \pm 4$  percent (Table 2). Mean total germination percentages were greatest in the 84-day treatment for acorns incubated at 15/6° ( $25 \pm 7$  percent), 20/10° ( $23 \pm 3$  percent), and 25/15° C ( $43 \pm 3$  percent) incubation temperatures. All ungerminated acorns in all treatments were viable (Table 2).

Nuttall oak acorn germination was affected by the interaction of winter submergence duration and incubation temperature ( $df = 12$ ,  $F = 3.58$ ,  $P = .0005$ ). Mean total germination percentages for acorns in the 30/20° C temperature regime increased with 42 days of winter submergence and were consistently highest among incubation temperatures in all submergence treatments (Table 3). At the 20/10° and 25/15° C incubation temperatures, germination percentages increased with 63 days ( $df = 4$ ,  $F = 17.47.96$ ,  $P < .0001$ ) and 84 days ( $df = 4$ ,  $F = 70.07$ ,  $P < .0001$ ) of winter submergence, respectively. Some acorns at the 15/6° C incubation temperature germinated in the 63- and 84-day treatments; however, percentages were  $\leq 10 \pm 2$  percent (Table 3).

The interaction of winter submergence duration and incubation temperature ( $df = 12$ ,  $F = 3.04$ ,  $P = .0021$ ) affected nonviability percentages in Nuttall oak acorns. All nongerminants in the control were viable. However, in the 21-day treatment,  $51 \pm 4$  percent of all acorns in the 30/20° C incubation temperature were nonviable (Table 3). Among incubation temperatures in the 42-day treatment, nonviability percentages ranged from  $3 \pm 2$  percent (25/15° C) to  $39 \pm 4$  percent (30/20° C). All ungerminated acorns in all incubation temperatures in the 63- and 84-day treatments were nonviable (Table 3).

### Discussion

Seed dormancy break is a dynamic process, and in the vast majority of species with seeds that possess physiological dormancy

**Table 1. Mean ( $\pm$ SE) total germination, viability, and nonviability percentages for cherrybark oak (*Quercus pagoda*) acorns receiving 0, 21, 42, 63, and 84 days of winter submergence in a greentree reservoir, followed by 8 weeks of incubation at 15/6°, 20/10°, 25/15°, and 30/20° C.**

Winter submergence (days)		Incubation alternating temperature regime (° C)			
		15/6	20/10	25/15	30/20
Germination (percent)	0	5 $\pm$ 3 <sup>Aa</sup>	23 $\pm$ 4 <sup>Ba</sup>	62 $\pm$ 7 <sup>Ca</sup>	80 $\pm$ 4 <sup>Da</sup>
	21	5 $\pm$ 2 <sup>Ab</sup>	49 $\pm$ 4 <sup>Bb</sup>	64 $\pm$ 4 <sup>Ca</sup>	83 $\pm$ 3 <sup>Da</sup>
	42	42 $\pm$ 3 <sup>Ac</sup>	68 $\pm$ 5 <sup>Bc</sup>	85 $\pm$ 4 <sup>Bb</sup>	83 $\pm$ 5 <sup>Ba</sup>
	63	82 $\pm$ 1 <sup>Ad</sup>	88 $\pm$ 5 <sup>Ad</sup>	82 $\pm$ 3 <sup>Ab</sup>	81 $\pm$ 4 <sup>Aa</sup>
	84	84 $\pm$ 6 <sup>Ad</sup>	85 $\pm$ 2 <sup>Ad</sup>	87 $\pm$ 3 <sup>Ab</sup>	75 $\pm$ 5 <sup>Aa</sup>
Viable (percent)	0	95 $\pm$ 3 <sup>Aa</sup>	77 $\pm$ 4 <sup>Ba</sup>	38 $\pm$ 7 <sup>Ca</sup>	20 $\pm$ 4 <sup>Da</sup>
	21	75 $\pm$ 2 <sup>Ab</sup>	51 $\pm$ 4 <sup>Bb</sup>	36 $\pm$ 4 <sup>Ca</sup>	17 $\pm$ 2 <sup>Da</sup>
	42	18 $\pm$ 3 <sup>Ac</sup>	11 $\pm$ 3 <sup>Ac</sup>	15 $\pm$ 4 <sup>Ab</sup>	17 $\pm$ 5 <sup>Aa</sup>
	63	0 <sup>Ad</sup>	0 <sup>Ad</sup>	0 <sup>Ac</sup>	0 <sup>Ab</sup>
	84	3 $\pm$ 2 <sup>Ad</sup>	0 <sup>Ad</sup>	1 $\pm$ 1 <sup>Ac</sup>	1 $\pm$ 1 <sup>Ab</sup>
Nonviable (percent)	0	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>
	21	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>
	42	40 $\pm$ 4 <sup>Ab</sup>	21 $\pm$ 6 <sup>Bb</sup>	0 <sup>Ca</sup>	0 <sup>Ca</sup>
	63	18 $\pm$ 1 <sup>Ac</sup>	12 $\pm$ 5 <sup>Ac</sup>	18 $\pm$ 3 <sup>Ab</sup>	19 $\pm$ 4 <sup>Ab</sup>
	84	13 $\pm$ 6 <sup>Ac</sup>	15 $\pm$ 2 <sup>Ac</sup>	12 $\pm$ 3 <sup>Ab</sup>	24 $\pm$ 5 <sup>Bb</sup>

Note: Means with dissimilar uppercase letters are significantly different within a submergence duration, and means with dissimilar lowercase letters are significantly different within a given incubation temperature and acorn fate (Tukey's HSD,  $P = .05$ ).

**Table 2. Mean ( $\pm$ SE) total germination and viability percentages for willow oak (*Quercus phellos*) acorns receiving 0, 21, 42, 63, and 84 days of winter submergence in a greentree reservoir, followed by 8 weeks of incubation at 15/6°, 20/10°, 25/15°, and 30/20° C.**

Winter submergence (days)		Incubation alternating temperature regime (° C)			
		15/6	20/10	25/15	30/20
Germination (percent)	0	0 <sup>Aa</sup>	0 <sup>Aa</sup>	6 $\pm$ 3 <sup>Aa</sup>	50 $\pm$ 8 <sup>Ba</sup>
	21	0 <sup>Aa</sup>	4 $\pm$ 2 <sup>Aa</sup>	20 $\pm$ 5 <sup>Bb</sup>	46 $\pm$ 12 <sup>Ca</sup>
	42	3 $\pm$ 2 <sup>Aa</sup>	6 $\pm$ 3 <sup>Aa</sup>	11 $\pm$ 6 <sup>Aa</sup>	14 $\pm$ 4 <sup>Ab</sup>
	63	0 <sup>Aa</sup>	6 $\pm$ 1 <sup>Aa</sup>	22 $\pm$ 9 <sup>Bb</sup>	52 $\pm$ 5 <sup>Ca</sup>
	84	25 $\pm$ 7 <sup>Ab</sup>	23 $\pm$ 3 <sup>Ab</sup>	43 $\pm$ 3 <sup>Bc</sup>	52 $\pm$ 2 <sup>Ba</sup>
Viable (percent)	0	100 $\pm$ 0 <sup>Aa</sup>	100 $\pm$ 0 <sup>Aa</sup>	94 $\pm$ 3 <sup>Aa</sup>	50 $\pm$ 8 <sup>Ba</sup>
	21	100 $\pm$ 0 <sup>Aa</sup>	96 $\pm$ 2 <sup>Aa</sup>	80 $\pm$ 5 <sup>Bb</sup>	54 $\pm$ 12 <sup>Ca</sup>
	42	97 $\pm$ 2 <sup>Aa</sup>	94 $\pm$ 3 <sup>Aa</sup>	89 $\pm$ 6 <sup>Aa</sup>	86 $\pm$ 4 <sup>Ab</sup>
	63	100 $\pm$ 0 <sup>Aa</sup>	94 $\pm$ 1 <sup>Aa</sup>	78 $\pm$ 9 <sup>Bb</sup>	48 $\pm$ 5 <sup>Ca</sup>
	84	75 $\pm$ 7 <sup>Ab</sup>	77 $\pm$ 3 <sup>Ab</sup>	57 $\pm$ 3 <sup>Bc</sup>	48 $\pm$ 2 <sup>Ba</sup>

Note: Means with dissimilar uppercase letters are significantly different within a submergence duration, and means with dissimilar lowercase letters are significantly different within a given incubation temperature and acorn fate (Tukey's HSD,  $P = .05$ ).

**Table 3. Mean ( $\pm$ SE) total germination, viability, and nonviability percentages for Nuttall oak (*Quercus texana*) acorns receiving 0, 21, 42, 63, and 84 days of winter submergence in a greentree reservoir, followed by 8 weeks of incubation at 15/6°, 20/10°, 25/15°, and 30/20° C.**

Winter submergence (days)		Incubation alternating temperature regime (° C)			
		15/6	20/10	25/15	30/20
Germination (percent)	0	0 <sup>Aa</sup>	0 <sup>Aa</sup>	4 $\pm$ 2 <sup>Aa</sup>	33 $\pm$ 4 <sup>Ba</sup>
	21	1 $\pm$ 1 <sup>Aa</sup>	4 $\pm$ 2 <sup>Aa</sup>	14 $\pm$ 4 <sup>Ba</sup>	39 $\pm$ 6 <sup>Ca</sup>
	42	0 <sup>Aa</sup>	6 $\pm$ 3 <sup>Ba</sup>	9 $\pm$ 2 <sup>Ba</sup>	55 $\pm$ 2 <sup>Ca</sup>
	63	7 $\pm$ 2 <sup>Aa</sup>	15 $\pm$ 4 <sup>Ab</sup>	15 $\pm$ 7 <sup>Aa</sup>	43 $\pm$ 5 <sup>Ba</sup>
	84	10 $\pm$ 2 <sup>Ab</sup>	21 $\pm$ 1 <sup>Bb</sup>	46 $\pm$ 5 <sup>Cb</sup>	51 $\pm$ 8 <sup>Ca</sup>
Viable (percent)	0	100 $\pm$ 0 <sup>Aa</sup>	100 $\pm$ 0 <sup>Aa</sup>	96 $\pm$ 2 <sup>Aa</sup>	66 $\pm$ 4 <sup>Ba</sup>
	21	99 $\pm$ 1 <sup>Aa</sup>	96 $\pm$ 2 <sup>Aa</sup>	86 $\pm$ 4 <sup>Aa</sup>	9 $\pm$ 3 <sup>Bb</sup>
	42	93 $\pm$ 3 <sup>Aa</sup>	74 $\pm$ 7 <sup>Bb</sup>	82 $\pm$ 1 <sup>Bb</sup>	6 $\pm$ 2 <sup>Cb</sup>
	63	0 <sup>Ab</sup>	0 <sup>Ac</sup>	0 <sup>Ac</sup>	0 <sup>Ab</sup>
	84	0 <sup>Ab</sup>	0 <sup>Ac</sup>	0 <sup>Ac</sup>	0 <sup>Ab</sup>
Nonviable (percent)	0	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>
	21	0 <sup>Aa</sup>	0 <sup>Aa</sup>	0 <sup>Aa</sup>	51 $\pm$ 4 <sup>Bb</sup>
	42	6 $\pm$ 3 <sup>Aa</sup>	14 $\pm$ 7 <sup>Ab</sup>	3 $\pm$ 2 <sup>Aa</sup>	39 $\pm$ 4 <sup>Bb</sup>
	63	93 $\pm$ 2 <sup>Ab</sup>	85 $\pm$ 4 <sup>Ac</sup>	85 $\pm$ 7 <sup>Ab</sup>	57 $\pm$ 5 <sup>Bb</sup>
	84	90 $\pm$ 2 <sup>Ab</sup>	79 $\pm$ 1 <sup>Bc</sup>	54 $\pm$ 5 <sup>Cc</sup>	49 $\pm$ 8 <sup>Cb</sup>

Note: Means with dissimilar uppercase letters are significantly different within a submergence duration, and means with dissimilar lowercase letters are significantly different within a given incubation temperature and acorn fate (Tukey's HSD,  $P = .05$ ).



(PD), dormancy break is characterized by germination occurring in an expanding range of temperatures following increasing stratification time. Additionally, increasing germination percentages within these respective temperatures are observed (Baskin and Baskin 2004, 2014). In the laboratory, these parameters are most often evaluated at 4 weeks of incubation; however, in this study, incubation time was extended to 8 weeks to allow more time to investigate the fate of nongerminants.

Among the three red oak species, dormancy break was achieved only in cherrybark oak acorns. Following 63 days of winter submergence, germination percentages were  $\geq 80$  percent in all incubation temperatures and did not increase with 84 days of submergence. These germination percentages are consistent with those reported for cherrybark oak acorns following the same period of cold stratification on moist medium (Hawkins 2018). In other words, winter submergence of at least 63 days may substitute for cold stratification in breaking dormancy in cherrybark oak acorns. Guo et al. (2002) reported 81–97 percent germination (at approximately 20° C post-submergence) for cherrybark oak acorns following only 0–30 days of winter submergence. However, acorns received 45 days of cold stratification prior to the treatments (Guo et al. 2002). Prior cold stratification for the control (0 days) and the additive effect of cold stratification plus winter submergence would be expected to yield higher germination percentages relative to treatments of comparable submergence duration in this study. Additionally, in the Guo et al. (2002) study, acorns were collected from the ground during late autumn (November) and thus had received some stratification prior to collection.

Viability loss in cherrybark oak acorns began with 42 days of winter submergence and only at the two lowest incubation temperatures. This indicates that viability loss was temperature-dependent following winter submergence and occurred prior to dormancy break in the acorn cohort. On the other hand, the percentage of nonviable acorns in the 63- and 84-day treatments was relatively low (12–24 percent) and existed in all incubation temperatures. These data suggest that acorn viability loss with longer flood durations is not dependent upon post-drop down temperatures. However, it is not possible to determine whether acorn loss occurred during or after winter submergence.

Larsen (1963) and Guo et al. (2002) reported 45–55 percent total germination for willow oak acorns when incubated at a single temperature (20–27° C) following varying submergence durations at water temperatures ranging from 8 to 22° C. Similarly, germination percentages for willow oak acorns ( $\leq 52$  percent) in this study were low, particularly relative to those of cherrybark oak. A significant increase in total germination percentages among the three lower incubation temperatures occurred only with 84 days of winter submergence, which suggests a further increase in germination percentages was possible with increased winter submergence. This pattern of germination response for acorns cold stratified on moist medium would indicate a failure of stratification time in satisfying dormancy break requirements. Therefore, either dormancy break in willow oak acorns requires a period of cold stratification (or winter submergence) longer than that of cherrybark oak, or presuming acorns require less than 84 days of cold stratification for dormancy break, winter submergence does not substitute for cold stratification.

Of note is the significant decrease in willow oak acorn total germination in the 25/15° and 30/20° C temperature regimes following the 42-day treatment. A reduction in total germination at these incubation temperatures has not been observed in acorns of this species

following a gradient of cold stratification treatments of the same duration (Hawkins, unpublished results); nor was it observed in the cherrybark or Nuttall oak acorns in this study. Therefore, this occurrence presents two possibilities. The depression in germination percentages is an artifact of this study, or this is a species-specific response to this length of flood duration, water temperature, incubation temperature, or an interaction of these variables. Effective cold stratification temperatures of seeds on a moist substrate are within the range of 0–10° C (Stokes 1965), and water temperatures during the 42-day treatment were  $\leq 10$ ° C for the latter 32 days of the treatment. Therefore, it is unlikely that the decrease in germination percentages for this treatment is solely the result of water temperature.

Germination percentages ( $\leq 62$  percent) in Nuttall oak acorns were similar to that of willow oak and thus were low relative to those of cherrybark oak acorns. Although increasing winter submergence durations resulted in increasing germination percentages across a wider range of temperatures, nongerminants were nonviable in the 63- and 84-day winter submergence treatments without respect to incubation temperature. These results are in contrast to those of Briscoe (1961), who found that submergence at water temperatures of 3–4° C had no effect on Nuttall oak acorn germination. The discrepancy is most likely related to submergence time, which was terminated at 34 days in Briscoe's (1961) study.

Mature Nuttall oak trees are moderately flood-tolerant and may withstand up to 3 years of inundation (Gardiner 2001). This level of flood tolerance was not exhibited in acorns of this species. In winter flood durations of 21 and 42 days, the percentage of acorns found to be nonviable was dependent on incubation temperature, indicating acorn loss occurred post-submergence. However, with 63 and 84 days of winter submergence, all nongerminants (49–93 percent of total) were nonviable, regardless of incubation temperature. Within the methodology of this study, it is impossible to determine whether acorns in these two winter submergence treatments were rendered nonviable during or after submergence. Regardless of the time of occurrence, Nuttall oak acorns appear to be more susceptible to viability loss than those of cherrybark and willow oak when subject to winter submergence.

## Conclusions

The dynamics of red oak acorn germination in response to hydrologic regime is influenced by dormancy break requirements, which vary among species of *Lobatae*. In bottomland forests, and within the constraints of the variables in this research, if winter submergence is of sufficient length for dormancy break in cherrybark oak acorns, a high percentage of acorns will readily germinate post-drop down across a range of ambient temperatures. Concurrently, the percentage of germinants in a cohort of willow acorns may be less than that of cherrybark oak because of dormancy break requirements not being met, and immediate germination post-drop down would be strongly dependent on ambient temperature. However, retention of viability in willow oak nongerminants allows for further stratification post-drop down, and acorns may continue to germinate as the cohort approaches dormancy break. Similarly, germination in Nuttall oak acorns would be expected to be low relative to cherrybark oak acorns. However, the potential for post-drop down germination in Nuttall oak acorns may be offset because of viability loss during winter submergence durations greater than 42 days.

## Literature Cited

- ANDERSON, P.H., AND S.R. PEZESHKI. 1999. The effects of intermittent flooding on seedlings of three forest species. *Photosynthetica* 37:543–552.
- BASKIN, C.C., AND J.M. BASKIN. 2005. Underdeveloped embryos in dwarf seeds and implications for assignment to dormancy class. *Seed Sci. Res.* 15:357–360.
- BASKIN, C.C., AND J.M. BASKIN. 2014. *Seeds: Ecology, biogeography and evolution of dormancy and germination*. 2nd ed. Elsevier/Academic Press, San Diego, CA. 1586 p.
- BASKIN, J.M., AND C.C. BASKIN. 2004. A classification system for seed dormancy. *Seed Sci. Res.* 14:1–16.
- BASKIN, J.M., C.C. BASKIN, AND X. LI. 2000. Taxonomy, anatomy and evolution of physical dormancy in seeds. *Plant Spec. Biol.* 15:139–152.
- BONNER, F.T., AND J.A. VOZZO. 1987. *Seed biology and technology of Quercus*. USDA Forest Service Gen. Tech. Rep. SO-66. 21 p.
- BRENT JR., F.V. 1986. *Soil Survey of Noxubee County, Mississippi*. USDA, Soil Conservation Service, Washington, DC. 198 p.
- BRISCOE, C.B. 1961. Germination of cherrybark and Nuttal(l) oak acorns following flooding. *Ecology* 42:430–431.
- GARDINER, E.S. 2001. Ecology of bottomland oaks in the southeastern United States. *Int. Oaks* 12:48–55.
- GARDINER, E.S., AND K.W. KRAUSS. 2001. Photosynthetic light response of flooded cherrybark oak (*Quercus pagoda*) seedlings grown in two light regimes. *Tree Physiol.* 21:1103–1111.
- GRAVATT, D.A., AND C.J. KIRBY. 1998. Patterns of photosynthesis and starch allocation in seedlings of four bottomland hardwood tree species subjected to flooding. *Tree Physiol.* 18:411–417.
- GUO, Y., M.G. SHELTON, AND B.R. LOCKHART. 1998. Effects of flood duration and season on germination of black, cherrybark, northern red, and water oak acorns. *New Forests* 15:69–76.
- GUO, Y., M.G. SHELTON, AND E. HEITZMAN. 2002. Effects of flood duration and depth on germination of cherrybark, post, southern red, white, and willow oak acorns. P. 622 in *Proceedings of the eleventh biennial southern silviculture research conference*, K.W. OUTCALT, (ed.). USDA Forest Service Gen. Tech. Rep. SRS-48.
- HAWKINS, T.S. 2018. Regulating acorn germination and seedling emergence in *Quercus pagoda* (Raf.) as it relates to natural and artificial regeneration. *New Forests* doi: 10.1007/s11056-018-9667-z.
- HAWKINS, T.S., J.L. WALCK, AND S.N. HIDAYATI. 2011. Seed ecology of *Lindera melissifolia* (Lauraceae) as it relates to rarity of the species. *J. Torrey Bot. Soc.* 138:298–307.
- HODGES, J.D. 1997. Development and ecology of bottomland hardwood sites. *Forest Ecol. Manag.* 90:117–125.
- INSAUSTI, P., A. SORIANO, AND R.A. SANCHEZ. 1995. Effects of flood-influenced factors on seed germination of *Ambrosia tenuifolia*. *Oecologia* 103:127–132.
- LARSEN, H.S. 1963. Effects of soaking in water on acorn germination of four southern oaks. *For. Sci.* 9:236–241.
- MORAVKOVÁ, L., P. ZÁKRAVSKÝ, AND Z. HOUDOVÁ. 2002. Germination response to temperature and flooding of four Central European species of *Bolboschoenus*. *Preslia. Praha* 74:333–343.
- OBROUCHEVA, N.V. 2010. Distinct regulatory patterns of seed dormancy release and germination commencement. *Seed Sci. Technol.* 38:265–279.
- PEZESHKI, S.R., AND J.H. ANDERSON. 1996. Responses of three bottomland species with different flood tolerance capabilities to various flooding regimes. *Weil. Ecol. Manag.* 4:245–256.
- SAS INSTITUTE, INC. 2007. *The SAS system for Windows, release V9.4*. SAS Institute, Cary, NC.
- STOKES, P. 1965. Temperature and seed dormancy. P. 746–803 in *Encyclopedia of plant physiology*, Vol. 15/2, W. RUHLAND (ed.). Springer-Verlag, Berlin and New York.
- TANNER, J.T. 1986. Distribution of tree species in Louisiana bottomland forests. *Castanea* 51:168–174.
- WALLS, R.L., D.H. WARDROP, AND R.P. BROOKS. 2005. The impact of experimental sedimentation and flooding on the growth and germination of floodplain trees. *Plant Ecol.* 176:203–213.