

Research Paper

Cite this article: Hawkins TS (2020). Dormancy break and germination requirements in acorns of two bottomland *Quercus* species (Sect. *Lobatae*) of the eastern United States with references to ecology and phylogeny. *Seed Science Research* **30**, 199–205. <https://doi.org/10.1017/S0960258520000264>

Received: 25 October 2019
Revised: 22 May 2020
Accepted: 22 June 2020
First published online: 16 July 2020

Key words:

acorn dormancy; acorn germination; *Lobatae*; oak regeneration; *Quercus nigra*; *Quercus phellos*

Correspondence:

Tracy S. Hawkins,
E-mail: tracy.hawkins@usda.gov

Dormancy break and germination requirements in acorns of two bottomland *Quercus* species (Sect. *Lobatae*) of the eastern United States with references to ecology and phylogeny

Tracy S. Hawkins 

USDA Forest Service, Center for Bottomland Hardwoods Research, Box 9681, Mississippi State, MS 39762, USA

Abstract

Quercus species are ecologically and economically important components of deciduous forests of the eastern United States. However, knowledge pertinent to a thorough understanding of acorn germination dynamics for these species is lacking. The objectives of this research were to determine dormancy break and germination requirements for acorns of two eastern United States bottomland species, *Quercus nigra* and *Quercus phellos* (Section *Lobatae*), and to present results within ecological and phylogenetic contexts. Three replicates of 50 acorns of each species received 0 (control), 6, 12 or 18 weeks of cold stratification, followed by incubation in alternating temperature regimes of 15/6, 20/10, 25/15 and 30/20°C. Eighteen weeks of cold stratification were not sufficient for dormancy break in *Q. nigra* acorns. Cumulative germination percentages at 4 weeks of incubation were $\geq 77\%$, but only in incubation temperatures of 25/15 and 30/20°C. Dormancy break in *Q. phellos* acorns was achieved with 18 weeks of cold stratification, and cumulative germination percentages were $\geq 87\%$ at 4 weeks of incubation in all test temperature regimes. Gibberellic acid solutions were not an effective substitute for cold stratification in either species. Phylogenetically, *Q. nigra* and *Q. phellos* are closely related species and, ecologically, both grow in the same habitat. Acorns of both species possess deep physiological dormancy (PD), but dormancy break and germination requirements differ in acorns of these two *Quercus* species.

Introduction

The oaks (*Quercus* spp.) are described as the most ecologically and economically important tree species in North America (Bonner and Karrfalt, 2008; Cavender-Bares, 2016). Approximately 58 *Quercus* species are native to the United States and 50 of these grow in the eastern United States (Stein et al., 2003). The *Quercus* species of the eastern United States are represented by two groups commonly referred to as the red oaks and the white oaks. Within the *Quercus* subgenus, the red oaks are placed in section *Lobatae*, and the white oaks are in section *Quercus*. Phylogenetically, these sections represent two major American oak clades that are of northern temperate origin and diverged ca. mid-Eocene (Hipp et al., 2018). Thereafter, sympatric parallel diversification of *Lobatae* and *Quercus* occurred as species moved southward during the Oligocene to latter Miocene, and ecological diversification proceeded within clades (Hipp et al., 2018). Indeed, in the eastern United States, representatives of both sections are common canopy components in mesophytic to xeric upland forests (Abrams, 1992), and in bottomland forests that often are subject to periodic inundation (Hodges, 1994; King and Keeland, 1999; Gardiner, 2001).

To date, we have only a broad understanding of acorn germination dynamics for a relatively few eastern United States *Quercus* species, and the majority of past studies have focused on those of section *Lobatae*. For example, Bonner and Vozzo (1987) reported that the length of cold stratification required to ‘enhance acorn germination’ varied among nine *Quercus* species of *Lobatae*. Peterson (1983) found that increased lengths of cold stratification reduced the time to maximum germination in *Quercus nigra* L. acorns, and Hopper et al. (1985) quantified this same trend in acorns of *Quercus rubra* L. We can infer from these studies that physiological dormancy (PD) is present in acorns of many species of section *Lobatae*; however, it is unclear whether different levels of PD are present.

Seeds may possess non-deep, intermediate or deep PD (Baskin and Baskin 2004, 2014). The level of PD is identified by seed dormancy break and germination response to the length of stratification (warm and/or cold) and to treatment with gibberellic acid (GA₃). In other words, seeds with deep PD require longer periods of stratification (>12 weeks) for dormancy break than those with non-deep PD (≤ 12 weeks). Those with intermediate PD require 8–12 weeks of stratification. Additionally, GA₃ promotes germination in seeds with non-deep PD,

may or may not promote germination in those with intermediate PD and would not be effective in seeds with deep PD (Baskin and Baskin, 2014).

Quercus pagoda grows in bottomland hardwood forests of the eastern United States and often in association with other *Quercus* species such as *Q. laurifolia* Michx., *Q. nigra* and *Quercus phellos* L. (Gardiner, 2001). Acorns of *Q. pagoda* Raf. (Section *Lobatae*) require at least 12 weeks of cold stratification for dormancy break and possess non-deep PD (Hawkins, 2019a). However, past research (e.g. Peterson, 1983; Hopper et al., 1985; Bonner and Vozzo, 1987) suggests that different levels of PD may be present among species of section *Lobatae*, and thus, acorn dormancy break requirements of *Q. pagoda* may differ from that of other sympatric *Quercus* species. The primary objective of this research was to identify dormancy break and germination requirements in acorns of *Q. nigra* and *Q. phellos*. This component of the research will determine the level of PD in acorns of these two species and allow for comparison of dormancy break and germination requirements among three bottomland *Quercus* species. This will add to our limited knowledge of the ecology of acorn germination dynamics within a bottomland forest habitat and begin to elucidate phylogenetic associations of dormancy types or level of PD among species of section *Lobatae*.

The second objective of this study was to determine if varying acorn treatments, such as length of cold stratification and/or post-stratification temperature in a controlled environment, are required for interspecific germination synchrony in acorns of *Q. nigra* and *Q. phellos*. In the last 70 years, a significant decline in *Quercus* spp. abundance in the eastern United States forests has resulted from a culmination of timbering practices, e.g. high grading (McGee, 1972), fungal pathogens (Bruhn et al., 2008; Kelley et al., 2009), insect damage (Stephen et al., 2001; Starkey et al., 2004; Haavik et al., 2012) and changes in fire disturbance patterns (Guyette et al., 2002; Arthur et al., 2012). In turn, much attention has focused on mitigating *Quercus* spp. decline by employing artificial regeneration practices in *Quercus* depleted forests and on former agricultural sites (Stanturf et al., 2000; Dey et al., 2008). The success of these regeneration efforts may be enhanced by synchronizing acorn germination to produce even-aged *Quercus* seedlings (Dey et al., 2008).

Materials and methods

The species

Q. nigra grows along the Coastal Plain from southern New Jersey and Delaware; south to Florida; west to eastern Texas; north to Missouri and east to Virginia (Vozzo, 1990). This species grows on a variety of sites ranging from poorly drained bottomlands to ridges and high flats (Vozzo, 1990; Gardiner, 2001). Mature trees are moderately flood-tolerant (Gardiner, 2001), and acorns have been shown to retain viability up to 30 days during winter or spring submergence (Guo et al., 1998). However, the effects of submergence on *Q. nigra* acorn viability have not been tested beyond 30 days.

Q. phellos ranges from New York; west to Missouri; south to Texas, east to Florida and north to Delaware (Schlaegel, 1990). The species grows on a variety of soil types in bottomland forests, but most often is found on high flats and loamy ridges of recent alluvium (Schlaegel, 1990; Gardiner, 2001). Mature *Q. phellos* trees are moderately flood-tolerant (Gardiner, 2001), and acorns of this species have been reported to retain viability for up to 84 days of winter submergence (Hawkins, 2019b).

Sampling and conditions

In November 2012, mature acorns were harvested directly from randomly selected *Q. phellos* trees growing in Starkville, Mississippi (33.4504°N, 88.8184°W) and from *Q. nigra* trees growing in Amory, Mississippi (33.9843°N, 88.4881°W). Acorns were immediately taken to the laboratory and remained in covered 19-L buckets at ambient temperature (~24°C) for 16 h before receiving experimental treatments.

Five incubators were set at 12 h/12 h diurnal alternating temperature regimes that approximate seasonal temperatures within the range of the two *Quercus* species. One of five incubators was set at 5/1°C (winter) and used for cold stratification. Temperature regimes for the remaining four incubators were 15/6°C (early spring/late fall), 20/10°C (late spring/early fall), 25/15°C (early/late summer) and 30/20°C (summer). All incubators were set to provide 12 h of light (50 $\mu\text{mol m}^{-2} \text{s}^{-1}$) during the high-temperature period and 12 h of uninterrupted darkness during the low-temperature period.

Effects of cold stratification on acorn dormancy break

Acorns were placed on sand moistened with distilled water in 16 cm \times 16 cm \times 5 cm clear plastic dishes. Plastic lids were placed on the dishes to retard moisture loss. Three replicates of 50 acorns per dish for each species were used in each treatment. Dishes were placed in a completely randomized design in each of the treatments. Acorns received 0 (control), 6, 12 or 18 weeks of cold stratification at 5/1°C. Acorns used as controls (0-week cold stratification) were placed directly in the test temperatures (15/6, 20/10, 25/15 and 30/20°C). Following each cold stratification treatment, acorns received an incubation time of 12 weeks in each of the test temperature regimes. Germination was recorded at 2-week intervals. Emergence of the radicle (at least 1 mm) was the criterion for germination. At the conclusion of the incubation period, acorns that did not germinate were assessed for viability. Acorns containing firm, white embryos were scored as viable, and those with soft, brown to grey embryos were scored as non-viable.

Effects of gibberellic acid (GA₃) on acorn dormancy break

To determine whether gibberellic acid (GA₃) substitutes for cold stratification, three replicates of 50 acorns for each species were placed in plastic containers (described in the cold stratification experiment) containing sand moistened with either distilled water (control), or a solution of 10, 100 or 1000 mg l⁻¹ GA₃. Acorns were incubated at 25/15°C for 12 weeks. This incubation temperature was used because it is too high to be effective for cold stratification (Stokes, 1965). Germination was recorded at 2-week intervals. At the conclusion of the incubation period, acorns that did not germinate were assessed for viability as described previously in the effects of cold stratification experiment.

Statistical analyses

Means and standard errors for germination proportions were calculated based on the number of viable acorns. Non-viable acorns in each replicate were $\leq 6\%$. For each *Quercus* species, a factorial repeated measures analysis of variance (ANOVA), based on the binomial distribution, was used to test for fixed effects and interaction of (1) length of cold stratification, length of incubation and incubation temperature in the germination experiment and (2)

solution concentration and length of incubation in the GA₃ experiment. Multiple comparison was used to compare germination proportions among incubation temperatures at a given incubation time. Tukey HSD was used for multiple comparison to control Type I error rates. The SAS procedure GLIMMIX was used to carry out all analyses (SAS Institute Inc., 2007). Following analyses, all proportions were converted to percentages for presentation.

Results

For purposes of clarity later in the Discussion section, results focus on germination response at the 4-weeks incubation time for identification of acorn dormancy break requirements and dormancy type (*sensu* Baskin and Baskin, 2014), and at the 12-weeks incubation time to further assess germination responses to main effects and/or their interactions over time.

Effects of cold stratification on acorn dormancy break

Germination in *Q. nigra* acorns was affected by all variables (length of stratification, incubation temperature and incubation time) and their interactions (Table 1). In the control (0-week cold stratification), mean cumulative germination percentages at 4 weeks of incubation were $\leq 7 \pm 3\%$, and ranged from $1 \pm 1\%$ (15/6°C) to $58 \pm 9\%$ (30/20°C) at 12 weeks of incubation (Fig. 1A). With 6 weeks of cold stratification, followed by 4 weeks of incubation, mean germination was higher for acorns incubated in the 30/20°C temperature regime ($48 \pm 4\%$; d.f. = 3, $F = 96.81$, $P < 0.0001$) than those at the lower incubation temperatures ($\leq 10 \pm 1\%$) (Fig. 1B). Thereafter, during incubation times of 4–12 weeks, mean germination percentages for acorns receiving 6 weeks cold stratification and incubation temperatures of 25/15 and 30/20°C increased to 90 ± 3 and $100 \pm 0\%$, respectively (Fig. 1B). During this same incubation time range, mean germination percentages for acorns incubated at 20/10 and 15/6°C increased to 45 ± 11 and $8 \pm 3\%$, respectively (Fig. 1B).

Q. nigra acorns receiving 12 weeks of cold stratification and 4 weeks of incubation germinated to a higher percentage than those in the lower incubation temperatures (d.f. = 3, $F = 136.71$, $P < 0.0001$) (Fig. 1C). However, with 12 weeks of incubation, there was no difference in mean cumulative germination percentages for acorn in the 25/15 and 30/20°C temperature regimes (Fig. 1C), and these percentages were significantly higher (d.f. = 3, $F = 73.57$, $P < 0.0001$) than those for acorns in the two lower incubation temperatures (Fig. 1C). There was no difference in mean cumulative germination percentages for acorns receiving 18 weeks cold stratification and incubated for 4 weeks at 25/15 or 30/20°C (Fig. 1D). At 12 weeks incubation, there was no difference in mean cumulative germination percentages among acorns in the 20/10, 25/15, and 30/20°C incubation temperatures (Fig. 1D). Mean cumulative germination percentages for acorns incubated at 15/6°C did not exceed $47 \pm 6\%$, regardless of incubation time (Fig. 1D).

Q. phellos acorn germination was affected by length of cold stratification, length of incubation, incubation temperature and the interaction of these variables (Table 1). In the absence of cold stratification, acorns did not germinate at an incubation time of 4 weeks, regardless of incubation temperature, and cumulative germination was $\leq 29 \pm 1\%$ among incubation temperatures with 12 weeks of incubation (Fig. 2A). With 6 weeks of cold stratification and 4 weeks of incubation, cumulative germination for

Q. phellos acorns was highest ($43 \pm 1\%$) in the 30/20°C incubation temperature (d.f. = 3, $F = 289.75$, $P < 0.0001$) (Fig. 2B). Over the course of an additional 8 weeks of incubation, cumulative germination percentages (76 ± 2 to $87 \pm 3\%$) at 20/10, 25/15 and 30/20°C were greater (d.f. = 3, $F = 197.63$, $P < 0.0001$) than those at 15/6°C (Fig. 2B).

Germination percentages ranged from 12 ± 3 to $93 \pm 0\%$ for *Q. phellos* acorns receiving 12 weeks of cold stratification and 4 weeks of incubation (Fig. 2C). However, with 12 weeks of incubation, cumulative germination percentages ($\geq 97 \pm 2\%$) were not significantly different among incubation temperatures (Fig. 2C). Cumulative germination percentages $\geq 87 \pm 2\%$ among all incubation temperatures were achieved with 18 weeks cold stratification and 4 weeks of incubation, and there was no difference in cumulative germination percentages at 12 weeks of incubation (Fig. 2D).

Effects of gibberellic acid (GA₃) on acorn dormancy break

Length of incubation (d.f. = 6, $F = 38.76$, $P < 0.0001$), and the interaction GA₃ concentration and length of incubation (d.f. = 18, $F = 3.09$, $P = 0.0007$), had significant effects on cumulative germination percentages in *Q. nigra* acorns. Germination percentages ($0\text{--}5 \pm 3\%$) among the control (distilled water) and GA₃ solutions were low and not significantly different at 4 weeks of incubation (Fig. 3). However, at 6 weeks incubation, acorn cumulative germination percentages in the 100 and 1000 mg l⁻¹ GA₃ solutions were greater than those of the control and 10 mg l⁻¹ GA₃ solution. At 8 weeks incubation, acorn cumulative germination percentages were greater in the three GA₃ solutions than that of the control (Fig. 3). At the 12 weeks incubation time, there was no difference in acorn cumulative germination percentages among treatments and the control (Fig. 3).

The overall ANOVA showed that GA₃ concentration (d.f. = 3, $F = 0.01$, $P = 0.9993$), length of incubation (d.f. = 6, $F = 0.28$, $P = 0.9463$) or the interaction of these variables (d.f. = 18, $F = 0.24$, $P = 0.9991$) had no effect on germination of *Q. phellos* acorns. Over the course of 12 weeks of incubation, acorn cumulative germination percentages among GA₃ concentrations (and control) remained $\leq 9 \pm 1\%$ (Fig. 3).

Discussion

The process of seed dormancy break is characterized by changes in germination percentages over a range of post-stratification temperatures in response to a treatment such as cold or warm stratification. When dormancy break occurs, seed germination response to a specific treatment is observed within 2–4 weeks of seed placement in a range of incubation temperatures (Baskin and Baskin, 2004, 2014). This incubation time is used to identify seed dormancy break in response to a treatment since changes in germination percentages beyond this time may reflect the interaction of treatment and incubation time (Baskin and Baskin, 2014). In light of this, dormancy type for acorns of *Q. nigra* and *Q. phellos* was determined by acorn cumulative germination percentages at 4 weeks of incubation.

With 12 or 18 weeks of cold stratification and 4 weeks of incubation, cumulative germination percentages for *Q. nigra* acorns were $\geq 77\%$ but only in incubation temperatures of 25/15 and 30/20°C. However, with 18 weeks of cold stratification and 10 or 12 weeks of incubation, cumulative germination percentages for acorns incubated at 20/10, 25/15 and 30/20°C were not significantly different ($\geq 95 \pm 2\%$). This suggests that acorns of *Q. nigra*

Table 1. Results of ANOVAs showing the effects and interactions of length of cold stratification (0, 6, 12 and 18 weeks), length of incubation (2, 4, 6, 8, 10 and 12 weeks) and incubation temperature (15/6, 20/10, 25/15 and 30/20°C) on germination of *Q. nigra* and *Q. phellos* acorns

Source of variation	<i>Quercus nigra</i>			<i>Quercus phellos</i>		
	d.f.	<i>F</i>	<i>P</i>	d.f.	<i>F</i>	<i>P</i>
Length of stratification (<i>S</i>)	3	1560.10	<0.0001	3	7076.62	<0.0001
Incubation temperature (<i>T</i>)	3	1795.59	<0.0001	3	558.64	<0.0001
Length of incubation (<i>I</i>)	6	897.81	<0.0001	6	1954.14	<0.0001
<i>S</i> × <i>T</i>	9	99.27	<0.0001	9	105.98	<0.0001
<i>S</i> × <i>I</i>	18	64.09	<0.0001	18	332.25	<0.0001
<i>T</i> × <i>I</i>	18	73.90	<0.0001	18	22.93	<0.0001
<i>S</i> × <i>T</i> × <i>I</i>	54	19.25	<0.0001	54	33.63	<0.0001




Fig. 1. Mean (\pm SE) cumulative germination percentages for *Q. nigra* acorns receiving (A) 0 week (control), (B) 6 weeks, (C) 12 weeks or (D) 18 weeks of cold stratification at 5/1°C and incubated for 12 weeks in four alternating temperature regimes. Means at 4 or 12 weeks incubation with dissimilar letters are significantly different (Tukey's HSD, $P = 0.05$).

may require >18 weeks of cold stratification to germinate to higher percentages over a wider range of temperatures (<20/10°C) and achieve dormancy break. On the other hand, dormancy break in *Q. phellos* acorns occurred with 18 weeks of cold stratification. Cumulative germination percentages for acorns of this species were $\geq 87\%$ at 4 weeks of incubation in all test temperature regimes. Gibberellic acid was not an effective substitute for cold stratification in either species, as germination percentages were $\leq 5 \pm 3\%$ for both species at the 4 weeks incubation time. Collectively, these data indicate that acorns of *Q. nigra* and *Q. phellos* possess deep PD (*sensu* Baskin and Baskin, 2004, 2014).

In the *Lobatae* clade, *Q. nigra* and *Q. phellos* are closely related species, appearing on separate, but adjacent branches stemming

from a node with an estimated divergence in the mid-Miocene (Hipp et al., 2018). *Q. nigra* is a sister taxon to *Q. arkansana* Sarg., and *Q. phellos* is a sister taxon to *Q. elliotii* Wilbur (syn. *Q. pumila* Walter) (Hipp et al., 2018). Both *Q. arkansana* and *Q. elliotii* are upland forest species and, thus, occupy a different ecological niche than *Q. nigra* and *Q. phellos*. Dormancy type for *Q. elliotii* is not known; however, cumulative germination percentages in *Q. arkansana* are significantly greater for acorns receiving 90–120 days of cold stratification (Wirges and Yeiser, 1984). This suggests that acorns of *Q. arkansana* may possess deep PD, and the level of PD may be the same as that of the sister taxon, *Q. nigra*. Summarily, deep PD is present in two closely related *Quercus* species of similar habitat (*Q. nigra* and




Fig. 2. Mean (\pm SE) cumulative germination percentages for *Q. phellos* acorns receiving (A) 0 week (control), (B) 6 weeks, (C) 12 weeks or (D) 18 weeks of cold stratification at 5/1°C and incubated for 12 weeks in four alternating temperature regimes. Means at 4 or 12 weeks incubation with dissimilar letters are significantly different (Tukey's HSD, $P=0.05$).




Fig. 3. Mean (\pm SE) cumulative germination percentages for *Q. nigra* (solid line) and *Q. phellos* (broken line) acorns receiving 0-week cold stratification and incubated for 12 weeks at 25/15°C in distilled water (control), or a solution of 10, 100 or 1000 mg l⁻¹ of gibberellic acid (GA₃). Means at 12 weeks incubation with dissimilar letters are significantly different (Tukey's HSD, $P=0.05$).

Q. phellos), and potentially the same is closely related *Quercus* species of dissimilar habitats (*Q. phellos* and *Q. arkansana*).

It should be noted that *Q. pagoda*, a bottomland species that often grows with *Q. nigra* and *Q. phellos*, appears on a separate, less diverse branch of the *Lobatae* clade than *Q. nigra* and *Q. phellos*, and one that diverged from the latter two species branch ca. mid-Miocene (Hipp et al., 2018). *Q. pagoda* is sister taxon to *Q. ilicifolia*, which grows on dry, sandy barrens, or rocky hillsides.

Acorns of *Q. pagoda* possess non-deep PD (Hawkins, 2019a), and those of *Q. ilicifolia* possess an unknown level of PD, and possibly some level of epicotyl dormancy (Allen and Farmer, 1977). Although, epicotyl dormancy is not expressed in *Q. pagoda* acorns, the time between germination and epicotyl emergence increases with decreasing post-germination temperature, but this is unrelated to the length of cold stratification received (Hawkins, 2019a). This suggests similar (PD), although not identical, dormancy types between sister taxa (*Q. pagoda* and *Q. ilicifolia*) that grow in dissimilar habitats, and different levels of PD in less closely related species (*Q. pagoda* vs *Q. nigra* and *Q. phellos*) that grow together in mixed stands in the eastern United States bottomland forests.

Differences in dormancy type or level of PD in congeners with respect to species relatedness or habitat are not an atypical occurrence in temperate deciduous forests (Li et al., 1999; Hawkins et al., 2010a). However, recognition of dormancy break and germination requirements in *Quercus* species within and among habitats is highly relevant to large-scale oak research and artificial regeneration efforts. In both instances, comparative morphometric measurements are sometimes used to assess seedling response to abiotic variables (Anderson and Pezeshki, 2000) and/or identify traits that enhance the success of seedling establishment of outplanted nursery generated seedlings (Dey et al., 2008). Comparative measurements of an even-aged seedling cohort, particularly within the first months following seedling emergence, ensures that morphological and/or physiological differences among seedlings are not age-related. Intraspecifically, germination synchrony and thus an even-aged seedling cohort may be achieved by ensuring dormancy break in acorns of the *Quercus* species of interest, followed by post-germination temperatures

appropriate for epicotyl emergence (Hawkins, 2019a). However, if dormancy break requirements differ between acorns of one or more species, it may be possible to subject acorns of the *Quercus* species to the same length of cold stratification, but then adjust the post-stratification temperature to achieve interspecific germination synchrony. For example, in this study, 18 weeks of cold stratification was sufficient for dormancy break in acorns of *Q. phellos*, but not for those of *Q. nigra*. However, with 18 weeks of cold stratification, $84 \pm 3\%$ of *Q. nigra* acorns and $97 \pm 2\%$ of *Q. phellos* acorns germinated within 2 weeks in an incubation temperature of 30/20°C (i.e. synchronous germination).

The interaction of variables (length of cold stratification, post-stratification temperature and time) that contribute to acorn dormancy loss make it difficult to predict the influence of different levels of PD on germination synchronicity, and subsequent seedling emergence among *Q. nigra* and *Q. phellos* acorns in a bottomland habitat. At the time of dispersal, acorns begin receiving some level of stratification. Although autumn dispersal of *Q. nigra* and *Q. phellos* acorns is relatively synchronous (Cypert and Webster, 1948; Hawkins, pers. observation), time elapsed from dispersal to spring temperatures that satisfy germination temperature requirements will vary with year and latitude. Similarly, the length of time that acorns receive temperatures effective for cold stratification (0–10°C; Stokes, 1965) may vary annually within and among latitudes throughout the species' ranges.

An additional abiotic influence on acorn germination dynamics of bottomland *Quercus* species is submergence from flooding which may vary annually in timing and duration (Hawkins et al., 2010b). For example, 63 days of winter submergence (water temperature range, 4–13°C) was an effective substitute for cold stratification and, thus, sufficient for dormancy break in *Q. pagoda* acorns (Hawkins, 2019b). In contrast, dormancy break was not achieved in *Q. phellos* acorns with 63 or 84 days of winter submergence (Hawkins, 2019b). Guo et al. (1998) reported that germination percentages for *Q. nigra* acorns receiving up to 30 days of winter submergence were not significantly greater than those receiving 0-day winter submergence. Additionally, submergence duration may vary within acorn cohorts due to the heterogeneous topography associated bottomland hardwood forests in the eastern United States (Gardiner, 2001; Gardiner et al., 2004).

Conclusions

PD is the most prevalent dormancy type in seeds of woody species in the canopy and subcanopy of the eastern United States temperate deciduous forests (Baskin and Baskin, 2014). Among species of the *Lobatae* section, acorn dormancy break and germination requirements may differ between species that possess the same level of PD, and *Quercus* species of the same habitat may possess different levels of PD. In this study, acorns of the closely related *Q. nigra* and *Q. phellos* possessed the same level of PD, and a different level of PD than that of a more distantly related *Quercus* species. However, an accurate description of biogeographical or phylogenetic associations of acorn dormancy type (or level of PD) among species of *Lobatae* will require ecologically relevant studies of more representatives of this section. Additionally, information gained from these studies may provide methods for refining artificial regeneration practices aimed at restoration of *Quercus* spp. in forests of the eastern United States.

Acknowledgements. The author thanks Rory Thornton for help with data collection, Dr. Charles Sabatia for direction in statistical analyses and Dr.

Jeffrey Walck and Dr. Siti Hidayati for review of an earlier draft of this manuscript.

References

- Abrams MD (1992) Fire and the development of oak forests. *BioScience* **42**, 346–353.
- Allen R and Farmer Jr RE (1977) Germination characteristics of bear oak. *Southern Journal of Applied Forestry* **1**, 19–20.
- Anderson PH and Pezeshki SR (2000) The effects of intermittent flooding on seedlings of three forest species. *Photosynthetica* **37**, 543–552.
- Arthur MA, Alexander HD, Dey DC, Schweitzer CJ and Loftis DL (2012) Refining the oak-fire hypothesis for management of oak-dominated forests of the eastern United States. *Journal of Forestry* **110**, 257–266.
- Baskin JM and Baskin CC (2004) A classification system for seed dormancy. *Seed Science Research* **14**, 1–16.
- Baskin CC and Baskin JM (2014) *Seeds: ecology, biogeography, and evolution of dormancy and germination* (2nd edn). San Diego, Elsevier Academic Press.
- Bonner FT and Karrfalt RP (2008) *The woody plant seed manual*. Washington, DC, U.S. Forest Service. Agriculture Handbook 727.
- Bonner FT and Vozzo JA (1987) *Seed biology and technology of Quercus*. U.S. Department of Agriculture Forest Service. Gen. Tech. Rep. SO-66.
- Bruhn JN, Wetteroff Jr JJ, Mihail JD, Kabrick JM and Pickets JB (2008) Distribution of *Armillaria* species in upland Ozark Mountain forests with respect to site, overstory species composition and oak decline. *European Journal of Forest Pathology* **30**, 43–60.
- Cavender-Bares J (2016) Diversity, distribution, and ecosystem services in the North American Oaks. *International Oaks* **27**, 37–48.
- Cypert E and Webster BS (1948) Yield and use by wildlife of acorns of water and willow oaks. *Journal of Wildlife Management* **12**, 227–231.
- Dey DC, Jacobs D, McNabb K, Miller G, Baldwin V and Foster G (2008) Artificial regeneration of major oak (*Quercus*) species in the eastern United States – a review of the literature. *Forest Science* **54**, 77–106.
- Gardiner ES (2001) Ecology of bottomland oaks in the southeastern United States. *International Oaks* **12**, 48–55.
- Gardiner ES, Hodges JD and Fristoe TC (2004) Flood plain topography affects establishment success of direct-seeded bottomland oaks, in Connor KF (Ed.) *Proceedings of the 12th Biennial Southern Silvicultural Research Conference*. Asheville, North Carolina, USDA Forest Service. Gen. Tech. Rep. SRS-71.
- Guo Y, Shelton MG and Lockhart BR (1998) Effects of flood duration and season on germination of black, cherrybark, northern red, and water oak acorns. *New Forests* **15**, 69–76.
- Guyette RP, Muzika RM and Dey DC (2002) Dynamics of an anthropogenic fire regime. *Ecosystems* **5**, 472–486.
- Haavik LJ, Jones JS, Galligan LD, Guldin JM and Stephen FM (2012) Oak decline and red oak borer outbreak: impact in upland oak-hickory forests of Arkansas, USA. *Forestry* **85**, 341–352.
- Hawkins TS (2019a) Regulating acorn germination and seedling emergence in *Quercus pagoda* (Raf.) as it relates to natural and artificial regeneration. *New Forests* **50**, 425–436.
- Hawkins TS (2019b) The influence of dormancy break requirements on germination and viability responses to winter submergence in acorns of three bottomland red oak (Sect. *Lobatae*) species. *Forest Science* **65**, 556–561.
- Hawkins TS, Baskin CC and Baskin JM (2010a) Morphophysiological dormancy in seeds of three eastern North American *Sanicula* species (Apiaceae subf. Saniculoideae): evolutionary implications for dormancy break. *Plant Species Biology* **25**, 103–113.
- Hawkins TS, Skojac Jr DA, Schiff NM and Leininger TD (2010b) Floristic composition and potential competitors in *Lindera melissifolia* (Lauraceae) colonies in Mississippi with reference to hydrologic regime. *Journal of the Botanical Research Institute of Texas* **4**, 381–390.
- Hipp AL, Manos PS, González-Rodríguez A, Hahn M, Kaproth M, McVay JD, Avalos SV and Cavender-Bares J (2018) Sympatric parallel diversification of major oak clades in the Americas and the origins of Mexican species diversity. *New Phytologist* **217**, 439–452.

- Hodges JD** (1994) The southern hardwood bottomland region and brown loam bluffs subregion, pp. 227–269 in Barrett JW (Ed.) *Regional silviculture of the United States* (3rd edn). New York, John Wiley and Sons, Inc.
- Hopper GM, Smith DW and Parrish DJ** (1985) Germination and seedling growth of northern red oak: effects of stratification and pericarp removal. *Forest Science* **31**, 31–39.
- Kelley MB, Fierke MK and Stephen FM** (2009) Identification and distribution of *Armillaria* species associated with an oak decline event in the Arkansas Ozarks. *Forest Pathology* **39**, 397–404.
- King SL and Keeland BD** (1999) Evaluation of reforestation in the Lower Mississippi River Alluvial Valley. *Restoration Ecology* **7**, 348–359.
- Li X, Baskin JM and Baskin CC** (1999) Physiological dormancy and germination requirements of seeds of several North American *Rhus* species (Anacardiaceae). *Seed Science Research* **9**, 237–245.
- McGee CE** (1972) From a defective hardwood stand to multiple use opportunity. *Journal of Forestry* **70**, 700–704.
- Peterson JK** (1983) Mechanisms involved in delayed germination of *Quercus nigra* L. seeds. *Annals of Botany* **52**, 81–92.
- SAS Institute, Inc.** (2007) *The SAS System for Windows, Release V9.4*. Cary, North Carolina, SAS Institute.
- Schlaegel BE** (1990) *Quercus phellos* L., willow oak, pp. 715–720 in R.M. Burns and Honkala BH (Eds) *Silvics of North America: Volume 2. Hardwoods*. Agriculture Handbook 654. Washington, DC, USDA Forest Service.
- Stanturf JA, Gardiner ES, Hamel PB, Devall MS, Leininger TD and Warren ME** (2000) Restoring bottomland hardwood ecosystems in the Lower Mississippi Alluvial. *Journal of Forestry* **98**, 10–16.
- Starkey DA, Oliveria F, Mangini A and Mielke M** (2004) Oak decline and red oak borer in the interior highlands of Arkansas and Missouri: natural phenomena, severe occurrences, pp. 217–222 in Spetich MA (Ed.) *Upland oak ecology symposium: history, current conditions, and sustainability*. Asheville, North Carolina, USDA Forest Service, Southern Research Station. Gen Tech. Rep. SRS-73.
- Stein JD, Binion D and Acciavatti R** (2003) *Field guide to native oak species of eastern North America*. Morgantown, West Virginia, USDA Forest Service. FHTET-2003-01.
- Stephen FM, Salisbury VB and Oliveria FL** (2001) Red oak borer, *Enaphalodes rufulus* (Coleoptera: Cerambycidae), in the Ozark mountains of Arkansas, USA: an unexpected and remarkable forest disturbance. *Integrated Pest Management Reviews* **6**, 247–252.
- Stokes P** (1965) Temperature and seed dormancy, pp. 746–803 in Ruhland W (Ed.) *Encyclopedia of plant physiology*, vol. **15**, part 2. Berlin, Springer.
- Vozzo JA** (1990) *Quercus nigra* L., water oak, pp. 701–703 in Burns RM and Honkala BH (Eds) *Silvics of North America: Volume 2. Hardwoods*. Agriculture Handbook 654. Washington, DC, USDA Forest Service.
- Wirges G and Yeisiker J** (1984) Stratification and germination of Arkansas oak acorns. *Tree Planters' Notes* **35**, 36–38.