

How seed orchard culture affects seed quality: Experience with the southern pines

by J.P. Barnett¹

Tree improvement programs have influenced significantly the quality of southern pine seeds produced when compared to collections from native stands. Seed orchard management practices such as fertilization can increase seed size and reduce seed dormancy. These result in the need for less complex pregermination treatments. Repeated cone collections from the same clones facilitate collections according to ripening (cone specific gravity), which can improve seed germination and storage. However, cultural practices may result in seed properties that are more sensitive to damage during processing procedures and result in lower quality unless special care is provided during this stage of handling. The effect of orchard management practices on seed quality also varies by species, with loblolly pine being less affected than longleaf pine.

Key words: *Pinus* spp., seed germination, seed dormancy, seed storage, cone maturity

Les programmes d'amélioration génétique ont eu une influence significative sur la qualité des semences de pins du sud lorsqu'on les compare aux semences recueillies dans les peuplements naturels. Les pratiques d'aménagement des vergers à graines telle la fertilisation peuvent accroître les dimensions de la graine et réduire la dormance. En conséquence, il est moins nécessaire de recourir à des traitements complexes de pré-germination. Les cueillettes répétées des cônes sur les mêmes clones facilitent les récoltes de graines pour ce qui est de la maturité (poids spécifique des cônes), ce qui peut améliorer la germination et l'entreposage. Toutefois, les pratiques de production peuvent modifier les propriétés des semences en les rendant plus susceptibles d'être endommager lors de la manipulation et entraîner une baisse de la qualité à moins qu'une attention spéciale n'y soit accordée au cours de la manipulation. L'effet des pratiques d'aménagement des vergers sur la qualité des semences varie selon l'espèce étant moins affecté que les pins à longues aiguilles.

Mots clés: *Pinus* spp., germination de la graine, dormance, entreposage des semences, maturité des cônes

Introduction

Remarkable progress was made in tree improvement of the southern pines (*Pinus* spp.) during the past four decades. Seed orchards produce seeds for the vast majority of the more than 1 billion pine seedlings produced annually in the southern United States (Dougherty and Duryea 1991). In fact, the more than 4,000 ha of southern pine seed orchards now produce seeds of most species in quantities in excess to our current needs. During the development of these orchards and incorporation of improved seeds into nursery practice, there were changes in seed management practices to adjust to differences in seed properties caused by more intensive tree culture.

Much of the definitive research on physiology of southern pine seeds was conducted with seeds collected from native stands. Through the years, there was a modification of practices to reflect the character of seeds from managed orchards. Since this was a gradual process, there has been no critical evaluation of the impact of tree improvement on seed quality. However, there were a number of changes in methodology that reflect the impact of more intensive tree culture. The purpose of this paper is to review the influence of seed orchard management on seed quality. Since there are few studies designed to compare these effects, many of the influences are documented from personal observation, evaluation of data from related studies, and from changes in seed processing practices through time.

Orchard Management

The goal in managing seed orchards is to produce high-quality seeds with a desirable genotype mix in quantities suf-

ficient for regeneration needs in the shortest time (Pait *et al.* 1991). Rapid growth of young trees is achieved by controlling weeds, insects, and water stress and ameliorating soil nutrient deficiencies. Management activities change as orchards begin to produce cones. Fertilization continues to maintain desirable levels of nutrients for overall tree vigor and cone production. Irrigation to reduce moisture stress, subsoiling to overcome compaction, and applications of insecticides to control insects are additional practices that may improve vigor and enhance cone and seed production of orchard trees.

Orchard management that included use of fertilizers, insecticides, and other cultural practices (e.g. mowing five times year⁻¹, subsoiling, and pruning or thinning) doubled loblolly pine (*Pinus taeda* L.) cone yields and tripled numbers of viable seeds cone⁻¹ over that produced in an untreated (mowed only) orchard (Fatzinger *et al.* 1985). Use of insecticides markedly improved yield of seeds cone⁻¹ over unprotected stands or orchards (Bramlett *et al.* 1977). However, there is little data to indicate that quality of seeds produced differs from unprotected seeds as a result of insecticide use.

Cone Collection and Seed Maturity

Wakeley (1954) reported the germination of southern pine seeds is directly related to cone maturity at the time of extraction and cone crops are mature enough to collect when specific gravity (SG) of 19 of 20 random sample cones drops below 0.89. The critical determinant of maturity is cone moisture content, but measuring specific gravity by flotation is the quickest way to estimate cone moisture. Wakeley (1954) determined specific gravity by flotation in oil, but another simple method is based on cone immersion in water (Barnett 1979a). Weight of cones that will float in water (SG < 1.00) is the difference in water level (cc) in a graduated cylinder before and after the addition

¹USDA Forest Service, Southern Research Station, Pineville, LA, USA 71360.

of a cone that is allowed to float. Volume is the difference between the initial water level and the level after the cone is submerged with a wire or small stick. Dividing cone weights by volume results in specific gravity.

Typically, cones were collected over a two- to three-week period and seed extraction was accomplished within a few weeks after collection. However, as the demand for southern pine seeds escalated and seed orchards came into production, this period was too short to allow collection of enough cones to provide an adequate seed supply. It became necessary then, to extend the collection period by beginning a few weeks before the traditional specific gravity maturity index.

An extended collection period is possible if slightly immature cones are stored or held prior to kilning (Barnett 1976a, McLemore 1975a). Collections of loblolly and slash (*P. elliottii* Engelm.) pine cones can begin two or three weeks before maturity, but seed yields from these immature cones may decrease even after cone storage. Early collections are advisable only if large quantities of seeds are needed immediately or if labor or the collectable crop is limited. Seeds from immature loblolly pine cones (which are collected before reaching a specific gravity of 0.88) are mature when extractable, and slash pine seeds continue ripening during cone storage. However, viability of longleaf pine (*P. palustris* Mill.) seeds from immature cones decreases during cone storage. Therefore, only mature longleaf cones should be collected. Once ripe, longleaf cones can be stored or precured for three to five weeks to increase seed yields without reducing viability, but the storage period for mature cones should not exceed eight weeks (McLemore 1961).

The method of cone storage has varied from burlap bags to seven hL wire-bound boxes, outdoors or in unheated indoor facilities. Although several studies of the influence of the type of cone storage on seed quality have been conducted (Barnett 1979b, Bonner 1987), there has been no clear indication of the superiority of any one method.

The greatest emphasis in southern pine seed orchard production has been on loblolly pine. It is fortunate this species is, of the southern pines, the least affected by seed orchard practices that require extension of cone collection and processing. Such extended periods are feasible without any major reduction in seed quality of loblolly pine. However, seed quality of slash and longleaf pines can be markedly reduced by changes in collecting and processing variables.

Orchard managers normally determine the relative sequence and approximate dates of ripening of all their clones. Once this clonal cone ripening sequence is determined, collections should be scheduled to start with those maturing early and follow the order of ripening. The time between ripening dates of the earliest and latest clones in one orchard was documented as 50 days (Zoerb 1969). The use of this collection strategy will reduce the adverse effects of early cone collections.

An alternative to collecting cones by hand is the net retrieval system that is used to collect loblolly, Virginia (*P. virginiana* Mill.), and eastern white (*P. strobus* L.) pine seeds (Pait *et al.* 1991, Wynens and Brooks 1979). Polypropylene netting is spread on the orchard ground before cone opening. Seeds fall to the netting naturally as the cones open (McConnell and Edwards 1985). When most of the seed crop is judged to be on the netting, it is rolled up and the seeds are recovered. High seed moisture contents are common and can be troublesome during retrieval and processing. However, with proper care, good qual-

ity can be maintained in loblolly pine seeds for use immediately or after at least 1.5 years of storage (Bonner and Vozzo 1986).

Cone and Seed Processing

One would assume that orchard management would have little influence on how cones and seeds should be processed. However, management practices do change some cone and seed properties (McLemore 1975b) and these can increase the susceptibility of seeds to extremes in processing.

Extraction

Seeds are usually extracted from southern pine cones in forced-draft kilns. Temperature and duration of kilning are critical for southern pine cones, particularly longleaf: temperatures of 46°C or more markedly reduce germination (Rietz 1941) and those under 32°C may not result in effective cone opening (Bonner 1987). Optimal temperatures are 35°C to 41°C. Increases in the length of treatment may also reduce viability. A general recommendation is that cones and seeds should be removed from kilns as soon as cones open sufficiently to release the seeds.

Cleaning

After seeds are extracted, they must be dewinged, cleaned and dried. The wings on seeds of all southern pines, except longleaf, are completely removed by brushing and tumbling in mechanical dewingers. The structure of longleaf seeds makes dewinging difficult; the wings are mechanically reduced to stubs, so dewingers must be carefully regulated to prevent injury to these thin-coated seeds. Wing removal that is done carefully and does not damage the seed coat has no effect on seed storability (Barnett 1969; Belcher and King 1968), but dewinging is a common cause of seed injury and loss of viability. Orchard management practices generally result in larger seeds and thus increase seed sensitivity to damage during processing activities such as dewinging.

Before seeds are used or stored, empty seeds should be removed from the seedlot. This is the easiest means of upgrading a seedlot. This can be accomplished by mechanical cleaning equipment, or, when lots are small, as in progeny tests, it is often convenient to use flotation in water or organic solvents to separate unfilled seeds. In the appropriate liquid, sound seeds sink, while unsound ones float and can easily be skimmed off. Examples of appropriate solvents for seed separations are: water for loblolly pine, *n*-pentane for longleaf pine, and 95-percent ethanol for slash, shortleaf (*P. echinata* Mill.), sand (*P. clausa* [Chapm. ex Engelm.] Vasey ex Sarg.), and spruce (*P. glabra* Walt.) pines (Barnett and McLemore 1970). To maintain seed quality, flotation in ethanol should be delayed until just prior to use, because if the ethanol is not thoroughly removed by drying, seeds so treated rapidly lose viability in storage (Barnett 1971).

Sizing

The reported effects of seed size on germination and early seedling growth are conflicting. The operational objective of sizing is to produce a uniform crop of seedlings. Seed size has a more pronounced effect on germination than seedling development. Uniform speed of germination may, therefore, be the most important consideration in sizing. More recent tests under laboratory conditions of minimal environmental stress




Fig. 1. Mean hypocotyl length of loblolly pine germinants 28 days after germination (from Dunlap and Barnett 1983). Vertical bar represent one SE.

have shown that germinant size after 28 days of growth was strongly correlated with seed size (Dunlap and Barnett 1983). The faster germinating seeds in each size class produced larger germinants after 28 days of incubation (Fig. 1). All seeds reached a maximum germination rate by the sixth day, but smaller seeds were slower to initiate germination (Fig. 2). These results are in agreement with Venator's (1973) findings indicating faster growing Caribbean pine (*P. caribaea* var. *hondurensis* Barrett and Golfair) seedlings tend to develop from early germinating seeds. Consequently, seedling size and possibly uniformity of growth are primarily a function of germination patterns, which are partially determined by seed size.

Seed size varies by genetic and geographic source, and fertilization practices commonly used in seed orchards increase seed size (McLemore 1975b). When orchard collections are bulked into a single lot, sizing can result in the exclusion of certain families. Some seed managers may be tempted to discard the small-seed fraction of their lots; but, such a practice could eliminate certain families and narrow the genetic base of the planting stock (Silen and Osterhans 1979). All sizes of seeds should be used in seedling production.

Seed processing techniques have been developed specifically for loblolly pine since it is the species more often produced in southern pine seed orchards. Of the southern pines, loblolly seeds are the most dormant, hardseeded, and the least susceptible to damage during processing. It is not surprising then that application of the processing technology developed primarily for loblolly pine to a species like longleaf pine may result in damage that reduces seed quality and performance.

Seed Storage

Inconsistent cone crops require that seed managers store seeds to offset years of low production. Fortunately, seeds of southern pines are relatively easy to store for long periods. Viability of slash and shortleaf seeds stored for 50 years at above-freezing temperatures was 66 and 25%, respectively (Barnett and Vozzo 1985). Longleaf seeds, which are the most difficult




Fig. 2. Mean daily germination of loblolly pine seeds from the large, medium, and small classes incubated under standard laboratory conditions (from Dunlap and Barnett 1983). Vertical bar represent one SE.

of the southern pines to store, have been held for 20 years at low seed moisture and subfreezing temperatures without significant losses in viability (Barnett and Jones 1993).

Careful control of seed moisture content and storage temperatures is essential to maintain viability (Barnett and McLemore 1970, Barton 1961, Jones 1966). General recommendations for long-term storage are to dry seeds to $\leq 10\%$ moisture content and store at subfreezing temperatures. Seeds that are damaged or are known to have low vigor can be preserved by drying to 8 to 10% and lowering storage temperatures to about -18°C (Kamra 1967). The genetic source of the seeds seems to have little effect on their storage capabilities (Barnett and McLemore 1970).

Seed Dormancy and Pretreatment

Seed dormancy in the southern pines seems determined by the magnitude of seed coat constraint. The ratio of the seed coat to total seed weight provides a means of rapidly estimating relative seed dormancy among closely related species (Barnett 1972, 1976b; Carpita *et al.* 1983). The theory is that thicker and heavier seed coats of the more dormant seeds restrict imbibition by preventing swelling of the megagametophyte and embryo and thus limit water absorption sufficient for germination. Loblolly pine seeds are considered the most dormant, and longleaf pine seeds, the least dormant of the southern pines. Although not thoroughly documented, one of the more significant effects of seed orchard culture may be on seed dormancy. Fertilization increases seed size (McLemore 1975b). Dunlap and Barnett (1983) found larger seeds tend to germinate more quickly and produce larger germinants than smaller ones, although final germination is typically lower in the larger seeds. Therefore, seed orchard management results in seeds with less dormancy than those from native stands.

Maternal factors such as seed coat properties that influence the speed of germination can obscure the nature of genetic control of subsequent growth processes (Perry 1976). Seed dormancy varies by geographic location or ecotype (Barnett

1991). Less than 15% of the weight of a conifer seed is in the embryo, which is the only portion with a genetic component from the male plant. In nature, moist prechilling or stratification is usually optimized as a result of natural conditions, but in nursery production, the genetic component from the male parent may be obscured when managers do not optimize the prechilling needs of the seedlot.

Overcoming seed dormancy is one of the major steps to ensure prompt and uniform germination. Presowing treatments to speed germination are discussed in detail by several authors (Bonner *et al.* 1974, Tanaka 1984). Typically, prechilling is done after an 8 to 24 hour period of moisture imbibition. Fully imbibed seeds are placed in polyethylene bags and held at temperatures of 1° to 5°C. The length of treatment varies by the extent of dormancy present in the seeds.

In recent years, techniques other than prechilling were investigated in an attempt to accelerate the dormancy-breakage process or obtain more desirable germination patterns for genetically improved seeds (Barnett 1989). However, none of these newer presowing treatments have been developed to the point where they are as consistently effective as conventional moist prechilling.

Predicting Seed Performance

For decades, nursery managers and seed physiologists have sought techniques, generally with little success, that would more accurately predict seed performance in the nursery. In an evaluation of the problem, Barnett and McLemore (1984) found that laboratory germination tests performed on prechilled seedlots provide the best predictors of nursery-tree yield for southern pines. Germination tests are standardized by conducting them under optimum light and temperature conditions (AOSA 1980). However, tests conducted under optimum conditions do not reflect germination of dormant seeds on nursery beds where temperatures and photoperiods are often less than optimal (Table 1). Since orchard culture results in changes in seed properties, comparative tests of untreated and prechilled (various durations) seeds are needed to develop optimal recommendations. A technique to improve prediction of seed performance is to determine prechilling needs under stress conditions that relate to the nursery conditions (Barnett 1992). The prechilling period should be extended to minimize the effect of the less than optimal conditions on initial seedling development.

Conclusions

Tree improvement programs have significantly influenced the properties of southern pine seeds produced compared to seeds from native stand collections. The effects of orchard cultural practices include:

1. Varying responses among species to management practices. Fortunately, loblolly pine, which is the major southern species produced in tree improvement programs, is most tolerant to cultural manipulation. Longleaf pine seeds are the most sensitive to impacts of management.
2. Reductions in seed quality related to extended cone collections result from the processing of immature cones and seeds. Repeated annual collections from the same clones allow orchard managers to collect by ripening sequence and thus overcome some of the problems of collecting large quantities of immature cones.

Table 1. Effect of length and method of stratification of a mixed loblolly pine seedlot in two testing environments (McLemore 1969)

Days stratification	Stratified in refrigerator at 1°C		Stratified outdoors	
	Germination percent	Germination value ¹	Germination percent	Germination value
<i>Tested at 15°C with 11-h photoperiod</i>				
0	<1	0.0	<1	0.0
30	68	7.1	59	6.0
60	95	17.3	91	11.4
113	99	24.0	98	19.6
<i>Tested at 22°C with 16-h photoperiod</i>				
0	96	20.8	96	20.8
30	99	37.6	98	41.8
60	99	47.1	99	47.0
113	100	50.3	99	56.3

¹Germination values represent the speed as well as completeness of germination (Czabator 1962).

3. Increases in seed size, resulting from orchard fertilization, reduce dormancy levels in loblolly pine seeds, increase sensitivity of longleaf pine seeds to damage during processing, and change the nature of presowing treatments in most species.
4. Insignificant effects on seed storability.
5. The reevaluation of prechilling needs based on comparative germination tests is required and germination should be optimized for nursery-bed conditions.

The effects of culture treatments on seed quality noted for the southern pines indicate areas that should be a concern of researchers and orchard managers working in other regions and with other species. Seeds from orchards may require different treatment than for those from native stands if high quality is to be maintained.

References

- Association of Official Seed Analysts. 1980.** Rules for testing seeds. *J. Seed Technol.* 3(3): 1–126.
- Barnett, J.P. 1969.** Long-term storage of longleaf pine seeds. *Tree Planters Notes* 20(2): 22–25.
- Barnett, J.P. 1971.** Flotation in ethanol reduces storability of southern pine seeds. *For. Sci.* 17: 50–51.
- Barnett, J.P. 1972.** Seed coat influences dormancy of loblolly pine seeds. *Can. J. For. Res.* 2: 7–10.
- Barnett, J.P. 1976a.** Cone and seed maturation of southern pines. USDA Forest Service Res. Pap. SO-122. Southern Forest Experiment Station, New Orleans, LA. 11 p.
- Barnett, J.P. 1976b.** Delayed germination of southern pine seed related to seed coat constraint. *Can. J. For. Res.* 6: 504–510.
- Barnett, J.P. 1979a.** An easy way to measure cone specific gravity. Pp. 21–23 *In: Proceedings of the seed collection workshop, 16–18 May 1979, Macon, GA.* USDA Forest Service Tech. Bull. SA-TP 8. State and Private Forestry, Atlanta, GA.
- Barnett, J.P. 1979b.** Southern pine cone maturation and storage. Pp. 11–20 *In: Proceedings of the seed collection workshop, 16–18 May 1979, Macon, GA.* USDA Forest Service Tech. Bull. SA-TP 8. State and Private Forestry, Atlanta, GA.
- Barnett, J.P. 1989.** Seed, cultural practices, and seedling uniformity. *Forestry* 62 (Suppl.): 95–105.
- Barnett, J.P. 1991.** Relating the seed coat of *Pinus* to speed of germination, geographic variation, and seedling development. Pp. 266–275 *In: Proceedings of the 21st southern forest tree improvement conference, 17–20 June 1991, Knoxville, TN.* Southern Forest Tree Improvement Committee, Knoxville, TN.

- Barnett, J.P. 1992.** Production of shortleaf pine seedlings. Pp. 20–31 *In: Proceedings of the shortleaf pine regeneration workshop, 29–31 October 1991, Little Rock, AR. USDA Forest Service Gen. Tech. Rep. SO-90. Southern Forest Experiment Station, New Orleans, LA.*
- Barnett, J.P. and J.P. Jones. 1993.** Response of longleaf pine seeds to storage conditions and pregermination treatments. *South. J. Appl. For.* 17: 180–187.
- Barnett, J.P. and B.F. McLemore. 1970.** Storing southern pine seeds. *J. For.* 68: 24–27.
- Barnett, J.P. and B.F. McLemore. 1984.** Germination speed as a predictor of nursery seedling performance. *South. J. Appl. For.* 8: 157–162.
- Barnett, J.P. and J.A. Vozzo. 1985.** Viability and vigor of slash and shortleaf pine seeds after 50 years of storage. *For. Sci.* 31: 316–320.
- Barton, L.V. 1961.** Seed preservation and longevity. Leonard Hill Books, Ltd., London. 216 p.
- Belcher, E.W. and J. King. 1968.** Storage of dewinged longleaf pine seed is possible. Georgia Forestry Research Council Res. Pap. 51. Macon, GA. 4 p.
- Bonner, F.T. 1987.** Cone storage and seed quality in longleaf pine. USDA Forest Service Res. Note SO-341. Southern Forest Experiment Station, New Orleans, LA. 4 p.
- Bonner, F.T. and J.A. Vozzo. 1986.** Effects of the net retrieval system on quality of seeds collected in pine orchards of the southern United States. Pp. 91–97 *In: Proceedings international symposium on seed problems under stressful conditions. IUFRO P.G. P02.04.00 Schrift Forst. Bundesv. Berichte, Vienna.*
- Bonner, F.T., B.F. McLemore and J.P. Barnett. 1974.** Presowing treatment of seed to speed germination. Pp. 126–135 *In: Seeds of woody plants in the United States. C. S. Schopmeyer, Tech. Coord. USDA Forest Service Agricultural Handb. 450. Washington, DC.*
- Bramlett, D.L., E.W. Belcher, G.L. DeBarr, G.H. Hertel, R.P. Karrfalt, C.W. Lantz, T. Miller, K.D. Ware and H.O. Yates, III. 1977.** Cone analysis of southern pines, a guidebook. USDA Forest Service Gen. Technical Report SE-13. Southeastern Forest Experiment Station, Asheville, NC. 28 p.
- Carpita, N.C., A. Skaria, J.P. Barnett and J.R. Dunlap. 1983.** Cold stratification and growth of radicles of loblolly pine (*Pinus taeda*) embryos. *Physiologia Plantarum* 59: 601–606.
- Czabator, F.J. 1962.** Germination value: an index combining speed and completeness of pine seed germination. *For. Sci.* 8: 386–396.
- Dougherty, P.M. and M.L. Duryea. 1991.** Regeneration: an overview of past trends and basic steps needed to ensure future success. Pp. 3–7 *In: Forest Regeneration Manual, M.L. Duryea and P.M. Dougherty, Eds. Kluwer Academic Publishers, Dordrecht, The Netherlands.*
- Dunlap, J.R. and J.P. Barnett. 1983.** Influence of seed size on germination and early development of loblolly pine germinants. *Can. J. For. Res.* 13: 40–44.
- Fatzinger, C.W., H.O. Yates, III, W.J. Hammond and R. Hutto. 1985.** Insect caused cone and seed losses in treated and untreated loblolly pine seed orchards. USDA Forest Service Research Note SE-333. Southeastern Forest Experiment Station, Asheville, NC. 4 p.
- Jones, L. 1966.** Storing southern pine seed: what are the best moisture and temperature conditions? Georgia Forestry Research Council Res. Pap. 41. Macon, GA. 8 p.
- Kamra, S.K. 1967.** Studies on storage of mechanically damaged seed of Scots pine (*Pinus sylvestris* L.). *Studia Forestalia Suecica* 42. Royal College of Forestry, Stockholm, Sweden. 17 p.
- McConnell, J.L. and J.L. Edwards. 1985.** Net retrieval seed collection. Pp. 125–128 *In: Proceedings conifer tree seed in the Inland West symposium. USDA Forest Service Gen. Tech. Rep. INT-203. Intermountain Research Station, Ogden, UT.*
- McLemore, B.F. 1961.** Prolonged storage of longleaf cones weakens seed. USDA Forest Service Southern Forestry Notes 132. Southern Forest Experiment Station, New Orleans, LA.
- McLemore, B.F. 1969.** Long stratification hastens germination of loblolly pine seed at low temperatures. *J. For.* 67: 419–420.
- McLemore, B.F. 1975a.** Collection date, cone-storage period affect southern pine seed yields, viability. *Tree Planters Notes* 26(1): 24–26.
- McLemore, B.F. 1975b.** Cone and seed characteristics of fertilized and unfertilized longleaf pines. USDA Forest Service Research Paper SO-109. Southern Forest Experiment Station, New Orleans, LA. 10 p.
- Pait, J.A., III, D.M. Flinchum, and C.W. Lantz. 1991.** Species variation, allocation, and tree improvement. Pp. 207–231 *In: M.L. Duryea and P.M. Dougherty, Eds. Forest Regeneration Manual. Kluwer Academic Publishers, Dordrecht, Boston, London.*
- Perry, T.O. 1976.** Maternal effects on the early performance of tree progenies. Pp. 473–481 *In: M. G. R. Carnell and F. T. Last, Eds. Tree physiology and yield improvement. Academic Press, New York.*
- Rietz, R.C. 1941.** Kiln design and development of schedules for extracting seed from cones. U.S. Department of Agriculture Tech. Bull. 733. Washington, DC. 70 p.
- Silen, R. and C. Osterhans. 1979.** Reduction of genetic base by sizing of bulk Douglas-fir seed lots. *Tree Planters Notes* 30(1): 24–30.
- Tanaka, Y. 1984.** Assuring seed quality for seedling production: cone collection and seed processing, testing, storage, and stratification. Pp. 27–39 *In: Forest nursery manual: Production of bareroot seedlings, M. L. Duryea and T. D. Landis, Eds. Martinus Nijhoff/Dr. W. Junk Publishers, The Hague/Boston/Lancaster.*
- Venator, C.R. 1973.** The relationship between seedling height and date of germination in *Pinus caribaea* var. *hondurensis*. *Turrialba* 23: 473–474.
- Wakeley, P.C. 1954.** Planting the southern pines. U.S. Department of Agriculture Agric. Monograph 18. Washington, DC. 233 p.
- Wynens, J. and T. Brooks. 1979.** Seed collection from loblolly pine. Georgia Forestry Commission Res. Pap. 3. Macon, GA.
- Zoerb, M.H., Jr. 1969.** Clonal variation in time of cone ripening in loblolly pine. Union Camp Corp. Woodlands Res. Note 22. 3 p.