
Largest-Crown- Width Prediction Models for 53 Species in the Western United States

William A. Bechtold, *USDA Forest Service, Southern Research Station, P.O. Box 2680, Asheville, NC 28802.*

ABSTRACT: *The mean crown diameters of stand-grown trees 5.0-in. dbh and larger were modeled as a function of stem diameter, live-crown ratio, stand-level basal area, latitude, longitude, elevation, and Hopkins bioclimatic index for 53 tree species in the western United States. Stem diameter was statistically significant in all models, and a quadratic term for stem diameter was required for some species. Crown ratio and/or Hopkins index also improved the models for most species. A term for stand-level basal area was not generally needed but did yield some minor improvement for a few species. Coefficients of variation from the regression solutions ranged from 17 to 33%, and model R^2 ranged from 0.15 to 0.85. Simpler models, based solely on stem diameter, are also presented. *West. J. Appl. For.* 19(4):245–251.*

Key Words: Largest crown width, tree crown width, crown diameter, crown modeling.

Crown-width models are commonly separated into two categories—models for open-grown trees and models for stand-grown trees. The dimensions of crowns in open settings approach maximum biological potential, while those of stand-grown trees are generally smaller due to the influence of competition. Terminology developed by crown modelers in the western United States identifies models based on open-grown trees as “maximum crown width” (MCW) models, and those derived from stand-grown trees as “largest crown width” (LCW) models (Hann 1997). Both types of models relate to the horizontal silhouette of a crown as defined by the vertical projection of its longest branch tips, hence the terms “maximum” and “largest.” MCW and LCW approximate the mean diameter of this silhouette from field measurements of crown extension along two or more axes passing through the tree crown.

MCW models predict potential crown size and are primarily used to develop tree stocking guides (Smith and Gibbs 1970) and crown competition indices (Krajicek et al. 1961). LCW models predict the actual size of tree crowns in forest settings, resulting in a variety of applications that include estimations of crown surface area and volume (Zarnoch et al. 2004), forest canopy cover (Gill et al. 2000), tree-crown profiles (Hann 1999), and wildlife habitat indices (Hays et al. 1981).

LCW prediction models are appealing because the direct measurement of crown diameters in the field is costly,

particularly for extensive inventories. The measurement of mean crown diameter with a logger’s tape averages more than one minute per tree (Bechtold et al. 2002). The objective of this article is to use extensive tree- and stand-level data gathered by the USDA Forest Service Forest Health Monitoring program (FHM) in the western United States to develop regional LCW prediction models for as many tree species as possible. A similar study has recently been conducted for species endemic to the eastern United States (Bechtold 2004).

Previous Studies

Significant relationships between crown width and stem diameter are well established for open-grown and stand-grown trees of many species (Krajicek et al. 1961, Dawkins 1963, Hetherington 1967). Simple linear relationships between crown width and stem diameter are often adequate, but quadratic expressions of stem diameter are known to improve crown-width models for some species (Paine and Hann 1982). Although diameter at breast height (dbh) is the most common variable used in crown-width prediction models, LCW (and occasionally MCW) models have been supplemented with additional tree-level and stand-level variables. Moeur (1981) used total height and crown length in models for 11 species in the northern Rocky Mountains, as did Hann (1997) for 15 species in western Oregon. Bechtold et al. (2002) found vertical crown ratio to be significant in models for 13 tree species in North Carolina. Bragg (2001) improved crown-diameter models for 20 species in the upper lake states by adding a term for basal area competition. Crown width also has been shown to vary by geographic location. Paine and Hann (1982) improved

NOTE: William A. Bechtold can be reached at (X28) 257-4357; Fax: (828) 257-4894; wabechtold@fs.fed.us. The USDA Forest Service, Forest Health Monitoring Program, provided funding for this research. This study was conducted and written by US government employees and is therefore in the public domain.

crown-width models for 11 of 15 species in southwest Oregon with the introduction of coordinates relating trees to a geographic reference point. To summarize, these studies show that measures of vertical tree dimension, stand density, and geographic location can improve crown-width models for some species over the use of stem diameter alone.

Methods

The Data

Between 1992 and 1999, the FHM program established a network of 1/6-ac plots systematically distributed across eight western states (California, Colorado, Idaho, Nevada, Oregon, Utah, Washington, and Wyoming). In addition to crown diameters, a variety of other tree and stand parameters were measured for use as indicators of forest ecosystem productivity and sustainability. The FHM plot network has since been integrated with the Forest Inventory and Analysis (FIA) sampling grid (Stolte 2001). Between 1992 and 1999, some plots were remeasured multiple times during successive inventories. To avoid problems with autocorrelation, only the most recent measurement of each tree was used for this analysis. After deleting species with less than 25 observations and applying additional screening restrictions as discussed below, the FHM dataset yielded a total of 21,689 observations from 983 forested plots across 8 western states.

The crown diameters used for this analysis conform to the LCWs of stand-grown trees. To ensure that only stand-grown trees were included, those with an "open grown" crown class were deleted. For each sampled tree with a stem diameter of at least 5.0 in., field crews measured (with logger's tapes) the horizontal diameter of the widest axis of the crown, plus the dimension perpendicular to the widest axis. The arithmetic mean diameter calculated from these two field measurements is the dependent variable in the prediction equations that follow.

For most species, dbh was used as the independent variable for stem diameter. For species with shrub-like form designated as "woodland," diameter at root collar (drc) was substituted for dbh as the measure of stem diameter (USDA Forest Service 2002).

Live-crown ratio was investigated as a measure of vertical crown dimension potentially correlated with the crown diameters of species encountered in this study. Tree length, crown length, and height to crown base are similar variables used by other modelers but not available in the FHM dataset. The crown ratios used in this analysis adhere to the rules for "uncompacted" live crown ratio as specified by the USDA Forest Service (2002). The term "uncompacted" means that estimates of crown ratio were not reduced to compensate for gaps between the base of the live crown and the top of a tree.

Stand-level basal aredac was selected to quantify the effect of stand density on crown diameter. Basal areas were computed from all live tally trees with stem diameters ≥ 5.0 in. For woodland species, drc was substituted for dbh in the basal area calculations.

Latitude, longitude, and elevation are potentially useful for integrating the effect of geographic location. Because there is much interaction between these variables in the complex topography of the western United States, an index comprised of all three was identified as an additional candidate variable. Hopkins (1938) studied the phenologic occurrence of springtime and concluded that relative to a given geographic position, spring is delayed by 1 day for every 100 ft of elevation, 4 days for every 1 degree of northward latitude, and by 1 1/4 days for every 1 degree of westward longitude. Based on these relationships, Hopkins bioclimatic index (i.e., the number of days spring is delayed) was computed for each tree sampled relative to the mean elevation (5,549 ft), latitude (42.16 degrees), and longitude (-116.39 degrees) of all plots in the 8-state region:

$$HI = \left(\frac{(E - 5449)}{100} \right) 1 + (LAT - 42.16)4 + (-116.39 - LON)1.25 \quad (1)$$

where:

- E = elevation (ft);
- LAT = latitude (decimal degrees); and
- LON = longitude (decimal degrees).

A positive H value means that spring is delayed relative to the reference position, while a negative value indicates that spring is advanced.

Regression Models

Mean LCW (in ft) was modeled as a function of one or more of the following terms associated with stem diameter, vertical crown dimension, stand density, and geographic location:

- D = dbh (in.), or drc (in.) for woodland species;
- CR = live crown ratio (%);
- BA = stand-level basal area (ft²/ac);
- LAT = latitude (decimal degrees);
- LON = longitude (decimal degrees);
- E = elevation (ft); and
- HI = Hopkins index (days).

The candidate variables then were evaluated with a series of fixed and stepwise regressions designed to identify the best model for each species. The ranges of the variables used in the final models resulting from the regression analyses are provided in Table 1.

Results and Discussion

Stem diameter and crown diameter are known to be highly correlated, so stem diameter was entered first into the ordinary least squares (OLS) regression:

$$LCW = b_0 + h_1(D), \quad (2)$$

where b_0 and b_1 are regression parameters estimated from the data.

Table 1. Ranges of data used to fit crown-width prediction models for 53 species in the western United States.

| Species ^a | n | Crown width ^b (ft) | | Stem diameter ^c (in.) | | Crown ratio (%) | | Basal area ^d (ft ² /ac) | | Latitude (decimal deg) | | Longitude (decimal deg) | | Elevation (ft) | | Hopkins index ^d (days) | | | |
|--------------------------|------|-------------------------------|-----|----------------------------------|------|-----------------|-----|---|-----|------------------------|-------|-------------------------|---------|----------------|-------|-----------------------------------|------|-----|----|
| | | Min | Max | Min | Max | Min | Max | Min | Max | Min | Max | Min | Max | Min | Max | Min | Max | | |
| Softwood species | | | | | | | | | | | | | | | | | | | |
| Pacific silver fir | 218 | 4 | 33 | 5.0 | 35.8 | 5 | 99 | 2 | 493 | 43.16 | 4X.96 | -1 | 23.95 | -120.52 | 2100 | 5 300 | -9 | 26 | |
| White fir | 855 | 3 | 35 | 5.0 | 62.6 | 10 | Y9 | 4 | 445 | 34.08 | 46.1 | -1 | 23.54 | -104.95 | 322 | 10200 | -40 | 19 | |
| Grand fir | 610 | 4 | 35 | 5.0 | 43.9 | 5 | 99 | 12 | 496 | 40.54 | 48.91 | -124.33 | -114.58 | 295 | 6400 | -48 | 20 | | |
| Corkbark fir | 68 | 5 | 15 | 5.2 | 18.3 | 45 | 99 | 70 | 222 | 37.02 | 41.44 | -1 | 12.17 | -105.15 | 8200 | 11200 | 1 | 28 | |
| Subalpine fir | 1262 | 3 | 2X | 5.0 | 27.4 | 5 | 99 | 3 | 280 | 37.44 | 48.86 | -123.05 | -105.67 | 1600 | 11500 | -14 | 44 | | |
| California red fir | 160 | 4 | 36 | 5.0 | 52.3 | 20 | 99 | 35 | 423 | 33.72 | 41.2X | -123.46 | -1 | 16.67 | 4600 | 9000 | -35 | 26 | |
| Shasta red fir | 63 | 6 | 26 | 5.0 | 40.1 | 40 | 99 | 38 | 28X | 41.47 | 43.70 | -123.05 | -123.05 | 5500 | 6600 | 3 | 17 | | |
| Noble fir | 50 | 4 | 29 | 5.2 | 46.0 | 20 | 99 | X | 5 | 354 | 43.94 | 48.86 | 62.35 | -1 | 19.96 | 2100 | 5600 | -1 | 32 |
| Port-Orford cedar | 7X | 3 | 22 | 5.0 | 13.4 | 15 | 99 | 199 | 270 | 42.85 | 43.15 | -124.46 | -123.88 | 800 | 2800 | -35 | -15 | | |
| California juniper (w) | 28 | 6 | 31 | 5.2 | 42.6 | 35 | 99 | 47 | 101 | 34.X2 | 41.56 | -121.03 | -1 | 18.28 | 4000 | 5837 | -41 | 4 | |
| Western juniper (w) | 302 | 3 | 36 | 5.0 | 45.3 | 10 | 99 | 8 | 218 | 36.35 | 44.93 | -121.91 | -1 | 13.01 | 200 | 9000 | -71 | 27 | |
| Utah juniper (w) | 402 | 1 | 30 | 5.0 | 29.7 | 10 | 99 | 8 | 221 | 34.32 | 44.93 | -121.91 | -104.86 | 4623 | X600 | -30 | 27 | | |
| Rocky mtn. juniper (w) | 144 | 5 | 29 | 5.0 | 38.1 | 5 | 99 | 5 | 277 | 37.02 | 44.91 | -1 | 12.46 | -103.60 | 3900 | X725 | -37 | 19 | |
| One-seed juniper (w) | 98 | 2 | 2X | 5.7 | 36.5 | 10 | 99 | 11 | 512 | 37.13 | 38.53 | -105.09 | -103.1 | 4900 | 6700 | -43 | -17 | | |
| Western larch | 183 | 3 | 2X | 5.0 | 23.1 | 5 | 99 | 24 | 300 | 44.42 | 4X.97 | -121.20 | -1 | 14.92 | 295 | 6400 | -25 | 20 | |
| Incense cedar | 220 | 3 | 32 | 5.0 | 42.0 | 5 | 99 | 12 | 445 | 35.72 | 44.71 | -123.79 | -11X.33 | 1100 | 6800 | -33 | 11 | | |
| Engelman spruce | 1205 | 3 | 29 | 5.0 | 35.9 | 5 | 99 | 2 | 391 | 37.02 | 48.89 | -123.84 | -105.15 | 295 | 12000 | -25 | 44 | | |
| Sitka spruce | 53 | 7 | 43 | 5.1 | 39.2 | 20 | 99 | 12 | 307 | 40.54 | 47.77 | -124.46 | -123.30 | 0 | 2X00 | -46 | -5 | | |
| Whitebark pine | 97 | 4 | 29 | 5.0 | 24.2 | 10 | 99 | 46 | 270 | 37.54 | 48.62 | -123.05 | -110.01 | 6200 | 10700 | 6 | 44 | | |
| Bristlecone pine | 26 | 7 | 25 | 5.0 | 18.9 | 65 | 99 | 24 | 121 | 37.31 | 39.45 | -114.65 | -105.06 | 8410 | 10700 | 9 | 25 | | |
| Common pinyon (w) | 27X | 5 | 27 | 5.0 | 25.5 | 30 | 99 | 3 | 500 | 35.12 | 40.14 | -118.28 | -103.32 | 4600 | 8725 | -40 | 11 | | |
| Lodgepole pine | 2761 | 1 | 40 | 5.0 | 62.2 | 5 | 99 | 2 | 315 | 36.35 | 4X.90 | -124.18 | -105.41 | 200 | 10900 | -23 | 43 | | |
| Limber pine | 164 | 3 | 34 | 5.0 | 20.7 | 5 | 99 | 3 | 256 | 37.02 | 45.64 | -1 | 15.76 | -105.15 | 6400 | 11200 | 2 | 43 | |
| Jeffrey pine | 108 | 5 | 44 | 5.1 | 48.1 | 5 | 99 | 12 | 345 | 36.10 | 42.19 | -123.75 | -1 | 18.33 | 1700 | 9000 | -38 | 15 | |
| Sugar pine | 118 | 5 | 49 | 5.0 | 44.9 | 15 | 99 | 2X | 445 | 34.08 | 42.73 | -124.10 | -1 | 16.94 | 1100 | X200 | -47 | 11 | |
| Western white pine | 84 | 4 | 34 | 5.1 | 3X.4 | 15 | 99 | 15 | 423 | 36.21 | 48.92 | -124.18 | -115.87 | 200 | 9200 | -25 | 32 | | |
| Ponderosa pine | 1413 | 1 | 46 | 5.0 | 41.3 | 5 | 99 | 4 | 414 | 36.81 | 4X.91 | -123.57 | -104.07 | 300 | 9800 | -56 | 41 | | |
| Grey pine | 37 | 6 | 54 | 5.1 | 34.X | 30 | 95 | 9 | 164 | 35.70 | 41.18 | = | 122.90 | -1 | 1X.05 | 400 | 4900 | -69 | -4 |
| Singleleaf pinyon (w) | 323 | 4 | 30 | 5.0 | 22.1 | 20 | 99 | 7 | 206 | 34.32 | 40.26 | -1 | 19.60 | -1 | 12.16 | 200 | 8900 | -71 | 15 |
| Douglas-fir | 4088 | 1 | 66 | 5.0 | 68.7 | 5 | 99 | 2 | 696 | 37.13 | 48.97 | -124.46 | -104.95 | 100 | 11000 | -49 | 67 | | |
| Redwood | 55 | 3 | 31 | 5.1 | 35.6 | 5 | 99 | 104 | 353 | 37.07 | 40.68 | = | 124.07 | -122.17 | 400 | 1800 | -55 | -39 | |
| Western redcedar | 439 | 3 | 38 | 5.0 | 62.0 | 15 | 99 | 13 | 493 | 37.90 | 48.97 | = | 124.42 | -109.76 | 200 | 8520 | -25 | 49 | |
| Western hemlock | 1008 | 4 | 54 | 5.0 | 63.5 | 5 | 99 | 12 | 496 | 42.6X | 4X.96 | = | 124.42 | -1 | 15.93 | 0 | xsoo | -34 | 49 |
| Mountain hemlock | 200 | 1 | 33 | 5.0 | 32.4 | 5 | YY | 28 | 397 | 37.64 | 4X.72 | -123.55 | -1 | 15.23 | 1600 | 9200 | -10 | 27 | |
| Hardwood species | | | | | | | | | | | | | | | | | | | |
| Bigleaf maple | 106 | 8 | 57 | 5.1 | 34.1 | 10 | YY | 37 | 405 | 39.82 | 48.33 | -124.06 | -121.38 | 100 | X500 | -36 | 49 | | |
| Rocky mtn. maple (w) | 70 | 8 | 39 | 5.0 | 26.2 | 25 | 99 | 10 | 2X8 | 40.75 | 48.64 | -123.48 | -1 | 10.95 | 1600 | 7200 | -3X | 14 | |
| Bigtooth maple (w) | 4X | 4 | 20 | 5.0 | 14.7 | 30 | YY | 4 | 145 | 37.34 | 41.13 | -112.75 | -11 | 1.33 | 5400 | 8275 | -11 | 11 | |
| Red alder | 409 | 3 | 54 | 5.0 | 28.3 | 10 | 99 | 6 | 386 | 40.54 | 4X.82 | -124.46 | -1 | 17.30 | 0 | 8100 | -46 | 53 | |
| White alder | 37 | 8 | 35 | 5.1 | 18.1 | 25 | 90 | 49 | 270 | 38.75 | 45.61 | -123.92 | -120.98 | 700 | 3000 | -56 | -2 | | |
| Pacific madrone | 164 | 1 | 43 | 5.0 | 29.7 | 5 | 99 | 13 | 369 | 37.21 | 45.33 | -124.20 | -121.06 | 700 | 6000 | -55 | 15 | | |
| Curleaf mtn-mahogany (w) | 227 | 3 | 29 | 5.0 | 24.0 | 10 | YY | 10 | 174 | 34.32 | 45.05 | -121.91 | -109.70 | 495 | 9200 | -37 | 27 | | |
| Tanoak | 534 | 2 | 41 | 5.0 | 31.0 | 5 | 99 | 57 | 353 | 37.07 | 42.52 | -124.37 | -121.06 | 300 | 6000 | -55 | 15 | | |
| Quaking aspen | 1383 | 1 | 34 | 5.0 | 19.9 | 5 | 99 | 2 | 256 | 37.02 | 4X.42 | -1 | 19.27 | -104.07 | 3200 | 10X00 | -9 | 33 | |
| Narrowleaf cottonwood | 44 | 8 | 35 | 5.1 | 16.6 | 30 | 90 | 17 | 150 | 37.05 | 38.85 | -10X.65 | -106.15 | 5400 | 8000 | -26 | -2 | | |
| Coastal live oak | 87 | 3 | 53 | 5.0 | 40.6 | 5 | YY | 60 | 147 | 34.51 | 3X.62 | -122.63 | -1 | 19.81 | 500 | 2000 | -73 | -54 | |
| Canyon live oak | 440 | 5 | 49 | 5.0 | 53.2 | 5 | 99 | 10 | 696 | 34.08 | 42.43 | -124.20 | -1 | 16.94 | 700 | 8200 | -60 | -5 | |
| Blue oak | 184 | 5 | 61 | 5.0 | 29.4 | 15 | 99 | 4 | 172 | 35.23 | 40.75 | -1 | 22.99 | -1 | 18.55 | 400 | 5900 | -69 | -2 |
| Gambel oak (w) | 24X | 1 | 10 | 5.0 | 15.4 | 5 | 99 | Y | 149 | 37.02 | 40.16 | -1 | 13.28 | -104.73 | 6000 | 8500 | -23 | 11 | |
| Oregon white oak | 126 | 6 | 30 | 5.0 | 22.4 | 15 | YY | 22 | 200 | 38.3X | 45.87 | -123.34 | -120.75 | 900 | 3700 | -54 | -11 | | |
| California black oak | 239 | 4 | 52 | 5.1 | 40.3 | 5 | 99 | 20 | 345 | 35.37 | 42.63 | -123.65 | -1 | 18.29 | 1100 | 6200 | -47 | -8 | |
| Valley oak | 29 | 5 | 47 | 5.1 | 21.3 | 10 | 99 | 63 | 172 | 35.70 | 39.20 | -122.99 | -120.96 | 900 | 1800 | -64 | -44 | | |
| Interior live oak | 79 | 2 | 37 | 5.0 | 19.5 | 15 | YY | 10 | 77 | 35.99 | 40.70 | -123.21 | -1 | 1X.06 | 600 | 7300 | -60 | -5 | |
| California laurel | 2X | 10 | 44 | 5.1 | 1X.7 | 30 | 99 | 24 | 271 | 3X.31 | 43.63 | -124.06 | -120.30 | 900 | 6000 | -54 | 15 | | |

^a Species designated (w) are woodland species where stem diameter is diameter at rootcollar (drc) instead of diameter at breast height (dbh)
^b Largest crown width of stand-grown trees (i.e., mean crown diameter).
^c Stand-level basal area (trees with stem diameters 5.0-in. and larger).
^d Hopkins index is the number of days spring is delayed relative to latitude 42.16°, longitude -116.39°, and elevation 5449 ft.

Examination of the residuals from the regression solutions indicated heteroscedasticity with respect to *D* for many species. A weighted least squares (WLS) approach thus was used for this and subsequent regressions to counter the effect of increasing variation with increasing stem diameter. Appropriate weights were determined by modeling

the variance of the residuals from the OLS solutions as a function of *D*, as follows:

1. The LCW models were solved using OLS regression.
2. The absolute values of the OLS residuals (*K*) were modeled as a function of *D*:

$$|R| = b_0 + b_1(D).$$

3. The LCW models were solved again using WLS, where each observation was weighted by w_i , the reciprocal of the estimated variance with respect to D_i :

$$w_i = 1/(b_0 + b_1(D_i))^2,$$

where $i = 1 \dots n$.

The D coefficients for pacific yew (*Taxus brevifolia*) and golden chinkapin (*Castanopsis chrysophylla*) were not statistically significant at a P value of 0.05. Because the ability to develop biologically justifiable models was doubtful, and the numbers of observations were limited (30 each), these two species were deleted from the analysis. The number of species available for modeling thus was reduced to 53 from a previous total of 55.

Further examination of the residuals from Equation 2 indicated that a quadratic term might improve the models for some species. All species were thus re-fitted with the model

$$LCW = b_0 + b_1(D) + b_2(D^2), \quad (3)$$

using WLS regression, and the quadratic term was retained for 14 species where the P -value associated with the D^2 coefficient was significant at $P = 0.05$.

On fixing D and D^2 in those models where significant (i.e., retaining these terms in subsequent regressions), all models then were re-fitted with an additional term for crown ratio (CR):

$$LCW = b_0 + b_1(D) + b_2(D^2) + b_3(CR). \quad (4)$$

CR then was retained for 39 species where its coefficient was significant ($P = 0.05$). At this stage, the signs of all coefficients were consistent and biologically reasonable. The coefficients associated with D were all positive, confirming a positive correlation between stem diameter and crown diameter. The coefficients associated with D^2 all were negative, meaning that crown diameter approaches an upper biological limit as stem diameter increases. The coefficients associated with CR all were positive, indicating that large crowns tend to be large in all dimensions.

After fixing D , D^2 , and CR in models where these terms were significant, all models were then re-fitted with an additional term for stand-level basal area (BA):

$$LCW = b_0 + b_1(D) + b_2(D^2) + b_3(CR) + b_4(BA). \quad (5)$$

The BA term was statistically significant ($P = 0.05$) in models for 19 species. A negative correlation between stand density and crown diameter was expected, but the additional term exhibited a mixture of positive and negative coefficients. In the few models with BA coefficients that were negative and statistically significant, the partial R^2 values resulting from the addition of BA generally were less than 0.02. Because D and CR are tree-level variables highly correlated with stand density, the general instability and weak significance of the BA term was attributed to col-

linearity with D and CR . As a result, the utility of the BA term in any of these models is questionable, but it did yield minor improvement for a few species. The BA term thus was retained for nine species where it was statistically significant and the coefficient was negative.

Again after fixing D , D^2 , CR , and BA in models where significant, all models were re-fitted with stepwise regressions where additional terms for LAT , LON , and E were entered as candidates. The stepwise procedure selected one or two of these geographic variables as statistically significant for many species, but there was no clear consistency. Different geographic terms were selected for different species, coefficient signs fluctuated between positive and negative for a given geographic variable, and some of the model intercepts changed dramatically. Over-parameterization, as well as interactions among latitude, longitude, and elevation made it impractical to include up to three different terms for geographic location, so Hopkins' (1938) bioclimatic index was investigated as an alternative.

Because Hopkins index was developed in the eastern United States, its applicability to western states is uncertain. The relationship between latitude, elevation, and climatic condition seemed reasonable for the West, but the negative effect of westward longitude was suspect-possibly attributable to distance from the moderating climatic effect of the Atlantic Ocean in the region where the index was developed. Theorizing that crown diameters generally should be smaller under climatic conditions where spring is delayed, correlations between LCW and latitude, longitude, and elevation were checked to verify a negative correlation between LCW and elevation, a negative correlation between LCW and latitude, and a positive correlation between LCW and longitude. Among individual species, there was considerable fluctuation in the signs of the correlation coefficients for each of these variables (again attributed to latitude, longitude and elevation interactions); but when averaged across all 53 species, the means of the correlation coefficients exhibited signs that were consistent with Hopkins (193X).

With D , D^2 , CR , and BA fixed in models where significant, all models were then re-fitted with an additional term for Hopkins index (HI)

$$LCW = b_0 + b_1(D) + b_2(D^2) + b_3(CR) + b_4(BA) + b_5(HI), \quad (6)$$

and HI was retained for 31 species for which its coefficient was significant ($P = 0.05$). Most of the coefficients associated with HI were negative, confirming that tree crowns generally are smaller in harsher climates where spring is delayed, but a few of the coefficients were positive. This was attributed to the possibility that competing species may drop out of the stand-level species mix as climatic conditions become more extreme.

Equation 6 thus was chosen as the best biologically justifiable model attainable from the available data, with

Table 2. Model statistics and parameter estimates from crown-width prediction Equation 6, for 53 species in the western United States.

| species ^a | Equation 6 ^b : $LCW = b_0 + b_1(D) + b_2(D^2) + b_3(CR) + b_4(BA) + b_5(HI)$ | | | | | | | | |
|--------------------------|---|------|----|----------------------------------|---------|---------|---------|---------|----------|
| | Model statistics ^c | | | Parameter estimates ^d | | | | | |
| | RSQ | RMSE | CV | b_0 | b_1 | b_2 | b_3 | b_4 | b_5 |
| Softwood species | | | | | | | | | |
| Pacific silver fir | 0.43 | 3.8 | 2X | 7.7763 | 0.5960 | — | — | — | -0.0705 |
| White fir | 0.62 | 2.7 | 21 | 2.47X9 | 0.93 17 | -0.0128 | 0.0327 | — | -0.1178 |
| Grand fir | 0.58 | 3.3 | 22 | 3.0335 | 0.9752 | -0.0113 | 0.0548 | — | -0.0597 |
| Corkbark fir | 0.15 | 2.1 | 24 | 6.0730 | 0.3756 | — | — | — | — |
| Subalpine fir | 0.52 | 2.3 | 24 | 2.6068 | 0.6145 | — | 0.0417 | — | -0.0698 |
| California red fir | 0.78 | 2.6 | 21 | 2.3660 | 0.5472 | — | 0.0316 | — | -0.0702 |
| Shasta red fir | 0.83 | 2.2 | 19 | 4.0524 | 0.6423 | — | — | — | — |
| Noble fir | 0.72 | 3.0 | 20 | 2.7761 | 0.7311 | — | 0.0476 | — | -0.0756 |
| Port-Orford cedar | 0.38 | 2.6 | 25 | 1.0365 | 0.7943 | — | 0.0399 | — | — |
| California juniper (w) | 0.76 | 3.2 | 22 | -0.6303 | 1.6960 | -0.0225 | — | — | 0.1 166 |
| Western juniper (w) | 0.65 | 3.x | 2x | -0.0037 | 1.3526 | -0.0165 | — | — | — |
| Utah juniper (w) | 0.50 | 3.0 | 31 | -5.9542 | 1.1877 | -0.0256 | 0.0857 | -0.0095 | — |
| Rocky mtn. juniper (w) | 0.60 | 3.1 | 25 | 4.1599 | 1.3528 | -0.0233 | 0.0633 | — | -0.0423 |
| One-seed juniper (w) | 0.70 | 2.7 | 23 | -0.7915 | 0.5975 | — | 0.0890 | -0.0094 | — |
| Western larch | 0.36 | 3.8 | 31 | 1.5995 | 0.7675 | — | 0.0750 | — | — |
| Incense cedar | 0.67 | 2.x | 22 | 2.0872 | 0.92X 1 | -0.0094 | 0.0332 | — | — |
| Engelman spruce | 0.55 | 2.5 | 23 | 4.1348 | 0.5694 | — | 0.0403 | — | -0.1014 |
| Sitka spruce | 0.65 | 4.7 | 24 | 8.8087 | 0.7825 | — | — | — | — |
| Whitebark pine | 0.53 | 3.2 | 31 | 0.5223 | 0.7432 | — | — | — | 0.0829 |
| Bristlecone pine | 0.58 | 3.1 | 20 | 12.7069 | 0.9571 | — | 0.2177 | — | — |
| Common pinyon (w) | 0.67 | 2.6 | 21 | -5.4647 | 1.9660 | -0.0395 | 0.0427 | — | -0.0259 |
| Lodgepole pine | 0.61 | 2.7 | 28 | -1.5440 | 1.3828 | -0.0200 | 0.0396 | -0.0083 | — |
| Limber pine | 0.56 | 3.3 | 28 | 3.4094 | 0.8638 | — | 0.0592 | -0.0105 | -0.0956 |
| Jeffrey pine | 0.85 | 2.7 | 19 | 1.2784 | 0.7937 | — | 0.0334 | — | -0.0887 |
| Sugar pine | 0.77 | 3.7 | 25 | 3.1052 | 0.8049 | — | — | — | -0.1230 |
| Western white pine | 0.74 | 3.2 | 25 | 4.X643 | 0.6949 | — | — | — | -0.0974 |
| Ponderosa pine | 0.73 | 3.0 | 23 | -0.3459 | 1.1110 | -0.0080 | 0.0566 | -0.0094 | -0.0362 |
| Grey pine | 0.81 | 4.3 | 20 | -2.4909 | 1.0716 | — | 0.0648 | — | -0.1127 |
| Singleleaf pinyon (w) | 0.66 | 2.4 | 23 | 2.1556 | 0.8302 | — | 0.0299 | -0.0195 | -0.0272 |
| Douglas-fir | 0.57 | 4.2 | 26 | 3.2346 | 1.1158 | -0.0112 | 0.0442 | -0.0057 | -0.0237 |
| Redwood | 0.46 | 4.4 | 23 | -15.1653 | 0.3182 | — | 0.1349 | — | -0.4682 |
| Western redcedar | 0.37 | 4.2 | 25 | 5.2911 | 1.0612 | -0.0153 | 0.0469 | — | — |
| Western hemlock | 0.63 | 4.4 | 26 | -0.4624 | 1.0429 | -0.0078 | 0.1018 | — | -0.027 1 |
| Mountain hemlock | 0.60 | 3.3 | 29 | -0.3362 | 0.7142 | — | 0.04 14 | — | — |
| Hardwood species | | | | | | | | | |
| Bigleaf maple | 0.45 | 7.5 | 33 | -1.9386 | 1.2250 | — | 0.1622 | — | -0.1417 |
| Rocky mountain maple (w) | 0.34 | 5.0 | 2X | 5.9765 | 0.8648 | — | 0.0675 | — | — |
| Bigtooth maple (w) | 0.3 1 | 2.6 | 23 | 4.0040 | 1.0604 | — | — | — | — |
| Red alder | 0.60 | 4.7 | 2X | -0.7294 | 1.2885 | — | 0.1307 | — | — |
| White alder | 0.38 | 4.8 | 24 | 4.61 XX | 0.9 135 | — | 0.1019 | — | — |
| Pacific madrone | 0.44 | 4X | 33 | 4.9133 | 0.9459 | — | 0.061 1 | — | 0.0523 |
| Curleaf mtn-mahogany (w) | 0.48 | 2.x | 26 | 4.0 1 05 | 0.861 1 | — | — | — | -0.043 1 |
| Tanoak | 0.52 | 3.9 | 27 | 3. 11 50 | 0.7966 | — | 0.0745 | -0.0053 | -0.0289 |
| Quaking aspen | 0.59 | 2.5 | 21 | -0.5095 | 1.2318 | — | 0.0744 | — | 0.0233 |
| Narrowleaf cottonwood | 0.52 | 4.0 | 24 | 4.1687 | 1.5355 | — | — | — | 0.1275 |
| Coastal live oak | 0.76 | 5.6 | 27 | -16.1696 | 1.7456 | — | 0.0925 | — | -0.1956 |
| Canyon live oak | 0.45 | 4.0 | 2X | 0.2738 | 1.0534 | — | 0.0350 | — | -0.1385 |
| Blue oak | 0.76 | 4.5 | 23 | 2.71 10 | 1.5159 | — | 0.04 15 | -0.0271 | — |
| Gambel oak (w) | 0.32 | 2.5 | 27 | 0.3309 | 0.8918 | — | 0.05 10 | — | — |
| Oregon white oak | 0.42 | 3.7 | 24 | -1.3160 | 2.93 11 | -0.0866 | — | — | — |
| California black oak | 0.47 | 5.7 | 33 | 1.6306 | 0.9867 | — | 0.0556 | — | -0.1199 |
| Valley oak | 0.78 | 3.7 | 17 | -2.1068 | 1.9385 | — | 0.0860 | — | — |
| Interior live oak | 0.56 | 4.4 | 24 | 0.7146 | 1.5460 | — | — | — | -0.1121 |
| California laurel | 0.58 | 4.7 | 24 | 2.4247 | 1.3174 | — | 0.0786 | — | — |

^a LCW = largest crown width of stand-grown trees (i.e., mean crown diameter); D = dbh for non-woodland species, or diameter at rootcollar for woodland species; CR = uncompacted vertical crown ratio (percent); BA = stand-level basal area, trees with stem diameters 5.0-in. and larger (ft²/acre); and HI = Hopkins Index (days spring is delayed relative to latitude 42.16° longitude -116.39°, and elevation 544 ft).

^b Species designated (w) are woodland species where stem diameter (D) is diameter at rootcollar (drc).

^c RSQ = adjusted R²; RMSE = root mean squared error from the regression solutions; CV = coefficient of variation from the regression solutions; CV = (RMSE/mean LCW)*100, where mean LCW = mean crown diameter.

^d Terms where parameter estimates are missing were excluded from the regression solutions due to nonsignificance. Nonsignificant model intercepts were retained in the solutions.

terms included or excluded for various species on the empirical basis of whether or not their associated coefficients were significant (P = 0.05).

Fit statistics and parameter estimates from the solutions of Equation 6 are presented in Table 2. Parameter estimates with missing values were excluded from the regressions for

Table 3. Model statistics and parameter estimates from crown-width prediction Equation 3, for 53 species in the western United States.

| Species ^a | Equation 3 ^b : $LCW = b_0 + b_1(D) + b_2(D^2)$ | | | | | |
|---------------------------|---|------|----|----------------------------------|----------|----------|
| | Model statistics ^c | | | Parameter estimates ^d | | |
| | RSQ | RMSE | CV | b_0 | b_1 | b_2 |
| Softwood species | | | | | | |
| Pacific silver fir | 0.42 | 3.X | 2X | 7.3037 | 0.5909 | |
| White fir | 0.51 | 3.0 | 24 | 4.4965 | 0.923X | -0.0120 |
| Grand fir | 0.50 | 3.6 | 23 | 5.7545 | 1.1196 | -0.0147 |
| Corkbark fir | 0.15 | 2.1 | 24 | 6.0730 | 0.3756 | |
| Subalpine fir | 0.44 | 2.5 | 26 | 3.9629 | 0.6469 | |
| California red fir | 0.75 | 2.x | 23 | 4.7623 | 0.5222 | |
| Shasta red fir | 0.83 | 2.2 | 19 | 4.0524 | 0.6423 | |
| Noble fir | 0.67 | 3.3 | 21 | 6.3260 | 0.6588 | |
| Port-Orford cedar | 0.33 | 2.7 | 26 | 2.3625 | 0.9974 | - |
| California juniper (w) | 0.68 | 3.7 | 25 | -2.1213 | 1.7308 | -0.0243 |
| Western juniper (w) | 0.65 | 3.x | 2X | -0.0037 | 1.3526 | -0.0165 |
| Utah juniper (w) | 0.24 | 3.6 | 3X | 2.4349 | 0.9000 | -0.0171 |
| Rocky mtn. juniper (w) | 0.54 | 3.3 | 27 | 2.1431 | 1.3447 | -0.0228 |
| One-seed juniper (w) | 0.46 | 3.6 | 31 | 5.7367 | 0.4932 | |
| Western larch | 0.30 | 4.0 | 33 | 4.5 176 | 0.793 1 | |
| Incense cedar | 0.65 | 2.9 | 23 | 4.1207 | 0.9773 | -0.0107 |
| Engelman spruce | 0.47 | 2.x | 25 | 5.1218 | 0.5547 | |
| Sitka spruce | 0.65 | 4.7 | 24 | X.8087 | 0.7825 | |
| Whitebark pine | 0.53 | 3.2 | 31 | 2.653 1 | 0.8015 | |
| Bristlecone pine | 0.4 1 | 3.7 | 24 | 7.425 1 | 0.8991 | |
| Common pinyon (w) | 0.65 | 2.7 | 21 | -1.2638 | 1.9922 | -0.04 10 |
| Lodgepole pine | 0.54 | 2.9 | 30 | -1.1994 | 1.5154 | -0.0232 |
| Limber pine | 0.35 | 4.0 | 34 | 4.0181 | 0.8528 | |
| Jeffrey pine | 0.83 | 2.9 | 20 | 4.2675 | 0.77 14 | |
| Sugar pine | 0.72 | 4.1 | 27 | 4.X657 | 0.7890 | |
| Western white pine | 0.69 | 3.5 | 2x | 4.2840 | 0.6949 | |
| Ponderosa pine | 0.66 | 3.3 | 25 | 2.3081) | 1.1388 | -0.0089 |
| Grey pine | 0.7 1 | 5.2 | 25 | 4.3699 | 1.2524 | |
| Singleleaf pinyon (w) | 0.55 | 2.X | 26 | 2.5093 | 0.8503 | |
| Douglas-fir | 0.51 | 4.4 | 27 | 5.7753 | 1.0639 | -0.0 109 |
| Redwood | 0.27 | 5.1 | 27 | 12.0128 | 0.4576 | |
| Western redcedar | 0.33 | 4.3 | 26 | x.1993 | 1.1134 | -0.0165 |
| Western hemlock | 0.50 | 5.1 | 31 | 5.0036 | 1.1808 | -0.0 107 |
| Mountain hemlock | 0.58 | 3.3 | 29 | 3.2343 | 0.6927 | |
| Hardwood species | | | | | | |
| Bigleaf maple | 0.27 | X.6 | 38 | 10.0915 | 1.1139 | |
| Rocky mountain maple (w) | 0.2X | 5.2 | 29 | 10.5451 | 0.9493 | |
| Bigtooth maple (w) | 0.31 | 2.6 | 23 | 4.0040 | 1.0604 | |
| Red alder | 0.47 | 5.5 | 32 | 4.7027 | 1.3537 | |
| White alder | 0.26 | 5.2 | 26 | 9.7927 | 0.9006 | |
| Pacific madrone | 0.40 | 5.0 | 34 | 5.7785 | 0.9832 | |
| Curlleaf mtn-mahogany (w) | 0.46 | 2.9 | 27 | 3.5082 | 0.8770 | |
| Tanoak | 0.40 | 4.4 | 31 | 6.7864 | 0.8443 | |
| Quaking aspen | 0.50 | 2.x | 24 | 2.5515 | 1.2029 | |
| Narrowleaf cottonwood | 0.47 | 4.2 | 25 | 2.8848 | 1.5866 | |
| Coastal live oak | 0.74 | 5.x | 2X | 0.5740 | 1.8475 | |
| Canyon live oak | 0.35 | 4.4 | 30 | 6.1397 | 1.0109 | |
| Blue oak | 0.76 | 4.5 | 23 | 3.9281 | 1.5550 | |
| Gambel oak (w) | 0.23 | 2.7 | 2x | 3.0334 | 0.9834 | |
| Oregon white oak | 0.42 | 3.7 | 24 | -1.3160 | 2.93 1 1 | -0.0866 |
| California black oak | 0.38 | 6.2 | 36 | 7.0284 | 1.0470 | |
| Valley oak | 0.77 | 3.x | 17 | 2.9954 | 1.9137 | |
| Interior live oak | 0.52 | 4.6 | 25 | 5.1005 | 1.6359 | |
| California laurel | 0.47 | 5.2 | 27 | 7.3204 | 1.4420 | |

^a LCW = largest crown width of stand-grown trees (i.e., mean crown diameter); D = dbh for non-woodland species, or diameter at rootcollar for woodland species.

^b Species designated (w) are woodland species where stem diameter (D) is diameter at rootcollar (drc).

^c RSQ = adjusted R²; RMSE = root mean squared error from the regression solutions; CV = coefficient of variation from the regression solutions; CV = (RMSE/mean LCW*100).

^d Terms where parameter estimates are missing were excluded from the regression solutions due to nonsignificance. Nonsignificant model intercepts were retained in the solutions.

species where they were determined to be nonsignificant. Although some of the model intercepts were also nonsignificant, these were retained to ensure that the resulting models were BLUE (best linear unbiased estimators).

The root mean squared errors (RMSE) shown in Table 2 provide estimates of the error in crown-diameter predictions in terms of feet. RMSE, a common measure of model performance, is most useful for comparing similar models

for similar species among different studies. Comparisons among models involving dissimilar species are facilitated with the coefficient of variation (CV), which re-expresses RMSE as a percentage of the mean of the dependent variable:

$$CV = \frac{RMSE}{LCW} \times 100, \quad (7)$$

where

RMSE = the root mean squared error from the regression solution; and

LCW = mean LCW from the model predictions.

Comparisons of model performance among species has utility for applications such as the FIA program, which occasionally debates the merits of measuring versus modeling various inventory attributes. Species with models that yield low CVs and other satisfactory diagnostic statistics might be identified as candidates where model predictions can be used in lieu of field measurements. By species, coefficients of variation from the regression solutions ranged from 17 to 33%. The mean CV across all 53 species was 25%. Little difference was observed in the proportion of variation captured by the models among hardwood, softwood, woodland and nonwoodland species groups. The mean CV for each of these groups was about 25%.

Model R^2 values from the solution of Equation 6 ranged from 0.15 for corkbark fir (*Abies lasiocarpa* var. *arizonica*) to 0.85 for jeffrey pine (*Pinus jeffreyi*). Mean R^2 values across all species resulting from regression solutions of Equations 3, 4, 5, and 6 were 0.50, 0.55, 0.56, and 0.57, respectively. This suggests that the addition of crown ratio (to species where it was found to be statistically significant) increases partial R^2 values by an average of 0.05 across the 53 species tested here. The further addition of a term for basal area and geographic location (Hopkins index) each add another 0.01.

Because crown-ratio and geographic location data may not be available to some users and the gains from additional variables beyond stem diameter are only marginal for some species, the regression solutions from Equation 3 also are provided (Table 3). Gering and May (1995) built a case for simplicity when modeling the crown diameters of four species groups in Tennessee. Gill et al. (2000) concluded that dbh was the only predictor needed to model the crown radius of 13 species of western conifers, even though additional independent variables slightly improved some of their models. By individual species, gains in model precision resulting from the addition of crown ratio, basal area, and Hopkins index can be evaluated by comparing model statistics from Equation 3 in Table 3 with Equation 6 in Table 2.

Conclusions

Stem diameter is the strongest predictor of crown diameter for most tree species in the western United States. Stem diameter in quadratic form and additional terms for vertical crown ratio and geographic location improve the models for many species. Because stem diameter and crown ratio are correlated with stand density, an additional term for stand density is not generally needed but does yield minor improvement for a few species. Although model performance for some species can be enhanced by the addition of independent variables beyond stem diameter, the additional data may not be available to some potential users and the gains are marginal for some species, so there is also utility in presenting simpler models based solely on stem diameter.

Literature Cited

- BECHTOLD, W.A., M.E. MIELKE, AND S.J. ZARNOCH. 2002. Comparison of field methods and models to estimate mean crown diameter. *North. J. Appl. For.* 19(4):177-182.
- BECHTOLD, W.A. 2004. Largest-crown-width prediction models for X7 species in the eastern U.S. South. *J. Appl. For.* 27(4):269-278.
- BRAGG, D.C. 2001. A local basal area adjustment for crown width prediction. *North. J. Appl. For.* 18:22-28.
- DAWKINS, H.C. 1963. Crown diameters: Their relation to bole diameter in tropical forest trees. *Commonw. For. Rev.* 42:318-333.
- GERING, L.R., AND D.M. MAY. 1995. The relationship of diameter at breast height and crown diameter for four species in Hardin county, Tennessee. *South. J. Appl. For.* 19: 177-181.
- GILL, S.J., G.S. BIGING, AND E.C. MURPHY. 2000. Modeling conifer tree crown radius and estimating canopy cover. *For. Ecol. Manage.* 126:405-416.
- HANN, D.W. 1997. Equations for predicting the largest crown width of stand-grown trees in western Oregon. *For. Res. Lab., Oregon State Univ., Corvallis, OR. Res. Contr.* 17. 14 p.
- HANN, D.W. 1999. An adjustable predictor of crown profile for stand-grown Douglas-fir trees. *For. Sci.* 45:217-225.
- HAYS, R.L., C. SUMMERS, AND W. SEITZ. 1981. Estimating wildlife habitat variables. *USDI Fish and Wildlife Serv., FWS/OBS-81/47.* 111 p.
- HETHERINGTON, J.C. 1967. Crown diameter: Stem diameter relationships in managed stands of Sitka spruce. *Commonw. For. Rev.* 1967. 46:278-281.
- HOPKINS, A.D. 193X. Bioclimatics, a science of life and climate relations. *USDA Misc. Pub.* 280. Washington, DC. 188 p.
- KRAJCEK, J.E., K.A. BRINKMAN, AND S.F. GINGRICH. 1961. Crown competition: A measure of density. *For. Sci.* 7:35-42.
- MOEUR, M. 1981. Crown width and foliage weight of northern Rocky Mountain conifers. *USDA For. Serv., Res. Pap. INT-283.* 14 p.
- PAINE, D.P., AND D.W. HANN. 1982. Maximum crown width equations for southwestern Oregon tree species. *For. Res. Lab., Oregon State Univ., Corvallis, OR. Res. Pap.* 46. 19 p.
- SMITH, H.C., AND C.B. GIBBS. 1970. A guide to sugarbush stocking based on the crown diameter/d.b.h. relationship of open-grown sugar maples. *USDA For. Serv. Res. Pap. NE-171.* 8 p.
- STOLTE, K.W. 2001. Forest health monitoring and forest inventory analysis programs monitor climate change effect in forest ecosystems. *Human Ecol. Risk Assess.* 7(5): 1297-1316.
- USDA FOREST SERVICE. 2002. Forest inventory and analysis national core field guide, volume 1: Field data collection procedures for phase 2 plots, version 1.6. *USDA For. Serv. Int. Rep. For. Inv. Anal., Washington, DC.* 103 p.
- ZARNOCH, S.J., W.A. BECHTOLD, AND K.W. STOLTE. 2004. Crown condition as an indicator of forest health. *Can. J. For. Res.* 34: 1057-1070.