

BIOLOGY AND MANAGEMENT OF *LONGIDORUS AMERICANUS* IN A SOUTHERN USA NURSERY

M. M. CRAM¹ AND S. W. FRAEDRICH²

¹USDA FOREST SERVICE, FOREST HEALTH PROTECTION, ATHENS, GA, USA,

²USDA FOREST SERVICE, SOUTHERN RESEARCH STATION, ATHENS, GA, USA

mcram@fs.fed.us

ABSTRACT

A recently discovered nematode, *Longidorus americanus*, caused stunting of *Pinus taeda* seedlings at a forest-tree nursery in Georgia, USA. In growth chamber experiments, *L. americanus* significantly reduced the root size of *P. taeda* and *P. elliotii* seedlings. Although the root systems of *P. palustris* were not significantly reduced, it was found to be a host for *L. americanus*. The field where damage by *L. americanus* occurred was in continuous production of *Pinus* spp. and *Quercus alba* seedlings from 1990 to 1998. Fumigation by MC33 (methyl bromide 67%/chloropicrin 33%) in the spring of 1998 and again in 2000 remedied the problem for only one year; the seedlings were severely stunted during the second production years (1999 and 2001). Growth chamber tests found that *Quercus* spp. (*Q. alba*, *Q. rubra*, *Q. falcata*, *Q. virginiana*, and *Q. nigra*) were hosts for *L. americanus*. The populations of nematodes declined with *Q. acutissima*. Grasses used as cover crops in southern (USA) nurseries, including *Triticum aestivum*, *Secale cereale*, *Sorghum bicolor*, *Avena sativa* and *Panicum ramosum* are not hosts for *L. americanus*. Weeds such as *Cyperus esculentus* and *C. rotundus* that are common in southern (USA) nurseries were also not suitable hosts for the nematode. A fallow study conducted in a field and in growth chambers suggests that *L. americanus* does not survive in soil for extended periods (> 334 days) without a suitable host. Surveys for *L. americanus* have determined that the nematode occurs in oak forests that border the nursery and in an adjacent pine seed orchard.

Key words: *Longidorus americanus*, *Pinus* sp., *Quercus* sp., cover crops, *Cyperus* sp. fallow

INTRODUCTION

A new species of *Longidorus* discovered in 2000 was associated with stunting of *Pinus taeda* in a forest nursery located in southern Georgia, USA (FRAEDRICH, CRAM 2002). The nematode was recently described and named *Longidorus americanus* (HANDOO et al. 2005). Symptoms of seedling damage by *L. americanus* include chlorotic needles, reduced shoot growth and root systems that lack lateral and fine roots. The stunted seedlings were found in field 7S where similar damage had been observed since 1996 on both *P. taeda* and *P. elliotii* crops. The cause of this damage went undiagnosed until 2000 largely due to our reliance on a nematode testing laboratory that processed soil using a sugar flotation method. This technique is more suitable for smaller plant-parasitic nematodes. The adult nematodes of *L. americanus* are large (7 - 8 mm long) and require extraction methods specific for large nematodes (FLEGG 1967, SHURTLEFF, AVERRE 2000). Over the next 5 years, a series of surveys, dosage response tests, host range tests, and fumigation and fallow studies were conducted to investigate various aspects of the biology of the nematode and determine how best to manage this problem. In all our surveys and studies, nematodes were extracted using a technique developed by FLEGG (1967) with minor modifications by FRAEDRICH and CRAM (2002).

Tab. 1.

Field history of fumigation and crop rotations of pine, hardwoods, and cover crops for blocks 1 - 5 in field 7S of a south Georgia forest tree nursery (Byromville GA).

Date	Block 1 ⁺	Block 2	Block 3	Block 4	Block 5
1989	MC33 [‡]	MC33	MC33	MC33	MC33
1990	<i>P. taeda</i> *	<i>P. taeda</i>	<i>P. taeda</i>	<i>P. taeda</i>	<i>P. taeda</i>
1991	<i>P. taeda</i>	<i>P. taeda</i>	<i>P. elliotii</i>	<i>P. elliotii</i>	<i>P. elliotii</i>
1992	<i>P. taeda</i>	<i>P. taeda</i>	<i>P. elliotii</i>	Basamid/ <i>P. elliotii</i>	MC33/ <i>P. elliotii</i>
1993	<i>Q. alba</i> / <i>P. taeda</i>	<i>P. taeda</i>	Fallow	Fallow/ MC33	Fallow/ MC33
1994	<i>Q. alba</i>	<i>Q. alba</i>	<i>C. florida</i> / <i>L. styraciflua</i>	<i>P. elliotii</i>	<i>P. elliotii</i>
1995	<i>Q. alba</i>	<i>Q. alba</i>	<i>C. florida</i> / <i>L. styraciflua</i>	<i>P. virginiana</i>	<i>P. virginiana</i> / <i>P. clausa</i>
1996	<u><i>P. taeda</i></u> / <i>Q. alba</i>	<i>Q. alba</i>	<i>L. bicolor</i>	<i>L. bicolor</i>	<i>S. cereale</i>
1997	<u><i>P. taeda</i></u>	<i>P. taeda</i>	<i>L. bicolor</i>	<i>S. bicolor</i>	<i>S. bicolor</i>
1998	Fallow	MC33/ <i>P. elliotii</i>	MC33/ <i>P. elliotii</i>	Fumigation study/ <i>P. taeda</i>	Fumigation study/ <i>P. taeda</i>
1999	Fallow	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>	<i>P. taeda</i>	<i>P. taeda</i>
2000	MC33/ <i>P. taeda</i>	MC33/ <i>P. taeda</i>	MC33/ <i>P. taeda</i>	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>
2001	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>	<u><i>P. taeda</i></u>

⁺Text – stunted seedling damage

[‡]MC33 = methyl bromide 67% / chloropicrin 33%

**P. taeda* = *Pinus taeda*; *P. elliotii* = *Pinus elliotii*; *Q. alba* = *Quercus alba*;

P. virginiana = *Pinus virginiana*; *P. clausa* = *Pinus clausa*; *C. florida* = *Cornus florida*;

L. bicolor = *Lespedeza bicolor*; *L. styraciflua* = *Liquidambar styraciflua*;

S. cereale = *Secale cereale*

FIELD HISTORY

The crop and fumigation history of field 7S, provided in Table 1, is vital to understanding the development of the problem caused by *L. americanus*. Blocks 1 and 2 of this field were used to test rotations of pine and white oak from 1990 to 1996. This crop rotation was experimental and was not part of the normal crop rotation used by this nursery or other southeastern forest-tree nurseries. In 1996 and again in 1997, a few areas of stunted *Pinus taeda* seedlings occurred in block 1. Seedlings were examined and soil was sent to a nematode laboratory, but these efforts failed to find a cause for the damage. Blocks 2 and 3 in field 7S were fumigated in 1998 with 67% methyl bromide/33% chloropicrin (MC33) and planted to *P. elliotii*. Block 1 was too wet at the time of fumi-

gation and remained fallow during 1998. A fumigation study was also established in blocks 4 and 5, and sown with *P. taeda* (CRAM et al. 2002). In 1999, large areas of stunted seedlings occurred in block 2 and extended into block 3 (Fig. 1). Seedlings from the affected areas were examined for fungi and soil samples were sent to a nematode testing laboratory, but again the cause of the damage could not be identified. Blocks 1 to 3 were fumigated with MC33 in the spring of 2000 and sown in *P. taeda*.



Fig. 1. Stunted *Pinus taeda* in block 2 of field 7S, summer 1999

In 2000, small spots (3 - 9 m long) of stunted seedlings occurred in unfumigated control plots of the fumigation study in blocks 4 and 5 (FRAEDRICH, CRAM 2002). Soil samples were sent to a nematode lab, which reported low levels of *Pratylenchus*

sp. and *Criconebella* sp. Evaluations for pathogenic fungi were also inconclusive. Finally, we discovered large (5.4 - 9 mm long) plant-parasitic nematodes associated with stunted seedlings as we were examining unwashed roots; these nematodes were determined to be members of the genus *Longidorus*. We believe that the nematode was moved into these nonfumigated control plots during routine field operations before establishment of the study in 1998.

SURVEYS

1 Nursery blocks

In August and October of 2000, the populations of *L. americanus* were greatest in soil from the centers and margins of stunted areas compared to adjacent areas with healthy seedlings (FRAEDRICH, CRAM 2002). In August, 26 *L. americanus* per 25 g soil were obtained from the center of patches, 13.1 at the margins, and 0.5 and 0.2 at distances of 1.5 and 3 m from the margins, respectively. In October, there was an average of 25.8 *L. americanus* per 25 g of soil at the margins of patches, and only 1.5 *L. americanus* /25 g soil at locations 1.5 m outside the margin.

2 Outside the nursery field

Surveys were also performed from 2001 to 2003 in various locations within the nursery and in locations that border the nursery. These samples typically consisted of 8 - 10 cores at a 6" depth taken at specific locations and along transects. Soil samples were taken in 3 red cedar windrows, an oak (*Quercus* spp.) seed orchard, and oak and pine forests adjacent to the nursery. Sixteen composite soil samples were also obtained from *P. taeda* and *P. elliottii* seed orchards that border the nursery. *Longidorus americanus* was found in 37% soil samples from the pine seed orchards, and in an area of oak (primarily *Q. nigra*) trees that bordered the nursery.

Tab. 2.

Longidorus americanus population densities and root dry weights of southern pine species 26 weeks after infestation (experiment 2)[†]

Pine species	Initial needle	Final needle	Root dry [‡] weight (g)
	nematode		
	Number / container		
<i>Pinus taeda</i>	200	1257	0.159a
	0	0	0.295b
<i>Pinus elliotii</i>	200	1683	0.334a
	0	0	0.556b
<i>Pinus palustris</i>	200	820	0.681a
	0	3	0.825a

[†]Data obtained from FRAEDRICH et al. 2003 (Plant Dis., 87:1129:1132)

[‡]Means followed by the same letter do not differ significantly ($P \leq 0.05$) according to

t-test

Tab. 3.

Mean number of *Longidorus americanus* obtained from soil and roots of plant species 13 weeks after infestation with 100 nematodes/container

Plant species	<i>Longidorus</i> sp./400cc soil [†]	Total estimated <i>Longidorus</i> sp. per container
<i>Pinus taeda</i>	74a	295
<i>Quercus nigra</i> [‡]	94a	377
<i>Q. rubra</i>	56ab	223
<i>Q. akba</i> [‡]	42ab	168
<i>Q. falcata</i>	33ab	131
<i>Q. virginiana</i>	32ab	129
<i>Q. acutissima</i>	17bc	69
Fallow	1c	5

[†]Means followed by the same letter do not differ significantly ($\alpha = 0.05$) according to Tukey's HSD test. Square root transformation of nematode counts performed before analysis. Data analyzed as a randomized complete block design.

[‡]Means based on three replications

HOST RANGE STUDIES

1 Pine hosts

Population densities of *Longidorus americanus* increased on roots of *P. taeda* seedlings and damaged the root systems of seedlings in growth chamber tests. Root dry weights of seedlings decreased with respect to both the initial *L. americanus* dose, and the final population per container (FRAEDRICH, CRAM 2002). *Pinus elliotii* and *P. palustris* were also found to be hosts of *L. americanus* (FRAEDRICH et al. 2003). Population densities of *L. americanus* increased on roots of *P. elliotii* and *P. palustris*, but at the initial population densities used in these experiments only the root dry weights of *P. elliotii* seedlings were reduced by the nematode (Table 2). This lack of effect of the nematode on *P. palustris* may be related to the unique developmental characteristics of this pine

In the growth chamber study, the population of *L. americanus* also decreased exponentially over time ($P \leq 0.0001$) in fallow containers (FRAEDRICH et al. 2005 in press). *Longidorus americanus* was not detected after days 334 and 427. In containers planted with pine seedlings, the population density of *L. americanus* initially declined but subsequently increased after additional pine seedlings were added to containers. *Longidorus americanus* does not survive for extended periods in the upper 15 cm of fallow soil in the field (> 263 days) or in fallow containers under optimal conditions (> 334 days). The use of fallow or use of non-host cover crops should provide control of the *L. americanus* nematode in the southern Georgia nursery.

DISCUSSION AND CONCLUSIONS

Longidorus species have been found in southern pine nurseries and forests (HOPPER 1958, RUEHLE, SASSER 1962); however, there have been no reports of these nematodes damaging pine seedlings in nursery beds. In 2000, *Longidorus americanus* was discovered to cause stunting of *Pinus taeda* seedling at a south Georgia forest nursery (FRAEDRICH, CRAM 2002). *Pinus elliottii* and *P. palustris* were found to be hosts of *L. americanus* (FRAEDRICH et al. 2003).

Damage by *L. americanus* occurred in a field where the nursery had alternated *P. taeda* production with production of *Quercus alba*. Host suitability tests have found that *Q. alba* and other native oak species are hosts for *L. americanus*. The initial development of pine seedling stunting in block 1 of field 7S was most likely due to continuous cropping of pine and oak species over an extended period of time.

Surveys have found *L. americanus* outside the nursery under *Q. nigra* trees (FRAEDRICH, unpublished) and in an adjacent *P. taeda* and *P. elliottii* seed orchards. Locations such as these may provide the sources for reintroduction in nursery fields through soil and water movement (e. g. floods, equipment, wind and animals). It is possible that this nematode has been introduced to the nursery fields in the past, but that the typical rotations of tree seedling crops with grass cover crops has provided adequate control of this nematode.

In 1998 and 2000, the nursery attempted to remedy the problem of stunting in pine seedling crops by using methyl bromide fumigation. Fumigation suppressed disease development in the first seedling crop after fumigation, but the problem reappeared during the second year of production. Although fumigation has been found to depress high nematode populations (DROPKIN 1989), nematode populations can rebound quickly and significantly impact subsequent seedling crops (MCKENRY, THOMASON 1976). The rebound of nematode populations after fumigation was demonstrated by FRAEDRICH and DWINELL (2003). In this study, dazomet and metam sodium reduced the needle nematode to nondetectable levels in the upper 15 cm of soil but populations subsequently increased during loblolly pine production to levels comparable to those in control plots by the end of the growing season (FRAEDRICH, DWINELL 2003).

The nursery where the *Longidorus* problem was found typically rotates from pine production to cover crops every two years. In fact, the majority of southern nurseries rotate their tree seedling production with small grain crops yearly or biennially (CRAM, FRAEDRICH 1997). All of the small grain cover crops used at this nursery were found to be non-hosts for *L. americanus*. These results coupled with the findings that *L. americanus* does not appear to survive extended periods in fallow soil may help explain why this nematode problem has not been a reoccurring problem in other fields at the nursery. We believe that the nursery will not have a serious problem with *L. americanus* in the future if they maintain the practice of alternating cover crops and periods of fallow with pine seedling production on a regular basis.

REFERENCES

- CRAM, M. M., ENEBAK, S. A., DWINELL, L. D., FRAEDRICH, S. W. 2002. Chloropicrin, EPTC, and plant growth-promoting rhizobacteria for managing soilborne pests in pine nurseries. In: Dumroese, R. K., Riley, L. E., Landis, T. D., technical coordinators. National Nursery Proceedings 1999, 2000, 2001. Ogden (UT): USDA Forest Service, Rocky Mountain Research Station. Proceedings RMRS-P-24. p. 69-74.
- CRAM, M. M., FRAEDRICH, S. W. 1997. Survey of southern forest nurseries: fumigation practices and pest management concerns. In: Landis, T. D. & South, D. B., technical coordinators. National Proceedings: Forest and Conservation Nursery Associations-1996. Portland, OR: USDA Forest Service, Pacific Northwest Research Station. Proceedings PNW-GTR-389:19-27.
- DROPKIN, V. H. 1989. Introduction to Plant Nematology. 2nd ed. Wiley & Sons, Inc.
- FLEGG, J. J. M. 1967. Extraction of *Xiphinema* and *Longidorus* species from soil by a modification of Cobb's decanting and sieving technique. *Annals of Applied Biology*, 60:429-437.
- FRAEDRICH, S. W., CRAM, M. M. 2002. The association of a *Longidorus* species with stunting and root damage of loblolly pine (*Pinus taeda* L.) seedlings. *Plant Disease*, 86:803-807.
- FRAEDRICH, S. W., CRAM, M. M., HANDOO, Z. A. 2003. Suitability of southern pines, other selected crops, and nutsedge to a *Longidorus* sp. Associated with stunting of loblolly pine seedlings. *Plant Disease*, 87:1129-1132.
- FRAEDRICH, S. W., CRAM, M. M., HANDOO, Z. A. 2005. The effects of fallow on *Longidorus americanus*, a nematode associated with stunting of loblolly pine seedlings in Georgia, USA. *Nematology*, 7:487-493.
- FRAEDRICH, S. W., DWINELL, L. D. 2003. Effect of dazomet, metam sodium and oxymyl on production of loblolly pine seedlings and population densities of a *Longidorus* sp. *Phytopathology*, 93 (Supplement):S27 (abstract).
- HANDOO, Z. A., CARTA, L. K., SKANTAR, A. M., YE, T., SUBBOTIN, S., FRAEDRICH, S. W., CRAM, M. M. 2005. Morphological and molecular characterization of *Longidorus americanum* n. sp. (Nematoda: Longidoridae), a needle nematode parasitizing pine in Georgia. *Journal of Nematology*, 37:94-104
- HOPPER, B. E. 1958. Plant-parasitic nematodes in the soils of southern forest nurseries. *Plant Disease Reporter*, 42:308-314.
- MCKENRY, M. V., THOMASON, I. J. 1976. Dosage values obtained following pre-plant fumigation for perennials. II. Using special methods of applying methyl bromide and 1,3-dichloropropene nematicides. *Pesticide Science*, 7:535-544.
- RUEHLE, J. L., SASSER, J. N. 1962. The role of plant-parasitic nematodes in stunting of pines in southern plantations. *Phytopathology*, 52:56-68.
- SHURTLEFF, M. C., AVERRE, C. W., III. 2000. *Diagnosing Plant Diseases Caused by Nematodes*. APS Press, St. Paul, MN.

IUFRO Working Party 7.03.04

Diseases and insects in forest nurseries

The sixth meeting of the IUFRO Working Party 7.03.04
(Diseases and insects in forest nurseries)
was held from September 11 - 14, 2005,
in Uherské Hradiště, Czech Republic



Forestry and Game Management Research Institute Strnady

2007

Communicationes Instituti Forestalis Bohemicae

Volumen 23



Forestry and Game Management Research Institute Strnady

2007

ISSN 1211-2992
ISBN 978-80-86461-74-8