



2000

Annual  
International Research  
Conference  
on  
Methyl Bromide  
Alternatives  
and Emissions Reductions

November 6 - 9, 2000

Rosen Plaza Hotel  
9700 International Drive  
Orlando, Florida 32819 -1-8114

Sponsored by  
Methyl Bromide Alternatives Outreach

in cooperation with

US Environmental Protection Agency  
and  
US Department of Agriculture

**2000**

**Annual International Research  
Conference on Methyl Bromide  
Alternatives and Emissions Reductions**

*Sponsored by  
Methyl Bromide Alternatives Outreach*

*In cooperation with  
Crop Protection Coalition,  
U S Environmental Protection Agency  
And  
U S Department of Agriculture*

**Office: Methyl Bromide Alternatives Outreach**

**Gary L. Obenauf**  
144 W. Peace River Drive  
Fresno, CA. 93711-6953  
Tel: 559 447 2127  
Fax: 559 436 0692  
e-mail: [gobenauf@agrc.cncf.org](mailto:gobenauf@agrc.cncf.org)

**Rosemary Obenauf**  
144 W. Peace River Drive  
Fresno, CA. 93711-6953  
Tel: 559 447 2127  
Fax: 559 436 0692  
e-mail: [robenauf@agrc.cnchost.com](mailto:robenauf@agrc.cnchost.com)

## Effect of methyl bromide on *Bursaphelenchus xylophilus* in pine wood

L. David Dwinell

USDA Forest Service, Athens, GA 30602

One of the unintended consequences of the globalization of trade has been an increase in the frequency of introductions and the numbers of exotic species intercepted at ports of entry. This is largely due to the increase in volume of trade and a broadening of trading partners, especially with the Pacific Rim and Asia. During the years 1985 through 1998, the U.S. Department of Agriculture (USDA) Animal and Plant Health Inspection Service (APHIS) made 6,952 interceptions of exotic pests on wood articles at nearly all of 100 port locations throughout the United States. The intercepted insects originated from at least 95 countries worldwide. A global strategy needs to address the exotic pest problems. Whether dealing with the export or import of transported wood, the pest control principles would be quite similar.

Since 1986, research on decontaminating softwood chips, green lumber, and logs concentrated on the pinewood nematode, *Bursaphelenchus xylophilus* (PWN), and its *Monochamus* vectors. The PWN, which is native to North America, is a destructive pest of the pine forests of Japan and China. The nematode has also been introduced into Taiwan, South Korea, and Portugal. On January 1, 2000, China put into place regulations to ensure that solid wood packing material (dunnage, crating, packing cases, pallets, et al.) (SWPM) entering their country was not infested by the PWN. Recent interceptions of the PWN in SWPM by the Finnish Plant Quarantine Service have resulted in emergency legislation by Finland. Although the use of elevated temperatures (56°C for 30 minutes) to eradicate the PWN and its *Monochamus* vectors in SWPM is the preferred mitigation method, there are circumstances when fumigation with methyl bromide is the most feasible and cost effective method available. Pre-shipment and quarantine uses, which are allowed under the Montreal protocol, would be an interim solution for countries without the resources or facilities to use heat pasteurization to decontaminate SWPM without seriously disrupting trade.

### Methods and materials

In early June 2000, three shortleaf pines (*Pinus echinata*) were felled and their main stems were then sawn into 91-102-cm-long logs. The logs were left in the forest where they were subsequently colonized by pine sawyers (*Monochamus* spp.), vectors of the pinewood nematode. In early August, the logs were sawn into disks, cants, or boards. The disks were 1-, 2-, 3- and 6-cm in thickness and had a mean diameter of 19 cm. The cants were 12.7 x 12.7 x 96.5 cm. The boards were 2.5 x 10.2 x 91 cm and 5.0 x 10.2 x 91 cm.

The disks, cants and boards were divided into three treatment group. Each load or replicate consisted of one each of the four disk sizes, a cant, nine 2.5 x 10.2 x 91 cm boards, and nine 5.0 x 10.2 x 91 cm boards. Before and after fumigation, the disks, cants, and boards were sampled for the PWN by drilling three 2.5 cm deep holes with an auger bit. After fumigation, the cants were also sampled at 2.5-5.0-cm depth. The borings within a disk, cant or board were pooled and the Baermann funnel procedure was used to extract the nematodes (Dwinell et al. 1994). The

presence or absence of the PWN in the samples were recorded.

The fumigation was done in three chambers constructed of plastic pipes and covered with a polyethylene sheet. The chambers, each of which had a volume of 2.83 m<sup>3</sup>, were on a concrete pad. The edges of the cover were sealed with dirt prior to fumigation. The air temperature in the chambers during fumigation was recorded using Hobo data loggers (Onset Computer Corporation). The three fumigation treatments were 0, 0.45, and 0.90 kg methyl bromide/2.83 m<sup>3</sup> and the treatment duration was 24 hours. There were three replicates of each treatment.

## Results and discussion

Prior to fumigation, the PWN was extracted from all the disks, cants, and boards (Table 1). The PWN was not extracted from any of the disks, cants or boards which had been fumigated, regardless of rate. The PWN was extracted from the control treatment disks, cants, and boards. The temperature during fumigation averaged 46.2°C for the three chambers, and the mean maximum temperature was 56.6°C. High temperatures during fumigation probably contributed to the penetration and efficacy of methyl bromide. The data is consistent with earlier reports on using methyl bromide to eradicate the PWN in pine wood. For example, in 1994, I reported that methyl bromide at the rate of 240 g/m<sup>3</sup> would eradicate the PWN in small logs of eastern (*P. strobus*), slash (*P. elliotii* var. *elliotii*), and loblolly (*P. taeda*) pines. Wang et al. (1995) reported that the efficacy of methyl bromide to eliminate the PWN in logs was dependent on temperature, dosage, duration of fumigation, the wood size, moisture content of the wood, and the position of the wood in the stack. Methyl bromide, as a pre-shipment or quarantine use, shows promise as an interim or emergency treatment for decontaminating SWPM.

## References

- Dwinell, L.D. S. Avramidis, and J.E. Clark. 1994. Evaluation of radio-frequency/vacuum dryer for eradicating the pinewood nematode in green sawn wood. *Forest Products J.* 44:19-24.
- Dwinell, L.D. 1994. Evaluation of sodium borate for the control of nematodes and wood-borers in pine logs. In: *Ann. Intl. Res. Conference Methyl Bromide Alternatives and Emissions Reductions*, 13-16 November 1994, Kissimmee, FL. p. 95-1.
- Dwinell, L.D. 1996. Methyl bromide alternatives for decontaminating softwood chips, lumber, and logs. *Ann. Intl. Res. Conference Methyl Bromide Alternatives and Emissions Reductions*, 3-6 November 1996, Orlando, FL. pp. 64-1,3.
- Wang, Y., Y. Song, X. Zang, M. Ge, and J. Zhao. 1995. Study of the effect of fumigation of the wood infested by pine wood nematode with methyl bromide. In: *Intl. Symposium On Pine Wilt Disease*, 31 Oct. - 5 Nov., 1995, Beijing, China

Table 1. The effect of methyl bromide fumigation on *Bursaphelenchus xylophilus* (PWN) in pine disks, cants, and green lumber

Pine wood Dimensions	Methyl bromide (kg/2.83m <sup>3</sup> )					
	-0-		0.45		0.90	
	Number samples with PWN/total number of samples before and after fumigation <sup>a</sup>					
	Pre-	Post-	Pre-	Post-	Pre-	Post-
2.5 cm disk <sup>b</sup>	3/3	3/3	3/3	0/3	3/3	0/3
5.0 cm disk <sup>b</sup>	3/3	3/3	3/3	0/3	3/3	0/3
7.5 cm disk <sup>b</sup>	3/3	3/3	3/3	0/3	3/3	0/3
15.2 cm disk <sup>b</sup>	3/3	3/3	3/3	0/3	3/3	0/3
12.7 cm x 12.7 cm cant <sup>c</sup>	3/3	3/3	3/3	0/3	3/3	0/3
2.5 cm x 10.1 cm board <sup>d</sup>	9/9	9/9	9/9	0/9	9/9	0/9
5.0 cm x 10.1cm board <sup>d</sup>	9/9	9/9	9/9	0/9	9/9	0/9
<b>TOTALS</b>	<b>33/33</b>	<b>33/33</b>	<b>33/33</b>	<b>0/33</b>	<b>33/33</b>	<b>0/33</b>

<sup>a</sup> Each treatment combination was replicated three times.

<sup>b</sup> Disk thickness; the disks had a mean diameter of 19 cm.

<sup>c</sup> The cants had a mean length of 96.5 cm. After fumigation, the cants were sampled at 2.54 cm and 5.0 cm depths.

<sup>d</sup> The boards had a mean length of 91 cm.

The use of trade names or firm names in this publication is for the readers' information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

This publication reports research involving pesticides. It does not contain recommendations for their uses, nor does it imply that the uses discussed have been registered. All uses of pesticides must be registered by appropriate State or Federal agencies before they can be recommended.

Caution: Pesticides can be injurious to humans, domestic animals, desirable plants, fish, or other wildlife if they are handled or applied improperly. Use all pesticides selectively and carefully. Follow recommended practices for the disposal of surplus pesticides and pesticide containers.