

Reprinted from

Forest Ecology and Management

Forest Ecology and Management 126 (2000) 13-24

Growth and development of loblolly pine in a spacing trial planted in Hawaii

William R. Harms^{a,1}, Craig D. Whitesell^{b,1}, Dean S. DeBell^{c,*}

^aUSDA Forest Service, Southern Research Station, Center for Forested Wetlands Research, 2730 Savannah Highway, Charleston, SC 29414, USA

^bUSDA Forest Service, Pacific Southwest Research Station, Institute of Pacific Islands Forestry, 1151 Punchbowl Street, Honolulu, HI 96813, USA

^cUSDA Forest Service, Pacific Northwest Research Station, 3625, 93rd Avenue SW Olympia, WA 98512, USA

Received 13 July 1998; accepted 12 February 1999



Forest Ecology and Management

Aims and scope. Forest *Ecology* and Management publishes scientific articles concerned with forest management and conservation, and in particular the application of biological, ecological and social knowledge to the management of man-made and natural forests. The scope of the journal includes all forest ecosystems of the world. A refereeing process ensures the quality and international interest of the manuscripts accepted for publication. The journal aims to encourage communication between scientists in disparate fields who share a common interest in ecology and natural-resource management, and to bridge the gap between research workers and forest managers in the field to the benefit of both. The journal should be of interest to research workers, managers and policy makers in forestry, natural resources, ecological conservation and related fields.

FOUNDING EDITOR

Laurence L. Roche, Murroe, Ireland

EDITORS-IN-CHIEF

For the Americas, Australia, New Zealand and the Pacific:

R.F. Fisher

Temple-Inland Forest
PO. Drawer N
303 S. Temple Drive
Diboll TX 75941-0814, USA

BOOK REVIEWS EDITOR

Margaret R. Gale
School of Forestry and Wood Products
Michigan Technological University
1400 Townsend Drive
Houghton, MI 49931, USA

EDITORIAL ADVISORY BOARD

G. Abrahamsen, Agricultural University of Norway, Ås, Norway

R. Alfaro, Canadian Forestry Service, Victoria, B.C., Canada

F. Andersson, Swedish University of Agricultural Sciences, Uppsala, Sweden

P.M.S. Ashton, Yale University, New Haven, USA

P. Attiwill, University of Melbourne, Parkville, Vic., Australia

J. Boyle, Oregon State University, Corvallis, USA

S. Brown, US Environmental Protection Agency, Corvallis, USA

J.C. Calvo, Institute of Technology, Cartago, Costa Rica

R.-J. Chen, University of Hong Kong, Pokfulam Road, Hong Kong

J.D. Deans, Institute of Terrestrial Ecology, Penicuik, Midlothian, UK

R.M. DeGraaf, USDA Forest Service, University of Massachusetts, Amherst, MA, USA

S. Diamandis, Forest Research Institute, Thessaloniki, Greece
D.P. Dykstra, President, World Forestry Centre, 4033 SW Canyon Road, Portland, OR 97221 USA

E.P. Farrell, University College Dublin, Dublin, Ireland

P.M. Fearnside, Instituto Nacional de Pesquisas da Amazonia-INPA, Manaus-Amazonas, Brazil

R.D. Finlay, Department of Forest Mycology & Pathology, SLU, Box 7026, SE-75007 Uppsala, Sweden

P.H. Freer-Smith, Forestry Commission, Farnham, UK

O. Garcia, ENGREF, Nancy, France

D.W. Gilmore, University of Minnesota, North Central Experiment Station, 1861 Highway 169 East, Grand Rapids, MN 55744, USA

R.A. Goyer, Louisiana State University, Baton Rouge, LA, USA

J.B. Hall, University College of North Wales, Bangor, UK

For the rest of the world:

G.M.J. Mohren

Silviculture and Forest Ecology Group

Department of Environment Sciences

Wageningen Agricultural University

P.O. Box 342

6700 AH Wageningen, The Netherlands

F. Houllier, Campus International de Baillarguet, Laboratoire Associe de Modelisation des Plantes (AMAP), Montpellier, France

B.M. Kumar, Kerala Agricultural University, Kerala, India

J.P. Lassoie, Cornell University, Ithaca, NY, USA

J.N. Long, Utah State University, Logan, UT, USA

A.E. Lugo, International Institute of Tropical Forestry, Rio Piedras, PR, USA

J.A. Maghembe, SADCC / ICRAF Agroforestry Project, Zomba, Malawi

F. Makeschin, Institut für Bodenkunde und Standortslehre, Tharandt, Germany

D.C. Malcolm, University of Edinburgh, Edinburgh, UK

E. Mälkönen, Finnish Forest Research Institute, Vantaa, Finland

M.A.R. Nahuz, Instituto de Pesquisas Tecnológicas,

São Paulo, SP, Brazil

R. Päivinen, European Forestry Institute, Joensuu, Finland

S.G. Pallardy, University of Missouri, Columbia, MO, USA

R.F. Powers, Pacific Southwest Research Station, Redding, CA, USA

T. Pukkala, University of Joensuu, Joensuu, Finland

F.E. Putz, University of Florida, Gainesville, FL, USA

L. Rasmussen, RISO, Roskilde, Denmark

D.D. Reed, Michigan Technological University, Houghton, MI, USA

R. Sands, University of Canterbury, Christchurch, NZ

J.A. Stanturf, Stoneville, MS, USA

J.R. Toliver, USDA Forest Service, Washington, DC, USA

J.K. Venclay, Southern Cross University, PO Box 157,

Lismore NSW 2480 Australia

K. von Weissenberg, University of Helsinki, Helsinki, Finland

D. Whitehead, Manaaki Whenua Landcare Research, Lincoln, New Zealand

Publication information: Forest Ecology and Management (ISSN 03781127). For 2000 volumes 123–136 are scheduled for publication. Subscription prices are available upon request from the Publisher or from the Regional Sales Office nearest you or from this journal's website (<http://www.elsevier.nl/locate/foreco>). Further information is available on this journal and other Elsevier Science products through Elsevier's website: (<http://www.elsevier.nl>). Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by standard mail (surface within Europe, air delivery outside Europe). Priority rates are available upon request. Claims for missing issues should be made within six months of the date of dispatch.

Orders, claims, and product enquiries: please contact the Customer Support Department at the Regional Sales Office nearest you: New York: Elsevier Science, PO Box 945, New York, NY 10159-0945, USA; Tel. (+1) (212) 633 3730, [toll free number for North American customers: 1-888-4ES-INFO (437-4636)]; fax: (+1) (212) 633 3680; e-mail usinfo-f@elsevier.com

Amsterdam: Elsevier Science, PO Box 211, 1000 AE Amsterdam, The Netherlands; Tel.: (+31) 20 485-3757; fax: (+31) 20 485 3432; e-mail ninfo-f@elsevier.nl

Tokyo: Elsevier Science, 9-15, Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel. (+81) (3) 5561 5033; fax (+81) (3) 5561 5047; e-mail: info@elsevier.co.jp

Singapore: Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192; Tel. (+65) 434 3727; fax: (+65) 337 2230; e-mail: asiainfo@elsevier.com.sg

Rio de Janeiro: Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; phone: (+55) (21) 509 5340; fax: (+55) (21) 507 1991; e-mail: elsevier@campus.com.br [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]



ELSEVIER

Forest Ecology and Management 126 (2000) 13-24

Forest Ecology
and
Management

www.elsevier.com/locate/foreco

Growth and development of loblolly pine in a spacing trial planted in Hawaii

William R. Harms^{a,1}, Craig D. Whitesell^{b,1}, Dean S. DeBell^{c,*}

^aUSDA Forest Service, Southern Research Station, Center for Forested Wetlands Research, 2730 Savannah Highway, Charleston, SC 29414, USA

^bUSDA Forest Service, Pacific Southwest Research Station, Institute of Pacific Islands Forestry, 1151 Punchbowl Street, Honolulu, HI 96813, USA

^cUSDA Forest Service, Pacific Northwest Research Station, 3625, 93rd Avenue SW Olympia, WA 98512, USA

Received 13 July 1998; accepted 12 February 1999

Abstract

Loblolly pine (*Pinus taeda* L) was planted at four square spacings (1.8, 2.4, 3.0, and 3.7 m) on the Island of Maui in 1961, and measured periodically for 34 years. Patterns of stand growth and development were examined and compared with yield model estimates of stand characteristics of plantations of the same initial spacings, ages, and site index in the southeastern United States. The Hawaiian plantings had much higher survival at all spacings and sustained high diameter growth in the face of intense competition. At Age 34, the 1.8 m spacing had 1585 stems/ha averaging 24.1 m tall and 28.8 cm DBH; the widest spacing (3.7 m) had 725 stems/ha, 26.1 m tall and 38.2 cm DBH. The highest basal areas (~ 100 m²/ha) were double maxima attained in the southeastern United States and were reflected in similar differences in volume yields. The Hawaiian plantings demonstrate that growth potential of loblolly pine is far greater than is apparent from observations on plantations in its native habitat. To capture this potential in other situations, research must identify the tree, stand, and environmental characteristics associated with low mortality rates and high diameter growth in Hawaii, and, conversely, the factors that limit loblolly's potential in the southeastern United States. © 2000 Elsevier Science B.V. All rights reserved.

Keywords: *Pinus taeda*; Stand dynamics; Stockability; Self-thinning; Growth and yield

1. Introduction

Loblolly pine (*Pinus taeda*) is the most common and commercially important pine species in the south-

eastern United States. It is well adapted to the wide range of soils, sites, and environments that occur over its natural range, which extends from southern New Jersey south to central Florida and west to eastern Texas (Baker and Langdon, 1990). Because of its rapid growth and adaptability it has been planted in a number of other countries, including Argentina, Australia, Brazil, China, New Zealand, South Africa, and also in the state of Hawaii, and found to perform

*Corresponding author. Tel.: +1-360-753-7667; fax: +1-360-956-2346.

E-mail address: ddebell/r6pnw_olympia@fs.fed.us (D.S. DeBell)

¹Retired.

exceptionally well so that it is now grown commercially in many of these countries (Burns and Hu, 1983; Schultz, 1997). China and Brazil have the largest loblolly pine management programs. Annually they produce and plant more than 320 million seedlings (Schultz, 1997). The most successful plantings are usually between latitudes of 24° and 30° N or S, and at elevations of 500-900 m. In Brazil loblolly pine is grown on 15-25-year rotations for both fiber and solid wood products. Stand yields of 15-25 m³/ha/year are common and yields up to 35 m³/ha/year have been reported. Height growth curves indicate heights of 27-32 m by Age 20 on the best sites and 19-23 m on average sites (Machado, 1980). In South Africa loblolly pine is grown at rotational lengths of 18-35 years. Mean annual increment is ca. 18 m³/ha/year but can exceed 28 m³/ha/year on the best sites. Trees can reach a height of 30 m and a DBH of 45 cm at 35 years (Schultz, 1997). There are numerous published reports on individual aspects of the growth of loblolly pine planted in the subtropics, e.g., seed source (Baldanzi, 1978; Falkenhagen, 1978), environment (Burns and Hu, 1983; Harms et al., 1994; Schultz, 1997), site index (Machado, 1980), thinning (Machado et al., 1990), and carrying capacity and thinning (Strub and Bredenkamp, 1985), but few data or analyses of the quantitative aspects of stand dynamics and growth, especially as compared to growth of native plantations, have been published. One exception is a series of reports that examine the growth of a spacing trial planted in Hawaii to evaluate the potential for loblolly pine to augment the state's softwood timber requirements. This spacing trial has been measured periodically for 34 years, and has provided a rich record of stand growth and development of an exceptionally fast-growing plantation. Two of the reports evaluate early survival, height and diameter growth in relation to spacing at ages 7 and 11 years (Whitesell, 1970, 1974), a third evaluates stand productivity and stockability differences between loblolly pine in Hawaii and the southeastern US at Age 25 years (DeBell et al., 1989), and a fourth examines tree, stand, and environmental components of growth and stockability at Age 26 years (Harms et al., 1994).

In this paper, we examine the Hawaii data for the broader, overall, effects of initial spacing on the temporal patterns of stand growth and development through a full rotation. Survival, height and diameter

growth; basal area and volume growth and yield; and stand structural characteristics from the time of establishment to Age 34 years are reported. To put the growth of this plantation in perspective, we compare the various components of its development with a southeastern US 'standard' derived from a loblolly pine yield model (Hafley et al., 1982).

2. Methods

The spacing trial was established in 1961 as a research study in the Olinda Forest Reserve on the island of Maui (Whitesell, 1970). The site is 1140 m asl, on the northeast side of east Maui at latitude 20°49' N. The mean air temperature on Maui is 14°C and varies 3°C from summer to winter; average annual rainfall is 1143 mm with a wet winter and spring and a relatively dry summer (Harms et al., 1994). Growth is possible throughout the year. The soil is Olinda loam, in the subgroup Entic Dystrandepts. Typically, the surface layer is a dark reddish-brown granular loam ca. 15 cm thick over a dark reddish-brown and yellowish-red subangular blocky silty clay loam subsoil. This soil developed in volcanic ash over andesite or basalt. It is slightly acid (pH 6.2-6.4) in both the surface layer and subsoil. Olinda soil is well drained, with a moderately rapid permeability (Foote et al., 1972). Average site index of the plantation is 24 m (base Age 25 years).

The field layout of the plantation consists of four blocks each with four square spacings: 1.8, 2.4, 3.0, and 3.7 m, planted in a randomized 4 x 4 Latin square design. Each plot is square, 0.11 ha in area, with the 5 x 5 rows in the center forming a 25-tree measurement sub-plot. All sub-plots had two or more exterior isolation rows. At the time of planting, the ground cover was a 3-inch sod layer of kikuyu grass (*Pennisetum clandestinum* Hochst.), rattail grass (*Sporobolus capensis* [Wild.] Kunth), and gorse (*Ulex europaeus* L.). The grass was heavily grazed just before planting. The sod was removed at each planting spot, and holes 30 cm deep were dug for the seedlings. Planting stock was 1-1 nursery-grown seedlings of unknown seed source origin. The 21 trees that died during the first year were replaced.

Measurements were made at ages 4, 7, 11, 20, 25, 26, and 34 years. Diameter at breast height (DBH,

1.37 m) of all living sample trees was measured each time. Heights of all sample trees were measured at ages 4, 7, 11, and 26 years, and on a systematically chosen nine-tree sub-sample at ages 20, 25, and 34 years. At Age 26, 16 trees from the border rows covering the range of DBH were felled and measured to obtain stem volumes. The volumes were used to select a published loblolly pine volume table that could be used to derive acceptable volume estimates for the plantation. The equations published by Burkhart (1977) gave satisfactory estimates: a paired *t*-test of differences between observed and estimated volumes of the 16 sample trees was not significant ($p = 0.05$). A local volume table was developed for each spacing and year from ages 7 through 34 by applying Burkhart's equation to the plot height-sample data to obtain volume, and fitting volume/basal area regressions (Clutter et al., 1983). Plot volumes were computed by summing estimated volumes of the plot trees and expanding to the unit area. The study was installed and measured in imperial units and converted to metric units for publication.

In two previous papers, growth comparisons were made using data from a loblolly pine spacing trial in South Carolina (DeBell et al., 1989; Harms et al., 1994). The spacings in the South Carolina plantation were the same as those at Olinda and the ages at which data were collected were similar, but the site index of the South Carolina plantation was 21 m, 3 m shorter than the Olinda plantation. Rather than base the comparisons in this report on data with a 3 m difference in site index, we elected to make comparisons using the loblolly pine yield model developed by Hafley et al. (1982) for plantations in the Southeast. The model has been shown to predict observed stand development with respect to initial spacing quite well (Buford, 1991). Model input includes site index (base Age 25 years), trees per acre planted, and percent survival at Year 1. Our comparisons use estimates obtained from the yield model for site index 80 (24 m), the average site index observed for the Olinda plantation, and for the same spacings and measurement ages as the Olinda plantation. Survival at Year 1 was assumed to be 100%. All graphical and numerical information presented in this report for southeastern US plantations were derived from the Hafley et al. (1982) model.

3. Results

3.1. Stand development

The major stand attributes of the Olinda plantation are summarized in Table 1. Early development of the plantation was described by Whitesell, 1970, 1974. He observed that the tree crowns had closed in the 1.8 m spacing by the fourth year at an average stand height of 4.5 m, by the seventh year in the 2.4 m and 3.0 m spacings, at an average stand height of 8.8 m, and that they were nearly closed in the 3.7 m spacing by the 11th year, when the average stand height was 12.7 m. Length of the green crown in the 1.8 m spacing decreased from 64% of tree height at ages 7 to 29% at Age 26, and from 81% to 36% in the 3.7 m spacing during the same period.

3.1.1. Survival and mortality

Mortality in the Olinda plantation was density related; no causes other than competition for growing space were identified, i.e., except for minor mortality after planting, losses were confined to trees in the suppressed and intermediate crown classes, and number of trees dying varied directly with initial density (Long and Smith, 1984; Oliver and Larson, 1990). Self-thinning began earliest and trees died in greatest numbers in the 1.8 m spacing, but losses did not become substantial until after Age 20 (Fig. 1 and Table 1). By Age 34 an average of 47% of the trees in the 1.8 m plots had died, and although mortality had begun to occur in the 2.4 m and 3.0 m plots, only 19% and 15%, respectively, had died by the 34th year. Less than 3% of the trees in the 3.7 m plots had died by the 34th year (Table 1). Survival curves from the yield model predictions for plantations in the Southeast show the same trends and relationships to spacing as in Hawaii, but self-thinning begins at much younger ages in all spacings, and the rates of mortality, and the absolute numbers of trees lost are greater, resulting in a pronounced divergence of the curves from the Olinda data (Fig. 1). By Age 34 years, predicted mortality in the Southeast is 67% in 1.8 m spacings, 47% in 2.4 m spacings, and 30% and 18%, respectively, in 3.0 m and 3.7 m spacings. At both locations, the survival curves were progressively declining and converging as a result of self-thinning. At Age 34, the curves of the closest and widest spacings for the

Table 1
Means for stand components of the Olinda plantation, Maui, Hawaii, at age of measurement ^a

Spacing ^b (m)	Plantation age (years)					
	4	7	11	20	25	34
Number surviving (trees per ha)						
1.8 x 1.8	2990	2960	2841	2601	2063	1585
2.4 x 2.4	1615	1615	1615	1581	1463	1362
3.0 x 3.0	1066	1066	1055	1023	990	915
3.1 x 3.7	746	746	746	740	740	725
Cumulative mortality (%)						
1.8 x 1.8	0.0	1.0	5.0	13.0	31.0	47.0
2.4 x 2.4	3.9	3.9	3.9	5.9	12.9	18.9
3.0 x 3.0	1.1	1.1	2.1	5.1	8.2	15.1
3.7 x 3.7	0.0	0.0	0.0	0.8	0.8	2.8
Height (m)						
1.8 x 1.8	4.5	9.2	11.7	18.0	20.3	24.1
2.4 x 2.4	4.4	8.7	11.9	17.5	20.0	24.4
3.0 x 3.0	4.4	9.0	12.4	18.5	20.5	26.1
3.7 x 3.7	4.5	9.3	12.7	19.8	22.0	26.1
Quadratic mean diameter at breast height (cm)						
1.8 x 1.8	7.7	13.8	17.0	21.4	24.0	28.8
2.4 x 2.4	8.2	16.0	19.9	24.9	26.9	30.0
3.0 x 3.0	8.4	17.7	22.1	28.9	30.5	34.4
3.7 x 3.7	9.1	19.6	25.1	32.1	34.3	38.2
Basal area per ha (sq m)						
1.8 x 1.8	13.8	44.5	63.9	93.3	93.4	103.1
2.4 x 2.4	8.6	32.6	50.4	76.9	82.9	96.5
3.0 x 3.0	5.9	26.2	42.6	67.6	12.9	85.2
3.7 x 3.7	4.9	22.6	38.7	62.0	68.5	83.0
Total volume per ha (cu m outside bark)						
1.8 x 1.8	75	274	440	828	957	1324
2.4 x 2.4	38	182	321	668	824	1191
3.0 x 3.0	37	140	265	584	729	1079
3.7 x 3.7	20	92	251	576	717	1053

^a Each value is the average of four replicate plot means.

^b Trees per ha planted: 1.8 m = 2990; 2.4 m = 1680; 3.0 m = 1078; 3.1 m = 146.

Southeast were 363 trees/ha apart, while the Olinda curves were still 860 trees/ha apart.

3.1.2. Height

Spacing had no significant effect on average stand height, although trees in the 3.7 m spacing were consistently taller than those in the closer spacings as early as the 7th year (Harms *et al.*, 1994). By Age 25 average height of the 3.7 m spacing was 22 m, 2 m taller than the other spacings. At Age 34 average

heights of the 3.0 and 3.7 m spacings were equal and both these spacings were 2 m taller than the two closest spacings (Table 1 and Fig. 1).

We selected the site index curve needed to drive the yield model from among several published curves (Golden *et al.*, 1981; Pienaar and Shiver, 1980; Smalley and Bower, 1971). None of the curves were entirely satisfactory, but this was to be expected since height growth patterns are known to be sensitive to the particular edaphic and environmental factors present




Fig. 1. Age trends in survival, average height, and average diameter by spacing for the Olinda plantation (filled symbols), and for plantations in the southeastern US (open symbols).

on a site, and the conditions at Olinda differ substantially from the conditions found in the Southeast. Qualitative comparisons of the site index models against the observed data indicated that the model of Smalley and Bower (1971) most closely approximated the height growth characteristics of the Olinda plantation and therefore was chosen to drive the yield model. The average height growth patterns of the Olinda plantation and the estimated average height-age curves for plantations in the Southeast are displayed in Fig. 1. The curves show that height growth of the Olinda plantation was reduced after about Age 11 relative to the Southeast, and remained below the Southeast curves as age increased. The 3.7 m Olinda spacing, however, fell within the range of the Southeast curves throughout the period of observation.

3.1.3. Diameter

Differences among spacings in mean stand diameter were evident four years after planting, when trees in the 3.7 m spacing were already 18% larger in DBH than in the 1.8 m spacing (Table 1) and Whitesell (1970). The relative growth advantage of the wider spaced trees had increased to 52.8% by the 20th year, dropping back to 32.6% at the end of the 34th year. During the nine years between ages 11 and 20 the 3.7 m spacing grew 7.0 cm, 59.1% more than the 1.8 m spacing which grew 4.4 cm. During the nine years between ages 25 and 34 growth of the 3.7 m spacing was only 3.9 cm, 18.7% less than the 1.8 m

spacing which had grown 4.8 cm. This may be an indication that competition in the 3.7 m spacing had become more intense with time relative to the 1.8 m spacing, which had developed a substantial degree of differentiation among trees, i.e., the coefficient of variation of mean DBH at Age 34 was 37.8% in the 1.8 m spacing versus 26.6% in the 3.7 m spacing. The spacing effect on diameter is further demonstrated by the divergence of the DBH growth curves. From the 4th to the 20th year, divergence increased progressively with increase in spacing, but from the 20th through the 25th year growth of the three widest spacings declined while growth of the 1.8 m spacing increased. From the 25th through the 34th year, growth increased again except for the 2.4 m spacing which, for no apparent reason, continued its decline (Fig. 1).

The yield model estimates of DBH (Fig. 1) indicate lower overall diameter growth rates for the Southeast. The difference between Olinda and the Southeast for any one spacing tended to be greater at all ages than the difference between that spacing and the next wider or closer spacing within the same location. Furthermore, the difference in DBH between locations increased as spacing increased. At Age 25, average stand diameters in Hawaii were 2.9-5.9 cm larger than for the same spacings in the Southeast.

The relative response of diameter growth of loblolly pine in the Southeast to increased spacing is of the same order of magnitude as in the Olinda spacings.

The maximum growth advantage of 3.7 m spacings over 1.8 m spacings was 39.7% at Age 20, dropping to 27.3% at Age 34. Between ages 11 and 20, 3.1 m spacings were estimated to grow 7.6 cm, 43.4% more than 1.8 m spacings which were estimated to grow 5.3 cm. Between the ages of 25 and 34, 3.7 m spacings were estimated to grow 3.3 cm, 18.2% less than the 1.8 m spacings which were estimated to grow 3.9 cm.

3.1.4. Stand growth trajectories

The relationship between mean DBH and stand density describes a trajectory that is characteristic of developing even-aged stands. As trees grow, stands advance through various stages along a typical path that ultimately attains a self-limiting maximum mean tree size-density boundary along which the stands then grow (Harms, 1984; Long and Smith, 1984; Oliver and Larson, 1990). The limiting density boundary can be equated to the self-thinning rule of Yoda et al. (1963) or the stand density index of Reineke (1933). Stand trajectories for the Olinda plantation and plantations in the Southeast are plotted in Fig. 2. The limiting density boundary reference line for Olinda was obtained by using a quadratic mean DBH of 25 cm and 2100 as the maximum (limiting) number of trees per ha possible at that DBH, estimated from a plot of the DBH-density data from the 1.8 m spacing. The line was located on the graph using the Reineke (1933) equation with a slope coefficient of 1.605, which Hasenauer et al. (1994) determined to be applicable to loblolly pine plantations in the southeastern US. The reference line for the Southeast was calculated using a mean DBH of 25 cm and 1150 trees/ha, the average of the values for maximum number of trees reported for coastal plain and Piedmont sites by Hasenauer et al. (1994). Our assumption in placing these reference lines was that the slope of the limiting density line was the same for both Olinda and the Southeast.

The difference in levels of the reference lines is a function of the higher stockability of the Olinda plantation as discussed by DeBell et al. (1989). They used the maximum mean tree size-density boundary line as a measure of stockability which they define as the maximum number of trees that can be grown to a given size under a particular set of conditions (DeBell et al., 1989; Harms et al., 1994)




Fig. 2. Mean stand DBH-density trajectories in relation to spacing for the Olinda plantation (filled symbols), and for plantations in the Southeastern US (open symbols). Ages (years) are shown for the DBH-density pairs plotted for each location. Estimated limiting density boundaries are shown for Olinda (dash), and for the Southeast (solid).

Following stand establishment, and for a period of years that varied with initial spacing, the trajectories rose vertically as trees grew in diameter (Fig. 2). As competition intensified and self-thinning commenced, the trajectories began to curve, the curvature increasing as the trajectories approached closer to the limiting density boundary. Self-thinning and mean DBH growth had progressed sufficiently by Age 20 to place the 1.8 m spacing trajectory at Olinda and the 1.8 m and 2.4 m spacing trajectories in the Southeast at their respective limiting density boundaries (Fig. 2). At Age 20, the mortality at the limiting density boundary at Olinda was 13% and there were 2601 trees/ha remaining with a mean DBH of 21.4 cm; the corresponding mortality in the Southeast was 42% with 1725 trees/ha remaining with an associated mean DBH of 18.5 cm. Self-thinning and diameter growth had not progressed enough by Age 34 in the 2.4, 3.0, and 3.7 m spacings at Olinda for their trajectories to have reached the limiting density boundary. In the




Fig. 3. Frequency distribution by percentage of trees per ha at age 34, above a minimum DBH by spacing, for the Olinda plantation and plantations in the southeastern US.

Southeast, the 3.0 m but not the 3.7 m trajectory had reached the limiting density boundary.

3.1.5. Size-class differentiation

Stand differentiation, which results from differential growth among the component trees (Oliver and Larson, 1990), was examined by evaluating DBH size-class frequency distributions by 5 cm class widths. Frequency distributions at Age 34 are graphed by spacing and location in Fig. 3 to show percentage number of trees per ha present in a spacing above a minimum diameter class. The Olinda spacings fall into two groups, with the smaller trees in the 1.8 m and 2.4 m spacings and the larger trees in the wider spacings. Southeastern US plantations show a uniform progression across spacings, due probably to the internal structure of the yield model. The greatest difference between the Olinda plantation and the Southeast is in the absolute range of diameter classes. There is a 5 cm difference in the minimum diameter class between locations-10 cm for Olinda and 15 cm

for the Southeast, but the maximum diameter class for the 3.7 m spacing at Olinda is 60 cm as compared to 40 cm for the same spacing in the Southeast. The smaller minimum diameter and greater range in tree size at any one spacing at Olinda further reflects greater tolerance of that system to crowding.

3.2. Stand productivity

3.2.1. Basal area

Net basal area per ha and mean annual basal area increment (MAI) at Olinda were greatest at the 1.8 m spacing to Age 34, and progressively less at successively wider spacings (Table 1, Figs. 4 and 5). Basal area increased continuously but at a gradually reducing rate throughout the 34 years of record. However, between the ages of 20 and 25 there was a period of substantial reduction in the rate of increase in all spacings, but most notably in the 1.8 m spacing in which the change was almost flat (Fig. 4). There was an initial rapid increase in basal area and associated MAI across all spacings. The differences in basal area among spacings widened until about Age 20, after which the rate of increase declined and the differences among spacings decreased. MAI culminated between ages 4 and 7 years in the two closest spacings and between ages 7 and 11 in the widest spacings, decreasing thereafter and beginning to converge (Fig. 5).

Basal area accumulation patterns in southeastern US plantations are similar to Olinda, but at lower absolute levels, through Age 11. Basal area MAI in the Southeast culminates later than at Olinda: at about Age 11 at 1.8 m spacings and between 11 and 20 at the wider spacings (Fig. 5). Between ages 11 and 20 years, the basal area accumulation and MAI curves for the 1.8 m spacing begin to fall below the 2.4 m spacing, and by Age 34 have crossed the curves for the wider spacings so that at Age 34 there is less basal area at 1.8 m spacing than at the 3.7 m spacing (Figs. 4 and 5). After Age 20, the basal area curves of all spacings are flattening and converging. By Age 34 curves of all spacings in the Southeast have converged to a point where they are within $3.5 \text{ m}^2/\text{ha}$ of each other in basal area and within $0.1 \text{ m}^2/\text{ha}/\text{year}$ in MAI. This was not the case at Olinda: at Age 34 years, curves of the Olinda spacings were still $20.1 \text{ m}^2/\text{ha}$ in basal area and $0.6 \text{ m}^2/\text{ha}/\text{year}$ in MAI from convergence, although




Fig. 4. Trends in stand basal area and volume by spacing for the Olinda plantation (filled symbols), and for plantations in the southeastern US (open symbols).




Fig. 5. Trends in mean annual increment in basal area and volume by spacing for the Olinda plantation (filled symbols), and for plantations in the southeastern US (open symbols).

they appeared to be in the early stages of drawing together (Figs. 4 and 5).

The convergence and crossing-over of basal area and volume curves for different spacings have been observed in other loblolly pine spacing studies, in natural and planted loblolly pine stands, and for other conifers (Buford, 1991; Liegel et al., 1985; Harrison and Daniels, 1988; Hafley et al., 1982). The phenomenon is probably due to a physiological inability of trees at older ages in closely spaced stands to rapidly take advantage of growing space released by mortality

to increase their growth. With continuing mortality this can result in a net loss in basal area.

3.2.2. Volume

Volume yields increased with decreasing spacing, with the greatest differences among the three closest spacings (Fig. 4). The 3.0 m and 3.7 m spacing yields were similar throughout the period of measurement, as was volume MAI after Age 20 (Fig. 5). Unlike basal area, however, the yield curves for volume showed an essentially linear and parallel increase, to Age 34.

There was no indication of convergence or a yield maximum (Fig. 4). Growth rate of the 1.8 m spacing was consistently greater than the wider spacings, but was essentially flat after Age 7. There was a period of reduced growth between ages 20 and 25 similar to that shown in the basal area curves. MAI of the 1.8 m spacing attained an apparent maximum at Age 20, decreasing to Age 25, and increasing slightly thereafter to Age 34. Growth rates of the other spacings had not culminated, the curves having taken on a broad, flat, and slightly increasing form (Fig. 5).

The reduction in volume and basal area growth of the Olinda plantation between ages 20 and 25 was probably a consequence of intense competition that developed as the stands approached and, in the case of the 1.8 m spacing, reached the limiting density boundary (Fig. 2). The severity of competition is reflected in the substantial increase in mortality during this period (Fig. 1). The growth recovery after Age 25 is probably a result of growth increases of the surviving trees as they occupied the space released by the trees that died.

Volume yield and MAI for the Southeast have patterns similar to the basal area trends. Growth and yield both converge by Age 34, and the curves for the 1.8 m spacing cross the wider spacings by Age 20 (Figs. 4 and 5). Volume MAI culminates later than basal area: the two closest spacings reach maximum volume growth at about Age 20, the 3.0 m spacing at Age 25, and the 3.7 m spacing probably at about Age 34.

Net volume and basal area and volume yields for the Olinda plantation and the Southeast at Age 34 are compared in Table 2. Both volume and basal area accumulations are greatest at the 1.8 m spacing at Olinda, and least at this spacing in the Southeast. Moreover, maximum yields at Olinda of the most

productive spacing (1.8 m) were about twice the maximum yields of the most productive spacing (3.0 m) in the Southeast at this age.

4. Discussion

Experimental studies of the influence of initial spacing on tree growth and plantation development have a long history (Sjolte-Jorgensen, 1967). In particular, the responses of loblolly pine trees in the southeastern US to initial spacing have been investigated and well documented in numerous spacing trials (Arnold, 1978; Harms and Lloyd, 1981; Owens, 1974; Shephard, 1974; Sprinz et al., 1979), and yield studies (Buford, 1991; Clutter et al., 1984; Hafley et al., 1982) among others. The patterns of tree and stand responses to initial spacing in Hawaii are consistent with spacing effects observed in the Southeast as depicted by the Hafley et al. (1982) yield model in Fig. 1. However, the nature of the responses differ substantially in magnitude. This was evident in the 1.8 m spacing at the limiting density boundaries where the difference between Hawaii and the southeastern US amounted to 62% more trees surviving and a 15% greater diameter at Age 34 years. This translates into 114% more basal area and 140% more volume (Table 2). When averaged across the four spacings, basal area and volume yields of the Olinda plantation at Age 34 were almost double (1.8 x) the yields in the Southeast.

It is evident from the data that the critical factors accounting for differences in yield between Olinda and the Southeast are tree survival and diameter growth. Tree height was not a factor because site index was a constant. The nature of the effects can

Table 2

Observed yields of the Olinda spacings at Age 34, Maui, Hawaii, versus estimated yields of loblolly pine plantations in the southeastern US (SEUS) of the same age, spacing, and site index ^a

Item	Spacing (m)							
	1.8 x 1.8		2.4 x 2.4		3.0 x 3.0		3.7 x 3.7	
	Olinda	SEUS	Olinda	SEUS	Olinda	SEUS	Olinda	SEUS
Trees per ha	1585	976	1362	897	915	756	725	613
Basal area (m ² ha ⁻¹)	103.1	48.2	96.5	51.7	85.2	51.7	83.0	49.6
Total volume, o.b. (m ³ ha ⁻¹)	1324	552	1191	612	1079	624	1053	605

^a SEUS yields are from Hafley et al. (1982) for site index 80 (24 m) base Age 25.

be seen by comparing stand development trajectories for the two locations (Fig. 2). The trajectories show that loblolly pine trees on the Olinda site were slow to thin relative to the Southeast but yet were able to continue to increase their diameters under conditions of crowding and competition that cause significant self-thinning and growth reduction in stands of comparable spacings and site index in the southeastern US. The result is that for each of the four spacings the mean DBH attained for a given number of surviving trees was always greater for the Olinda plantation. Since yield is the product of mean tree size and number of trees per unit area, the net effect of this was the nearly twofold yield differences observed.

Explanations for the high stockability of the Olinda plantation have been proposed by Harms et al. (1994) based on a study of the crown architecture of the trees and the structure of the stands at Age 26. They found a two-tiered crown-class structure with a sub-dominant stand having live crowns that extended from below the live crown base of the dominant stand well up into the main canopy. This served to increase the occupancy of the canopy space with functioning leaf area. The Olinda trees also had long crowns with high foliage biomass and leaf area. The development of these characteristics was attributed to the high solar radiation intensities and high sun angles at the latitude of Hawaii that allow penetration of sunlight deep into the canopy. In this light environment trees in the lower crown classes were able to retain their foliage and maintain a positive carbon balance sufficient for survival and growth in the densest stands. Growth rates were further enhanced by a long growing season, favorable temperature regime and soil moisture conditions, and a lack of disease and insect problems. In comparison, from measurements in a similar plantation in the Piedmont of South Carolina, they found crown development and growth to be limited by a less favorable light climate, a shorter growing season, and more stressful temperature and soil moisture conditions (Harms et al., 1994).

5. Conclusions

The Olinda trial serves to emphasize two important points. First, that the growth potential of loblolly pine is far greater than is apparent from growth records on

plantations in the southeastern US, a fact long since established by its performance at other locations in the sub-tropics. The data show that the growth of even unimproved stock when grown in very favorable environments, such as found in Hawaii, can double that which is currently being achieved in managed plantations in its natural range. (Burns and Hu, 1983; DeBell et al., 1989; Schultz, 1997).

Second, to capture the growth potential of loblolly pine, research must identify and quantify the specific tree, and stand physiological, and environmental factors that are responsible for the high survival rates and accompanying high diameter growth rates evident in the Olinda plantation at the high levels of competition that are not tolerated in plantations of the same spacing in the Southeast. And, conversely, research must identify the site-specific factors that are limiting to loblolly pine in the Southeast. Of the environmental and site factors thought to be responsible for the high stockability of the Olinda site by Harms et al. (1994), i.e., high sun angle, high solar radiation intensity, long growing season, favorable soil conditions, high foliage nutrients, and freedom from pests, most are not naturally present in non-limiting amounts in the Southeast. However, the potential for loblolly pine to respond to favorable environments suggests that improvement in productivity should be possible. That improvement is possible is supported in a recent report by Pienaar and Shiver (1993). Their data show that at Age 8, yield of genetically-improved loblolly pine planted on marginal agricultural cropland in the Georgia Piedmont was more than doubled by the simple expedient of complete control of the herbaceous competition during plantation establishment. It is evident from the Olinda plantation data that efforts to enhance the growth of loblolly pine should target research to improving productivity by discovering methods for increasing numbers of trees per unit area that can be grown to a desired size.

Acknowledgements

The authors wish to acknowledge the Hawaii Department of Land and Natural Resources, Division of Forestry and Wildlife, and the professional and technical staff of the Pacific Southwest Forest and Range Experiment Station for their assistance in the field. Special recognition is given to three pioneer

foresters who established this field trial: Karl Korte, District Forester; Gerald Pickford, Forest Ecologist; and Russell K. LeBarron, Silviculturist.

References

- Arnold, L.E., 1978. Gross yield of roughwood products from a 25-year-old loblolly and shortleaf pine spacing study. For. Res. Rep. No. 78-7. Agric. Exp. Sta. Univ. of Illinois at Urbana-Champaign, pp. 4.
- Baker, J.B., Langdon, O.G., 1990. *Pinus taeda* L. loblolly pine. In: Bums, R.M., Honkala, B.H. (Eds.), *Silvics of North America*, vol. 1, Conifers. U.S. Dep. Agric. Forest Serv. Agric. Handbk. 654, pp. 497-512.
- Baldanzi, G., 1978. Experimentacao com *Pinus taeda* na Estacao de Pesquisas Florestais de Rio Negro, Parana. (Research on *Pinus taeda* at the Rio Nigro Forest Research Station Parana). Floresta 9, 5-7.
- Buford, M.A., 1991. Performance of four yield models for predicting stand dynamics of a 30-year-old loblolly pine (*Pinus taeda* L.) spacing study. For. Ecol. Manage. 46, 23-38.
- Burkhart, H.E., 1977. Cubic foot volume of loblolly pine to any merchantable top diameter. South. J. Appl. For. 1, 7-9.
- Bums, P.Y., Hu, S.C., 1983. Ecological and silvicultural factors associated with rapid growth of loblolly pine stands. In: Jones, E.P., Jr. (Ed.), *Proceedings of the Second Biennial Southern Silvicultural Research Conference*, November 4-5, 1982, Atlanta, GA. U.S. Dep. Agric. Forest Serv., Southeastern For. Exp. Sta., Asheville, NC, Gen. Tech. Rep. SE-24, pp. 261-268.
- Clutter, J.L., Fortson, J.C., Pienarr, L.V., Brister, G.H., Bailey, R.L., 1983. *Timber Management: A Quantitative Approach*. John Wiley and Sons, New York, pp. 333.
- Clutter, J.L., Harms, W.R., Brister, G.H., Rheney, J.W., 1984. Stand structure and yields of site-prepared loblolly pine plantations in the lower coastal plain of the Carolinas, Georgia, and North Florida. U.S. Dep. Agric. Forest Serv. Southeastern For. Exp. Sta., Asheville, NC, Gen. Tech. Rep. SE-27, pp. 173.
- DeBell, D.S., Harms, W.R., Whitesell, C.D., 1989. Stockability: a major factor in productivity differences between *Pinus taeda* plantations in Hawaii and the Southeastern United States. For. Sci. 35, 708-719.
- Falkenhagen, E.R., 1978. Thirty five-year results from seven *Pinus elliotii* (Engelman.) and *Pinus Taeda* (L.) provenance trials in South Africa. S. African For. J. 107, 22-36.
- Foote, D.E., Hill, E.L., Nakamura, S., Stephens, F., 1972. Soil survey of the islands of Kauai, Oahu, Maui, Molokai, and Lanai, State of Hawaii. U.S. Dep. Agric. Soil Conservation Service, Washington, DC, pp. 232.
- Golden, M.S., Meldahl, R., Knowe, S.A., Boyer, W.D., 1981. Predicting site index for old-field loblolly pine plantations. South. J. Appl. For. 5, 109-114.
- Hafley, W.L., Smith, W.D., Buford, M.A., 1982. A new yield prediction model for unthinned loblolly pine in plantations. N.C. State Univ. Sch. For. Res., Raleigh, NC. Bioecon. Modeling Proj. Tech. Rep. 1, pp. 65.
- Harms, W.R., Lloyd, E.T., 1981. Stand structure and yield relationships in a 20-year-old loblolly pine spacing study. South. J. Appl. For. 5, 162-165.
- Harms, W.R., 1984. Applying silvics to stand management. In: Karr, B.L., Baker, J.L., Monaghan, T. (Eds.), *The Loblolly Pine Ecosystem (West Region) Proceedings*, March 20-22, 1984, Jackson, Mississippi. Miss. Coop. Extension Serv., Mississippi State, MS. pp. 3240.
- Harms, W.R., DeBell, D.S., Whitesell, C.D., 1994. Stand and tree characteristics and stockability in *Pinus taeda* plantations in Hawaii and South Carolina. Can. J. For. Res. 24, 511-521.
- Harrison, W.C., and Daniels, R.F., 1988. A new biomathematical model for growth and yield of loblolly pine plantations. In: Ek, A.R., Shifley, S.R., Burk, T.E. (Eds.), *Forest Growth Modeling and Prediction: Proceedings IUFRO Conference*, vol. 1, 23-27 August 1987, Minneapolis, MN. U.S. Dept. Agric. Forest Serv., North Central For. Exp. Sta., St. Paul MN, Gen. Tech Rep. NC-120, pp. 293-304.
- Hasenauer, H., Burkhart, H., Sterba, H., 1994. Variation in potential volume yield of loblolly pine plantations. For. Sci. 40, 162-178.
- Liegel, L.H., Balmer, W.L., Ryan, G.W., 1985. Honduras pine spacing study trial results in Puerto Rico. South. J. Appl. For. 9, 69-75.
- Long, J.N., Smith, F.W., 1984. Relation between size and density in developing stands: a description and possible mechanisms. For. Ecol. Manage. 7, 191-206.
- Machado, S., do, A., 1980. Curvas de indice de sitio para plantacoes de *Pinus taeda* L. na regio central do Estado do Parana. (Site index curves for *Pinus taeda* L. plantations in the central region of Parana State, Brazil. Floresta 11, 4-18.
- Machado, S., do A., Toledo Marquesini, C.P., Bacovis, R., 1990. Effects of age, site, average DBH and density upon the diameter distributions of thinned products from *Pinus taeda* plantation in the state of Parana, Brazil. Research in forest inventory, monitoring, growth and yield: proceedings from sessions of S4.02 Forest Resource Inventory and joint sessions of S4.02 and S4.01 Mensuration, Growth and Yield at the World Congress of International Union of Forestry Research Organizations in Montreal, Canada, August 5-11, 1990. School of For. and Wildlife Resour., Virginia Polytechnic Inst. and State Univ., Pub. FWS 3-90, pp. 150-165.
- Oliver, C.D., Larson, B.C., 1990. *Forest Stand Dynamics*. McGraw-Hill, Inc., New York, NY, pp. 467.
- Owens, E.G., 1974. The effects of initial tree spacing and regeneration method on stand development in loblolly pine after 10 years. In: Williston, H.L., Balmer, W.E. (Eds.), *Symposium on Management of Young Pines Proceedings*, 1974, Alexandria, La.-Oct. 22-24, Charleston, S.C.-Dec. 3-5. U.S. Dept. of Agric. Forest Serv. Southern and Southeastern Forest Exp. Sta., pp. 340-349.
- Pienaar, L.V., Shiver, B.D., 1980. Dominant height growth and site index curves for loblolly pine plantations in the Carolina flatwoods. South. J. Appl. For. 4, 54-59.
- Pienaar, L.V., Shiver, B.D., 1993. Early results from an old-field loblolly pine spacing study in the Georgia piedmont with competition control. South. J. Appl. For. 17, 193-196.

- Reineke, L.H., 1933. Perfecting a stand-density index for even-aged forests. *J. Agric. Res.* 46, 627-638.
- Schultz, R.P., 1997. Loblolly Pine. The ecology and culture of loblolly pine (*Pinus taeda* L.). U.S. Dep. of Agric. Forest Serv. Agric. Handbk 713.
- Shepherd, R.K., Jr., 1974. An initial spacing study. In: Williston, H.L., Balmer, W.E. (Eds.), Symposium on Management of Young Pines Proceedings, 1974, Alexandria, La. -Oct. 22-24, Charleston, S.C.-Dec. 3-5. U.S. Dept. of Agric. Forest Serv. Southern and Southeastern Forest Exp. Sta., pp. 121-128.
- Sjolte-Jorgensen, J., 1967. The influence of spacing on the growth and development of coniferous plantations. *International Rev. Forest Res.*, vol. 2, Academic Press, New York, NY, pp. 43-94.
- Smalley, G.W., Bower, D.R., 1971. Site index curves for loblolly and shortleaf pine plantations on abandoned fields in Tennessee, Alabama and Georgia highlands. U.S. Dep. Agric. Forest Serv., Southern For. Exp. Sta., New Orleans, LA, Res. Note SO-126, pp. 4.
- Sprinz, P., Clason, T., Bower, D., 1979. Spacing and thinning effects on the growth and development of a loblolly pine plantation. In: Clason, T.R. (Ed.), For. Res. Rep., North Louisiana Hill Farm Exp. Sta., Homer, LA, pp. 1-42.
- Strub, M.R., Breckenkamp, B.V., 1985. Carrying capacity and thinning response of *Pinus taeda* in the CCT experiments. *S. African J. For.* 128, 6-11.
- Whitesell, C.D., 1970. Early effects of spacing on loblolly pine in Hawaii. U.S. Dep. Agric. Forest Serv., Pacific Southwest For. and Range Exp. Sta., Berkeley, CA, Res. Note PSW-223, pp. 3.
- Whitesell, C.D., 1974. Effects of spacing on loblolly pine in Hawaii after 11 years. U.S. Dep. Agric. Forest Serv., Pacific Southwest For. and Range Exp. Sta., Berkeley, CA, Res. Note PSW-295, pp. 4.
- Yoda, K., Kira, T., Ogawa, H., Hozumi, K., 1963. Self-thinning in overcrowded pure stands under cultivated and natural conditions. *J. Biol. Osaka City Univ.* 14, 107-129.

Forest Ecology and Management

Submission of manuscripts.

Manuscripts should be submitted in triplicate. Authors from the Americas, Australia, New Zealand and the Pacific are requested to send their manuscripts to *Forest Ecology and Management*, Dr. Richard F. Fisher, Department of Forest Science, Texas A&M University, College Station, TX77843-2135, USA; all other authors are requested to send their manuscripts to the Editorial Office, *forest Ecology and Management*, PO. Box 181, 1000 AD Amsterdam, The Netherlands.

Authors in Japan please note: Upon request, Elsevier Science Japan will provide authors with a list of people who can check and improve the English of their paper (before submission). Please contact our Tokyo office: Elsevier Science Japan, 1-9-15 Higashi-Azabu, Minato-ku, Tokyo 106-0044, Japan; tel. (+81)-3-5561-5033; fax (+81)-3-5561-5047.

Electronic manuscripts: Electronic manuscripts have the advantage that there is no need for the rekeying of text, thereby avoiding the possibility of introducing errors and resulting in reliable and fast delivery of proofs.

For the initial submission of manuscripts for consideration, hardcopies are sufficient. For the processing of *accepted papers*, electronic versions are preferred. After *final acceptance*, your disk plus two, final and exactly matching printed versions should be submitted together. Double density (DD) or high density (HD) diskettes (3.5 or 5.25 inch) are acceptable. It is important that the file saved is in the native format of the wordprocessor program used. Label the disk with the name of the computer and wordprocessing package used, your name, and the name of the file on the disk. Further information may be obtained from the Publisher.

Enquiries concerning manuscripts and proofs: questions arising after acceptance of the manuscript, especially those relating to proofs, should be directed to Elsevier Science Ireland Ltd., Elsevier House, Brookvale Plaza, East Park, Shannon, Co. Clare, Ireland; Tel. (+353-61) 709685; Fax: (+353-61) 709113.

Advertising information: Advertising orders and enquiries can be sent to: **USA, Canada and South America:** Mr Tino de Carlo, The Advertising Department, Elsevier Science Inc., 655 Avenue of the Americas, New York, NY 10010-5107, USA; phone: (+1) (212) 633 3815; fax: (+1) (212) 633 3820; e-mail: t.decarlo@elsevier.com **Japan:** The Advertising Department, Elsevier Science K.K., 9-15 Higashi-Azabu 1-chome, Minato-ku, Tokyo 106, Japan; phone: (+81) (3) 5561 5033; fax: (+81) (3) 5561 5047. **Europe and ROW:** Rachel Gresle-Farthing, The Advertising Department, Elsevier Science Ltd., The Boulevard, Langford Lane, Kidlington, Oxford OX5 1GB, UK; phone: (+44) (1865) 843565; fax: (+44) (1865) 843976; e-mail: r.gresle-farthing@elsevier.co.uk

Orders, claims, and product enquiries: please contact the Customer Support Department at the Regional Sales Office nearest you:

New York: Elsevier Science, PO Box 945, New York, NY 10159-0945, USA; Tel.: (+1) (212) 633 3730 [toll free number for North American customers: 1-888-4ES-INFO (437-4636)]; Fax: (+1) (212) 633 3680; E-mail: usinfo-f@elsevier.com

Amsterdam: Elsevier Science, PO Box 211, 1000 AE Amsterdam, The Netherlands; Tel.: (+31) 20 4853757; Fax: (+31) 20 4853432; E-mail: nlinfo-f@elsevier.nl

Tokyo: Elsevier Science, 9-15, Higashi-Azabu 1-chome, Minato-ku, Tokyo 106-0044, Japan; Tel.: (+81) (3) 5561 5033; Fax: (+81) (3) 5561 5047; E-mail: info@elsevier.co.jp

Singapore: Elsevier Science, No. 1 Temasek Avenue, #17-01 Millenia Tower, Singapore 039192; Tel.: (+65) 434 3727; Fax: (+65) 337 2230; E-mail: asiainfo@elsevier.com.sg

Rio de Janeiro: Elsevier Science, Rua Sete de Setembro 111/16 Andar, 20050-002 Centro, Rio de Janeiro - RJ, Brazil; Tel.: (+55) (21) 509 5340; Fax: (+55) (21) 507 1991; E-mail: elsevier@campus.com.br [Note (Latin America): for orders, claims and help desk information, please contact the Regional Sales Office in New York as listed above]

USA mailing info: *Forest Ecology and Management* (0378-1121) is published semi-monthly by Elsevier Science B.V. (PO. Box 211, 1000 AE Amsterdam, The Netherlands). Annual subscription price in the USA US\$ 2488.00 (valid in North, Central and South America), including air speed delivery. Application to mail at periodical postage rate is pending at Jamaica, NY 11431.

USA POSTMASTER: Send address changes to *Forest Ecology and Management*, Publications Expediting Inc., 200 Meacham Avenue, Elmont, NY 11003.

AIRFREIGHT AND MAILING in the USA by Publications Expediting Inc., 200 Meacham Avenue, Elmont, NY 11003.

***Forest Ecology and Management* has no page charges**

For a full and complete Guide for Authors, please refer to *Forest Ecology and Management*, Vol. 126, No. 1, pp. 87-90. The instructions can also be found on the World Wide Web: access under <http://www.elsevier.nl> or <http://www.elsevier.com>

LANDSCAPE AND URBAN PLANNING

An International Journal of Landscape Ecology,
Landscape Planning and Landscape design

Editor-in-Chief:

J.E. Rodiek, *College of Architecture, Texas A & M University, College Station, TX 77843-3137, USA*

AIMS AND SCOPE

A journal concerned with conceptual, scientific and design approaches to land use. By emphasizing ecological understanding and a multi-disciplinary approach to analysis and planning and design, it attempts to draw attention to the interrelated nature of problems posed by nature and human use of land. In addition, papers dealing with ecological processes and interactions within urban areas, and between these areas and the surrounding natural systems which support them, will be considered. Papers in which specific problems are examined are welcome. Topics might include but are not limited to landscape ecology, landscape planning and landscape design. Landscape ecology examines how heterogeneous combinations of ecosystems are structured, how they function and how they change. Landscape planning examines the various ways humans structure their land use changes. Landscape design involves the physical strategies and forms by which land use change is actually directed. *Landscape and Urban Planning* is based on

the premise that research linked to practice will ultimately improve the human made landscape.

Editorial Advisory Board:

I.D. Bishop, *Parkville, Vic., Australia*,
E.G. Bolen, *Wilmington, NC, USA*,
I.D. Bruns, *Schorndorf, Germany*,
J.B. Byrom, *Edinburgh, UK*,
T.C. Daniel, *Tucson, AZ, USA*,
R.M. DeGraaf, *Amherst, MA, USA*,
J.G. Fabos, *Amherst, MA, USA*,
S. Gonzalez Alonso, *Madrid, Spain*,
M. Hough, *Etobicoke, ON, Canada*,
P. Jacobs, *Montreal, PQ, Canada*,
D.S. Jones, *Melbourne, Vic., Australia*,
H. Lavery, *Milton, Qld., Australia*,
W.M. Marsh, *Flint, MI, USA*,
D.L. Mitchell, *Dallas, TX, USA*,
D.G. Morrison, *Athens, GA, USA*,
J.I. Nassauer, *St. Paul, MN, USA*,
M. Nelischer, *Guelph, ON, Canada*,
D.D. Paterson, *Vancouver, BC, Canada*,
A. Ramos, *Madrid, Spain*,
P. Shepard, *Claremont, CA, USA*,
O.R. Skage, *Alnarp, Sweden*,
R.C. Smardon, *Syracuse, NY, USA*,
G. Sorte, *Alnarp, Sweden*,
F. Stearns, *Rhineland, WI, USA*,

R.C. Szaro, *Washington, DC, USA*,
J.W. Thomas, *La Grande, OR, USA*,
P.J. Trowbridge, *Ithaca, NY, USA*,
T.H.D. Turner, *London, UK*,
M.J. Vroom, *Wageningen, The Netherlands*,
W.V. Wendler, *College Station, TX, USA*,
B.-E. Yang, *Seoul, Korea*,
E.H. Zube, *Tucson, AZ, USA*

ABSTRACTED/INDEXED IN
Applied Ecology Abstracts,
Biological Abstracts,
Current Contents B & S,
Environmental Periodicals
Bibliography, Geobase,
Geographical Abstracts,
LandSearch.

1994 SUBSCRIPTION DATA

Volumes **28-30** (in 9 issues)
Subscription price:
Dfl. 1080.00 (US \$584.00)
Incl. Postage
ISSN 0169-2046

Elsevier Science Publishers,
P.O. Box 211, 1000 AE Amsterdam,
The Netherlands
Fax: (020) 5803-203

Customers in the USA and Canada:
Elsevier Science Publishers
P.O. Box 945, Madison Square Station,
New York, NY 10160-0757, USA
Fax: (212) 633-3680

The Dutch Guildler price quoted applies worldwide, except in the Americas (North, Central and South America). The US Dollar price quoted applies in the Americas only. Non VAT registered customers in the European Community should add the appropriate VAT rate applicable in their country to the price



ELSEVIER
SCIENCE PUBLISHERS