

Invasion of exotic earthworms into ecosystems inhabited by native earthworms

P. F. Hendrix · G. H. Baker · M. A. Callaham Jr ·
G. A. Damoff · C. Fragoso · G. González ·
S. W. James · S. L. Lachnicht ·
T. Winsome · X. Zou

Published online: 1 September 2006
© Springer Science+Business Media B.V. 2006

Abstract The most conspicuous biological invasions in terrestrial ecosystems have been by exotic plants, insects and vertebrates. Invasions by exotic earthworms, although not as well studied, may be increasing with global commerce in agriculture, waste management and bioremediation. A number of cases has documented where invasive earthworms have caused significant changes in soil profiles, nutrient and organic matter

dynamics, other soil organisms or plant communities. Most of these cases are in areas that have been disturbed (e.g., agricultural systems) or were previously devoid of earthworms (e.g., north of Pleistocene glacial margins). It is not clear that such effects are common in ecosystems inhabited by native earthworms, especially where soils are undisturbed. We explore the idea that indigenous earthworm fauna and/or characteristics of their

P. F. Hendrix (✉) · T. Winsome
Institute of Ecology and Department of Crop & Soil
Sciences, University of Georgia, Athens,
GA 30602, USA
e-mail: hendrixp@uga.edu

G. H. Baker
CSIRO Entomology, GPO Box 1700, Canberra,
ACT 2601, Australia

M. A. Callaham Jr
USDA Forest Service, University of Georgia, Athens,
GA 30602, USA

G. A. Damoff
Arthur Temple College of Forestry, Stephen F.
Austin State University, Nacogdoches, TX 75962,
USA

C. Fragoso
Departamento Biología de Suelos, Instituto de
Ecología A.C., Km. 2.4 Carr. Antigua a Coatepec,
No. 351, Col. Congregación el Haya. AP 63,
Xalapa 91070 Veracruz, Mexico

G. González
USDA Forest Service, International Institute of
Tropical Forestry, 1201 Ceiba Street, Río Piedras,
PR 00926, USA

S. W. James
Natural History Museum and Biodiversity Research
Center, The University of Kansas, 1345 Jayhawk Blvd,
Lawrence, KS 66045-7163, USA

S. L. Lachnicht
USDA ARS NCSCRL, 803 Iowa Ave., Morris,
MN 56267, USA

X. Zou
Institute for Tropical Ecosystem Studies, University
of Puerto Rico, PO Box 23341, San Juan, PR 00931,
USA

native habitats may resist invasion by exotic earthworms and thereby reduce the impact of exotic species on soil processes. We review data and case studies from temperate and tropical regions to test this idea. Specifically, we address the following questions: Is disturbance a prerequisite to invasion by exotic earthworms? What are the mechanisms by which exotic earthworms may succeed or fail to invade habitats occupied by native earthworms? Potential mechanisms could include (1) intensity of propagule pressure (how frequently and at what densities have exotic species been introduced and has there been adequate time for proliferation?); (2) degree of habitat matching (once introduced, are exotic species faced with unsuitable habitat conditions, unavailable resources, or unsuited feeding strategies?); and (3) degree of biotic resistance (after introduction into an otherwise suitable habitat, are exotic species exposed to biological barriers such as predation or parasitism, “unfamiliar” microflora, or competition by resident native species?). Once established, do exotic species co-exist with native species, or are the natives eventually excluded? Do exotic species impact soil processes differently in the presence or absence of native species? We conclude that (1) exotic earthworms do invade ecosystems inhabited by indigenous earthworms, even in the absence of obvious disturbance; (2) competitive exclusion of native earthworms by exotic earthworms is not easily demonstrated and, in fact, co-existence of native and exotic species appears to be common, even if transient; and (3) resistance to exotic earthworm invasions, if it occurs, may be more a function of physical and chemical characteristics of a habitat than of biological interactions between native and exotic earthworms.

Keywords Native earthworms · Exotic earthworms · Biological invasions · Disturbance · Competition

Introduction

Research over the past century has shown that where earthworms are abundant, they significantly

influence soil processes and are integral to the functioning of terrestrial ecosystems. Documented effects of earthworms include accelerated plant litter decomposition, nutrient transformations and plant nutrient uptake; increased soil aggregation and porosity; and enhanced water infiltration and solute transport (see Satchell 1983; Lee 1985; Hendrix 1995; Edwards and Bohlen 1996; Lavelle et al. 1999; Edwards 2004). While these effects are usually considered desirable in agricultural soils, recent interest has focused on detrimental impacts of invasive, exotic earthworms on soil processes in wildland ecosystems (Hendrix and Bohlen 2002; Bohlen et al. 2004a,b; James and Hendrix 2004). Exotic earthworms are capable of significantly affecting soil profiles, nutrient and organic matter dynamics, other soil organisms, and plant communities. Impacts have been reported in tropical forests (Zou and González 1997; Zou and Bashkin 1998; González and Zou 1999; Fragoso et al. 1999; Liu and Zou 2002; Decaëns et al. 2004); chaparral shrublands (Graham and Wood 1991; Graham et al. 1995); grasslands (Stockdill 1982; James 1991; Callahan et al. 2001); and particularly in temperate forests (Langmaid 1964; Alban and Berry 1994; Scheu and Parkinson 1994; Steinberg et al. 1997; Burte-low et al. 1998; McLean and Parkinson 2000; González et al. 2003; Bohlen et al. 2004b; Hale et al. 2005; also see Frelich et al. this issue).

Most of the work on earthworm invasions has focused on a relatively few species (e.g., European lumbricids, *Amyntas* spp., *Pontoscolex corethrus*) that have achieved wide distributions and are now abundant in many ecosystems. Moreover, the most dramatic effects of exotic species on soil organic matter dynamics have been observed in areas previously uninhabited by earthworms (e.g., north of Pleistocene glacial margins; see Frelich et al., McLean et al., Migge-Kleian et al. and Tiunov et al., this issue) or where native populations have been reduced by disturbance (e.g., pastures in Australia and Puerto Rico; see Baker et al. and González et al. in this issue). Effects of exotic earthworms have not been as frequently reported from invasions of ecosystems inhabited by native earthworm assemblages where soils and vegetation are undisturbed (e.g., Abbott 1985; Kalisz and Dotson 1989; Lavelle and Pashanas

1989; Callaham and Blair 1999; Fragoso et al. 1995, 1999). These observations suggest that some characteristics of indigenous earthworm fauna and/or their native habitats may be resistant to invasion by exotic earthworm species and thus may reduce the impact of exotic species on soil processes. In this paper, we explore the interactions between native and exotic earthworms and factors that may facilitate or inhibit invasions by exotic species into areas inhabited by native earthworm species.

Interactions between native and exotic earthworms

Habitat disturbance or competitive exclusion

Exotic earthworms have been spread throughout the world, aided by human colonization and commerce for at least the past few centuries; several peregrine species are now prevalent in many soils impacted by human activity (Ljungstrom 1972; Lee 1985; Kalisz 1993; Fragoso 1995, 1999; Reynolds 1995; Bhadauria et al. 2000). Where these introductions have occurred in areas inhabited by indigenous earthworms, exotic earthworms may not persist, they may occur exclusively, or they may co-occur with the native earthworm species. Reasons for success or failure of establishment, or for varying densities of exotic species at any particular site may not be known with certainty, but probably include site characteristics (e.g., soil and climatic conditions), invasion history (e.g., frequency and duration of introductions), and characteristics of the exotic and native species involved. Site disturbance, including natural phenomena (e.g., tree fall, floods) that can create conditions favorable for establishment or proliferation of exotic species, may be a particularly important factor.

Since the time of early observations, mechanisms by which exotic earthworms come to dominate in certain ecosystems have been debated (Eisen 1900; Beddard 1912; Smith 1928; Lee 1961; Stebbings 1962). Do exotic species displace native species through direct or indirect competition, or do exotic species occupy vacant niches following disturbance and the demise of

native species? Kalisz and Wood (1995) summarized the prevailing idea that physical disturbance or habitat fragmentation are prerequisite to establishment of and domination by exotic earthworms in soils occupied by native species. The proposed sequence is (a) habitat disturbance, (b) decline or extirpation of native species, (c) introduction of exotic species, and (d) colonization of empty habitat by exotic species. By considering the currently observed state of any particular earthworm assemblage, we can trace several possible series of events that may have led to that state from a presumed indigenous community in a pristine ecosystem (Fig. 1).

Pathway A represents the extreme case described by Kalisz and Wood (1995), through which disturbance leads to exclusively exotic assemblages, as often observed with “anthropochorous” earthworms in agricultural soils (e.g., Parmelee et al. 1990; Baker et al. 2002). We can speculate that the same outcome may occur under less severe disturbance but perhaps with more aggressive exotic invaders, as in pathway B-1. Pathways B-2 and C-1 lead to the often observed co-occurrence of native and exotic species (Stebbins 1962; Abbott 1985; James 1991; Fragoso et al. 1999) through varying levels of habitat disturbance and invasion intensity. The B-2 case again assumes at least moderate levels of disturbance, which reduce native population density and alter habitat conditions prior to invasion. The C-1 pathway suggests that competitive displacement of native species by exotic species may occur even in relatively undisturbed ecosystems; this possibility, whereby forest fragmentation for example, may foster exotic invasions without direct habitat disturbance, was termed “invisible disturbance” by Kalisz and Wood (1995). The idea is controversial and is supported by little empirical data. Furthermore, whether co-occurrence is a stable condition or whether native or exotic species maintain dominance in any particular situation are interesting long-term questions, as noted by the question marks for “successful” invasion on these pathways in Fig. 1. Finally, pathway C-2 represents the idea that native earthworm assemblages or properties of their minimally disturbed habitats are resistant to invasion by exotic species. There




Fig. 1 Hypothesized sequences of invasion depending on degree of habitat disturbance and invasion success by exotic earthworms invading ecosystems inhabited by native earthworms. See text for description of pathways

also is very little information with which to test this idea, as discussed below.

Case studies

As noted earlier, much of the research on earthworm invasions has been conducted in ecosystems previously devoid of earthworms. However, a few studies have examined interactions between native and exotic earthworms (Table 1).

Abbott (1985) in Western Australia and Kalisz and Dotson (1989) in Kentucky, USA, found that exotic earthworms occurred only in severely disturbed forest sites, whereas native earthworms occurred in undisturbed to slightly disturbed sites, sometimes in association with exotics. They noted that the exotics had failed to disperse into undisturbed areas even decades after introduction. Dalby et al. (1998) concluded from microcosm studies that the European lumbricid, *Aporrectodea longa*, would not successfully invade forest soils inhabited by native megascolecid earthworms in South Australia because of its strong preference for nearby pasture soils. In the central Himalayas of India, Bhadauria et al. (2000) reported declines in endemic earthworm abundances with disturbance pressure in natural and regenerating forests; exotic species did occur in the undisturbed climax forest but certain exotic species were restricted to the regenerating forest.

Studies of an invasion of European lumbricids into a native prairie in Kansas, USA, suggested

competitive displacement of native *Diplocardia* spp. by exotic *Aporrectodea* spp. in disturbed areas, but continued dominance by the native species under natural conditions (James 1982; Callaham and Blair 1999); regular prescribed fires appeared to favor the diplocardians. Studies in California, USA, grasslands (Winsome 2003; Winsome et al. 2006) showed that exotic species predominated only in areas within fertilizer-amended pastures and on sedimentary soils within unamended pastures. Native species were present in abundance equal to or greater than exotic species in all other habitat types, including oak woodland reserves within the amended pastures and on serpentine soils. These results suggested that displacement of native by exotic species in these grasslands occurred only where resource quality and/or disturbance were at a maximum (i.e., presence of high-quality forage grasses in the heavily grazed, amended pastures). Damoff (2005) found co-occurrence of the exotic *Amyntas diffringens* with several native earthworm species in a secondary bottomland hardwood forest in eastern Texas, USA. *Diplocardia komareki* was the largest and deepest-burrowing of the native species and may have interacted least with *A. diffringens*; all other species appeared to occupy the same vertical position (large niche overlap) in the soil profile with *A. diffringens*.

In Puerto Rico, the exotic earthworm, *Pontoscolex corethrurus*, was found to dominate both disturbed and little-disturbed sites, whereas native

Table 1 Relative occurrence of native and exotic earthworm in ecosystems subjected to varying types and degrees of disturbance. Minus, single plus and double plus signs indicate absence, presence and dominance, respectively, of native or exotic earthworm species

Earthworms					
Location	Ecosystem type	Disturbance	Native Exotic	Earthworm species (native/exotic)	References
Western Australia	Eucalyptus forest	Minimal	+ -	“Indigenous” species/ <i>Aporrectodea trapezoides</i> ,	Abbott (1985)
		Forest clearing/replacement	+ +	<i>A. caliginosa</i> , <i>Eisenia fetida</i> , <i>Octolasion cyaneum</i> ,	
	Pasture	Converted	+ ++	<i>Microscolex dubius</i>	Mele and Carter (1999a, b)
South Eastern Australia	Crops Acidified crops	Converted	+ +	<i>Spenceriella</i> spp./	Mele and Carter (1999a, b)
		Converted	+ +	<i>A. caliginosa</i> , <i>A. trapezoides</i> ,	
		Converted	+ +	<i>A. tuberculata</i> , <i>A. rosea</i> , <i>O. cyaneum</i> , <i>L. rubellus</i> , <i>M. dubius</i> , <i>M. phosphoreus</i> , <i>Microscolex</i> spp.	
South Australia	Eucalyptus forest	Minimal	++ +	<i>Gemascocox lateralis</i> / <i>A. caliginosa</i> , <i>A. trapezoides</i>	Dalby et al. (1998)
India	Broad-leaf temperate forest Sublimax mixed forest Pasture/pine plantation	Undisturbed	++ +	<i>Drauidia</i> sp. <i>Eutyphoeus naniamu</i> , <i>Eu. festivus</i> ,	Bhadauria et al. (2000)
		Minimal	+ +	<i>Eu. waltonii</i> / <i>Amynthas corticis</i> , <i>Bimastos parvus</i> ,	
		Deforested/converted	+ ++	<i>O. tyrtaeum</i> , <i>Octochaetona beatrix</i>	
Ivory Coast	Savanna/gallery forest Pasture Agroecosystems	Minimal	+ -	13 spp., esp. <i>Chuniodrillus</i>	Lavelle (1978), Fragoso et al. (1999)
		Converted	+ -	spp., <i>Milsonia anomala</i> ,	
		Converted	+ -	<i>Dichogaster agilis</i> , <i>Dichogaster</i> spp./None	
Colombia	Savanna	Minimal	+ -	Indigenous acanthodrilids,	Jiménez et al. (1998, 2001)
		Grazed pasture	+ -	glossoscolectids, ocnerodrilids	
South Eastern Mexico	Tropical rainforest Tropical deciduous forest forest cloud forest	Minimal	++ +	69 native spp, esp.	Fragoso et al. (1995, 1999)
		Minimal	++ +	<i>Dichogaster bolau</i> ,	
		Minimal	++ +	<i>D. saliens</i> , <i>Balanteodrilus pearsei</i> , <i>Lavellodrillus parvus</i> , <i>Diploctrema murchieii</i> /26 exotic spp., esp. <i>Pontoscolex corethrusus</i> , <i>Polypheretima elongata</i>	

Table 1 continued

Location	Ecosystem type	Disturbance	Native	Exotic	Earthworm species (native/exotic)	References
	Crops	Deforested/converted	+	++		
	Tropical pasture	Deforested/converted	+	++		
	Tree plantation	Deforested/converted (exotic abundance increases with level of disturbance)	+	++		
Costa Rica	Tropical pasture (non-grazed)	Deforested/converted	+	++	<i>Glossodrilus nemoralis</i> / <i>P. corethrurus</i> , <i>Metaphire californica</i>	Fragoso et al. (1999)
Peru	Tropical rainforest	Minimal	++	+	“Native forest spp.”/	Lavelle and Pashanasi (1989), Fragoso et al. (1999)
	Agroecosystems	Deforested/converted	+	++	<i>P. corethrurus</i>	
Puerto Rico	Tropical rainforest	Minimal	++	+	<i>Estherella</i> sp., <i>E. gatesi</i> ,	Zou and Gonzalez (1997),
	Tropical cloud forest	Minimal	++	+	<i>E. montana</i> , <i>Trigaster</i>	González and Zou
	Tropical pasture	Deforested/converted	+	+	<i>longissimus</i> / <i>P. corethrurus</i> , <i>Amyntas rodericensis</i>	(1999), Hendrix et al. (1999a), Liu and Zou (2002), Lachnicht et al. (2002)
	Pine and mahogany plantations	Deforested/converted	–	+	None/ <i>P. corethrurus</i> ,	González et al. (1996)
	Secondary forests	Deforested/converted	+	+	<i>A. rodericensis</i> <i>Pontoscolex spiralis</i> , <i>E.gatesi</i> , <i>E.montana</i> / <i>P. corethrurus</i> ,	
					<i>A. rodericensis</i>	
Kentucky, USA	Deciduous forest	Minimal	+	–	<i>Bimastos</i> spp., <i>Diplocardia</i> spp.	Kalisz and Dobson
		Logged	+	+	<i>Eisenoides carolinensis</i> ,	(1989), Dobson and
		Severely disturbed	–	+	<i>Komarektonia etoni</i> / <i>A.</i>	Kalisz (1989), Kalisz
					<i>Allolobophora chloroica</i> ,	(1993), Kalisz and
					<i>Ap.</i>	Wood (1995)
Georgia, USA	Mixed forest	Moderate	+	++	<i>Bimastos</i> sp., <i>Diplocardia</i> sp./ <i>A.</i>	Parmelee et al. (1990),
		Deforested/cultivated	+	++	<i>caliginosa</i> , <i>Lumbricus</i>	Hendrix et al. (1992),
					<i>rubellus</i> , <i>L. terrestris</i> , <i>M.</i>	Callaham and Hendrix
					<i>dubius</i>	(1997)
Florida, USA	Longleaf pine/wiregrass savanna	Minimal	+	–	<i>Diplocardia</i> spp.	Hendrix et al. (1999b)
Kansas, USA	Tallgrass prairie	Minimal	+	–	<i>Bimastos welchi</i> , <i>Diplocardia</i>	James (1991), Callaham
		Fire suppression	+	+	spp./ <i>A. caliginosa</i> , <i>O.</i>	and Blair (1999), Calla-
		Conversion to pasture	+	++	<i>cyaneum</i> ,	ham et al. unpublished
	Gallery forest	Encroachment into prairie	+	++		

Table 1 continued

Earthworms		Location		Ecosystem type	Disturbance	Native	Exotic	Earthworm species (native/exotic)	References									
Missouri, USA	Riparian alluvium Mixed forest	Minimal	++	+	+	+	+	<i>Bimastos zeteki</i> , <i>Diplocardia</i> spp./ <i>A. trapezoides</i> , <i>O. lacteum</i>	Stebbins (1962)									
										Texas, USA	Bottomland hardwood forest	Moderate	+	+	+	+	<i>Diplocardia</i> spp./ <i>Amyntas diffringens</i>	Damoff (2005)
Oregon, USA	Oak savanna Pasture Temperate coniferous forest	Minimal Converted/fertilized Minimal Logging/campsites	+	+	-	-	<i>Argilophilus marmoratus</i> / <i>A. trapezoides</i> Indigenous Megascolecids/ <i>A. caliginosa</i>	Winsome et al. (2003) Hendrix and Cromack unpublished										

earthworms were present in undisturbed sites. Exotic earthworms occurred in mahogany and pine plantations as well as in naturally regenerated secondary forests; native species were only present in the secondary forests (González et al. 1996). *Pontoscolex corethrurus* also was present in the relatively undisturbed tabonuco forest (with selective logging) and cloud forest at the top of undisturbed Luquillo Mountains (Zou and González 1997; Liu and Zou 2002; Hendrix et al. 1999b; also see González et al. in this issue).

A survey of 84 cropping and pasture systems in southeastern Australia showed that exotic species were dominant, but a single native species co-occurred with exotics in both systems (Mele and Carter 1999a). The native species occurred with higher abundances in the less disturbed pasture systems, but was also the dominant species in acidified cropping systems (Mele and Carter 1999a, b). An examination of earthworm populations under different tillage or stubble management showed that less disturbance (no-tillage verses plowing) favors greater earthworm populations, and can preserve native species in both temperate (Parmelee et al. 1990; Mele and Carter 1999b) and tropical (Fragoso et al. 1999) ecosystems (also see Baker et al. and González et al. in this issue). In eastern Colombia, exotic earthworms were apparently excluded from native savannas converted into man-made pasture systems because native conditions were relatively maintained (Jiménez et al. 1998).

Finally, recent studies using stable isotopic techniques in a variety of ecosystems suggest the potential for direct competition between native and exotic species for food resources, based on overlap in ¹³C and ¹⁵N signatures (Hendrix et al. 1998, 1999a, b; Callaham et al. 2001; Lachnicht et al. 2002; Winsome 2003). These relatively short-term studies do not show actual displacement of native species; longer-term observations or studies of well-characterized chronosequences might be more conclusive.

Overall, these case studies suggest that exotic earthworms are able to invade and become established in a variety of ecosystems currently or previously inhabited by native earthworms. They also appear to co-occur with native earthworms in at least some, usually disturbed, conditions. As

discussed below, it is not clear if co-occurrence is persistent or only a transient situation.

Invasion resistance by native earthworms and their habitats

A number of general mechanisms have been proposed that explain why exotic species may succeed or fail to invade new habitats (Simberloff 1989; Williamson 1996; Mack et al. 2000). Specific to earthworms, Hendrix and Bohlen (2002) discuss several mechanisms that may be particularly important determinants of success or failure along the C-2 pathway in Fig. 1.

Propagule pressure

In any area vulnerable to invasion, it is possible that exotic species simply have not yet been introduced or had adequate time to spread from local points of introduction. Dispersal of earthworms is relatively slow ($10\text{--}15\text{ m y}^{-1}$; Hoogerkamp et al. 1983; Ghilarov and Perel 1984; also see Terhivuo and Saura, this issue), and years to decades may be required for proliferation of an exotic population after it has been introduced. Observations by Alban and Berry (1994) and Hale et al. (2005) suggest extended periods of time between introduction of European lumbricid species and their invasion of earthworm-free forests in Minnesota (also see Frelich et al. and Tiunov et al., this issue). Repeated introductions of an exotic species (i.e., high propagule pressure) may increase the likelihood of its establishment. However, other factors also influence invasion potential, including species characteristics such as fecundity and parthenogenesis; habitat characteristics such as dominant vegetation, soil and climatic conditions; and indigenous biota, such as predators, parasites and competitors (possibly including native earthworms). These factors may impart invasion resistance to a given habitat.

Habitat matching

Once introduced, exotic earthworms may fail to become established if they are not pre-adapted to a local habitat. A number of abiotic factors are

known to influence earthworm distribution and abundance, and hence the success of introduced species. Temperature and water regimes appear to be controlling factors for many invasive taxa on a global scale, for example limiting European lumbricids to temperate regions or *Pontoscolex corethrurus* to the tropics (Gates 1970; Fragoso et al. 1999). At local scales, soil properties such as texture, pH, Ca/Mg ratios, and soil organic matter content are important determinant of invasion success (Lee 1985; Edwards and Bohlen 1996).

Introduced earthworms also may not become established if resources in a new site are limiting to growth and reproduction. It has been suggested that habitat disturbance, such as fertilizer amendments or vegetation conversion, increase resource availability to anthropochorous earthworms thus enhancing their ability to invade disturbed sites (Fragoso et al. 1999; Winsome et al. 2006). Even in the absence of disturbance, it would be expected that an invader's feeding strategy would have to match the resource base in a new habitat for it to become established (e.g., epigeic species would be unsuccessful in areas devoid of surface litter).

In microcosm experiments, exotic earthworms have shown both reduced and increased survival, growth and reproduction in soils from invaded habitats, apparently depending on species and site characteristics. For example, *A. trapezoides*, *A. caliginosa* and *Octolasion cyaneum* did better in South Australian scrub vegetation soils containing indigenous earthworm casting than in nearby pasture soils in which they had become established (Lawson 1993). Conversely, *A. trapezoides* lost weight and *A. longa* failed to reproduce in Eucalyptus forest soils in Western and South Australia, respectively (Abbott 1985; Dalby et al. 1998). Abbott (1985) concluded that *A. trapezoides* was not well adapted to low organic matter content of these forest soils. In California grasslands, Winsome et al. (2006) found that invasive *A. trapezoides* was better adapted (in terms of growth) to heavily amended pasture soils than to relatively less enriched native grassland soils.

Exotic species' adaptations to temperature and precipitation regimes are reasonably predictive of invasion success on a broad scale (Lee 1985), but

predictions at a finer scale appear difficult because of the high degree of local variability within many ecosystems. Furthermore, while establishment of an exotic earthworm species in a new habitat may suggest that it has overcome the hurdles of propagule pressure and habitat matching, it does not necessarily guarantee successful invasion of intact native earthworm communities.

Biotic resistance

Once introduced into a habitat to which they are otherwise adapted, exotic species may fail to become established for biological reasons, such as predation (e.g., by birds, lizards or moles), parasitism (e.g., ecto- or intra-coelomic nematodes), or effective competition by resident native species, including indigenous earthworms. Indirect evidence of biotic resistance comes from studies in undisturbed ecosystems where well-adapted exotic earthworms are known to have been introduced or have become established nearby, but have failed to invade a particular habitat occupied by native earthworms (e.g., several of the minimally disturbed sites in Table 1). However, of those cases where natives occur exclusively, many appear to be explainable on the basis of habitat factors (e.g., low pH and coarse textured soils in Florida or Ivory Coast; serpentine soils in California oak savanna), which may be unfavorable to the exotic earthworms. Exceptions are forests studied by Abbott (1985), Lavelle and Pashanasi (1989) and Kalisz (1993), where there is no apparent reason why exotic species have not dispersed from old logging, homestead or cultivated sites into native earthworm communities within the forest.

Microcosm studies give some support to the biotic resistance hypothesis, and specifically to direct competitive interactions between native and exotic earthworms. Winsome et al. (2006) found that native *Argilophilus marmoratus* negatively affected *Aporrectodea trapezoides* growth and development in native California grassland soils, but not in enriched pasture soils nearby; *A. trapezoides* was the stronger competitor when resources were not limiting, but *A. marmoratus* was better adapted to the low-productivity grasslands and exacerbated the effects of resource limitation on *A. trapezoides* in the native habitat.

A further aspect of biotic resistance may relate to interactions between introduced earthworms and soil microflora. Daane and Häggblom (1999) found that earthworm cocoons in sterile medium did not develop as successfully as those in non-sterile medium, suggesting a functional linkage between earthworms and the ambient soil microflora. Furthermore, Gilot-Villenave (1994) has proposed that earthworms introduced into a new habitat may be impaired if they encounter an unfamiliar microflora, whereas cocoons of the same species may survive if they carry an indigenous microbial inoculum. If true, this phenomenon raises interesting questions for earthworm invasion ecology: Do sites inhabited by native earthworms maintain microbial populations unfavorable to exotic earthworms? Do wormless sites have a different microflora that offers less resistance (implying that earthworms can build resistance by modifying microfloras)? Do disturbed areas have depauperate (or even exotic) microfloras that do not offer this resistance to invasion by pre-hatched earthworms? Are there practical implications for intentional introduction of earthworms (e.g., for land reclamation efforts)? Some studies do suggest an internal or external “rumen” in earthworm feeding whereby soil or gut microbes facilitate catabolism and assimilation of organic substrates by earthworms (Lavelle et al. 1995; Brown and Doube 2004). An analysis of the microbial flora of earthworm gut material demonstrated that 12 phospholipid fatty acid markers occurred only in gut compartments and not in the bulk soil (Sampedro et al. 2003). Bacteria phylotypes isolated from intestinal tissue of *Lumbricus rubellus* were not detected in cast material or bulk soil, but it was suggested that the association was opportunistic rather than obligate (Singleton et al. 2003). There are few data with which to test the idea of obligate or antagonistic microbial associations with earthworms, or their implications for invasion ecology, but these are important questions for further research.

Differences between native and exotic earthworm assemblages

Based on general knowledge of earthworm ecology, some potential differences between native

and exotic earthworm populations might affect the likelihood and outcome of exotic invasions into native earthworm communities. First, native earthworm densities and fecundities may be lower than those of invasive species, even in undisturbed soils (Lee 1985; Fragoso 1999; Winsome 2003). This situation could give r-selected, rapidly growing exotic species populations a competitive advantage over native fauna for common resources. Second, at least some native earthworm assemblages appear to be dominated by endogeic species (Kalisz 1993; Fragoso et al. 1999), possibly providing open niches in the O-horizon which could be readily exploited by epigeic exotic species, for example *Amyntas agrestis* in deciduous forests occupied by native earthworms in north Georgia, USA (Callaham et al. 2003). Third, native earthworms may be better adapted to local conditions and thus have a competitive advantage over exotic species during periods when climatic conditions force exotics into dormancy, as observed in prairie soils in Kansas, USA (James 1991; Callaham et al. 2001). These situations are somewhat speculative, but may be involved in some cases of exotic invasions into native earthworm communities.

Co-existence of native and exotic earthworms

Table 1 summarizes information from studies that have assessed the status of native and exotic earthworm species in ecosystems under various degrees of disturbance. Native earthworms appear to occur exclusively or to predominate over exotic earthworms mostly in relatively undisturbed sites. Nonetheless, co-occurrence of native and exotic species, especially in disturbed or managed sites, appears to be common across a range of ecosystem types. Intensity of and time since disturbance appear to be important correlates of relative abundances of native and exotic species (Fragoso et al. 1999). Biotic resistance, if it exists in earthworm communities, may be more a matter of degree than an absolute outcome of native and exotic species interactions. From a practical standpoint, perhaps the more important questions are: under what circumstances do

native and exotic species co-exist in a given volume of soil, and are these situations persistent in the long term? Again, there are only limited data with which to address these questions.

One mechanism for co-existence of exotic and native species may be spatial partitioning of resources. In tropical forests of Chajul, Mexico, Fragoso (personal observation) found that exotic *P. corethrurus* was well established in a low-species-diversity earthworm community (6 species) in a poor forest soil (ferralitic) near a small village, and accounted for more than 84% of total earthworm abundance and biomass. Old alluvial soils, in which a tropical forest was well established, harbored a richer community (11 species) but with *P. corethrurus* still the most important species (41 and 35% of total abundance and biomass). Nonetheless, there was evidence of changes in the community in response to the presence of the invader. For example, *Balanteodrilus pearsei* (a very common species in southeast tropical Mexico) was relatively thin and small, compared to individuals in other populations located 70 km north, where *P. corethrurus* was absent. The vertical distribution of *P. corethrurus* was more superficial in the alluvial soils than in the ferralitic ones, suggesting that other mesohumic endogeic species inhabiting alluvial soils (e.g., *Ramiellona strigosa* and *Lavellodrilus illkus*) impeded *P. corethrurus* utilization of deeper strata. Thus, we can hypothesize that native endogeic species prevented the invasion of deeper soil by the exotic *P. corethrurus* which instead concentrated in the upper layers where it negatively affected the native polyhumic *B. pearsei*. Lachnicht et al. (2002) also observed spatial partitioning of the soil volume in microcosms derived from tabonuco forests in Puerto Rico. *Pontoscolex corethrurus* was active in the upper mineral soil and forest floor layers, whereas the native *Esthrella* sp. (possibly an anecic species) occupied the deeper mineral soil after a 30-day incubation.

Co-existence may also be facilitated by temporal separation of activity between native and exotic earthworms. For example, James (1991) and Callaham et al. (2001) suggested that native *Diplocardia* in tallgrass prairie soils were adapted to higher temperatures than were the invading European lumbricids, and thus maintained

activity during warmer periods when lumbricids became dormant. Regardless of the mode of action, these studies suggest the potential for co-existence of native and exotic earthworms and for resource partitioning in the same soil volume. However, it is unknown whether such co-existence is a transient or long-term phenomenon.

Finally, an intriguing aspect of co-existence is the possibility that native species actually facilitate the establishment of exotic species. Lawson's (1993) microcosm studies (reviewed above) suggested that several invasive European lumbricid species were better adapted to or perhaps able to more effectively exploit resources in soils containing castings from native earthworms than in the pasture soils they had successfully invaded. If this phenomenon is generally observed, it may change our view of exotic earthworm invasions in areas inhabited by indigenous earthworms.

Implications of native–exotic earthworm interactions for soil processes

A key functional question regarding exotic invasions into native earthworm communities is whether or not the impacts of exotic species on soil processes are altered in the presence of native species in their native habitats. Once again, there are very few data with which to address this question. The extreme-case affirmative answer would of course be where exotic species fail to establish after being introduced into a habitat. However, the more interesting situations would be where native and exotic species co-exist. If impacts are observed on soil processes, they might be expected to be related to relative population densities of natives and exotics at times of peak activity (Winsome et al. 2006).

As noted above, James (1991) suggested that native earthworms were better adapted to local soil and climatic conditions, and hence maintained longer periods of activity and effects on nutrient dynamics in tallgrass prairie soils than did invading European lumbricids. This is not necessarily always the case, however, for example where exotic species invade an area to which they are climatically well matched (e.g., Asian subtropical *A. agrestis* in Georgia, USA; Callahan

et al. 2003). Finally, the microcosms experiment by Lachniet et al. (2002) showed significant reductions in C and N mineralization rates induced by *P. corethrurus* when it was incubated in soils with native *Estherella* sp. compared to when it was incubated alone. Whether or not these effects would occur under field conditions is not known, but this is clearly a topic in need of further research.

Conclusions

Although the database is limited, a few conclusions can be drawn from this review. First, exotic earthworms do invade ecosystems inhabited by indigenous earthworms, even in the absence of obvious disturbance (Table 1). Kalisz and Wood (1995) referred to this phenomenon as “invisible” disturbance, such as forest fragmentation in which native species may experience local extinction in small remnants. In some cases, native species appear to remain dominant, usually in undisturbed soils; in other cases (both disturbed and minimally disturbed soils), the exotics predominate (Fragoso et al. 1995, 1999). A number of factors are probably involved in these outcomes, including physical and ecological characteristics of the habitat, biological characteristics of native and exotic earthworm species, influences of other indigenous biota, and time and frequency of invasions. In many cases, there must certainly be an element of chance that introduction of exotic species even occurred. Thus, the challenge for developing predictive models of exotic earthworm invasions is significant.

Second, direct competitive exclusion of native earthworms by exotic earthworms seems plausible in theory, but is not easily demonstrated in practice. In fact, field studies suggest that co-existence of native and exotic species is common (Table 1), even if transient. Competitive interactions may occur, as suggested by several microcosm studies, but it also appears that at least some exotic earthworms may utilize resources not fully exploited by native species, especially in disturbed soils. Because many of the invasive species show considerable flexibility in their use of resources and/or microhabitats (Fragoso et al.

1999), they may be well adapted to establish populations within areas occupied by native earthworm communities. The example from Chajul, Mexico cited above illustrates a certain amount of flexibility by an exotic species. An extreme case of this flexibility is the observation by S. James (unpublished data) of *P. corethrurus* occupying arboreal habitats in a montane cloud forest on Nevis Island in the Lesser Antilles; the site had no native earthworms and *P. corethrurus* was found in the soil and in the trees. The possible combination of exotic species flexibility and “open” niche space left by native species raises interesting evolutionary questions.

Third, resistance to exotic earthworm invasions, if it occurs, may be more a function of physical and chemical characteristics of a habitat than of biological interactions with native earthworms.

Acknowledgements This project was supported by the National Research Initiative of the USDA Co-operative State Research, Education and Extension Service, grant number 2003-35107-13876, and by National Science Foundation grant number 0236276 to the University of Georgia Research Foundation, Inc.

References

- Abbott I (1985) Distribution of introduced earthworms in northern jarrah forest of Western Australia. *Aust J Soil Res* 23:263–270
- Alban DH, Berry EC (1994) Effects of earthworm invasion on morphology, carbon and nitrogen of a forest soil. *Appl Soil Ecol* 1:243–249
- Baker G, Carter P, Barrett V, Hirth J, Mele P, Gourley C (2002) Does the deep-burrowing earthworm, *Aporrectodea longa*, compete with resident earthworm communities when introduced to pastures in south-eastern Australia? *Eur J Soil Biol* 38:39–42
- Beddard FE (1912) Earthworms and their allies. Cambridge University Press, London, p 145
- Bhadoria T, Ramakrishnan PS, Srivastava KN (2000) Diversity and distribution of endemic and exotic earthworms in natural and regenerating ecosystems in the central Himalayas, India. *Soil Biol Biochem* 32:2045–2054
- Bohlen PJ, Groffman PM, Fahey TJ, Fisk MC, Suarez E, Pelletier DM, Fahey RT (2004a) Ecosystem consequences of exotic earthworm invasion of north temperate forests. *Ecosystems* 7:1–12
- Bohlen PJ, Scheu S, Hale CM, McLean MA, Migge S, Groffman PM, Parkinson D (2004b) Non-native invasive earthworms as agents of change in northern temperate forests. *Front Ecol Environ* 2:427–435
- Brown GG, Doube BM (2004) Functional interactions between earthworms, microorganisms, organic matter and plants. In: Edwards C (ed) *Earthworm ecology*. 2nd edn. CRC Press, Boca Raton, FL, pp 213–239
- Burtelow AE, Bohlen PJ, Groffman PM (1998) Influence of exotic earthworm invasion on soil organic matter, microbial biomass and denitrification potential in forest soils of the northeastern United States. *Appl Soil Ecol* 9:197–202
- Callaham MA Jr, Hendrix PF (1997) Relative abundance and seasonal activity of earthworms (Lumbricidae and Megascolecidae) as determined by hand-sorting and formalin extraction in forest soils on the southern Appalachian piedmont. *Soil Biol Biochem* 29:317–321
- Callaham MA Jr, Blair JM (1999) Influence of differing land management on the invasion of North American tallgrass prairie soils by European earthworms. *Pedobiologia* 43:507–512
- Callaham MA Jr, Blair JM, Hendrix PF (2001) Native North American and introduced European earthworms in tallgrass prairie: behavioral patterns and influences on plant growth. *Biol Fertil Soil* 34:49–56
- Callaham MA Jr, Hendrix PF, Phillips RJ (2003) Occurrence of an exotic earthworm (*Amyntas agrestis*) in undisturbed soil of the southern Appalachian Mountains, USA. *Pedobiologia* 47:466–470
- Daane LL, Häggblom MM (1999) Earthworm egg capsules as vectors for the environmental introduction of biodegradative bacteria. *Appl Environ Microbiol* 65:2376–2381
- Dalby PR, Baker GH, Smith SE (1998) Potential impact of an introduced lumbricid on a native woodland in South Australia. *Appl Soil Ecol* 9:351–354
- Damoff GA (2005) Are east Texas bottomland forest soils being re-engineered by the exotic invasive earthworm *Amyntas diffringens* (Baird, 1869)? American Society of Agronomy International Annual Meetings, November 6–10, 2005, Salt Lake City, UT. Abstract 192-9
- Decaëns T, Blanchart E, Fragoso C, Lavelle P, Jimenez JJ, Barros E, Chauvel A (2004) Soil macrofaunal communities in permanent pastures derived from tropical forest or savanna. *Agric Ecosyst Environ* 103:301–312
- Dotson DB, Kalisz PJ (1989) Characteristics and ecological relationships of earthworm assemblages in undisturbed forest soils in the southern Appalachians of Kentucky, USA. *Pedobiologia* 33:211–220
- Edwards CA (ed) (2004) *Earthworm ecology*, 2nd edn. St. Lucie Press, Boca Raton, FL
- Edwards CA, Bohlen PJ (1996) *The biology and ecology of earthworms*, 3rd edn. Chapman and Hall, London
- Eisen G (1900) *Researches in American Oligochaeta*, with especial reference to those of the Pacific Coast and adjacent islands. *Proc Calif Acad Sci* II:85–276
- Fragoso C, Barois I, James S (1995) Native earthworms of the north Neotropical region: current status and controversies. In: Hendrix PF (ed) *Earthworm ecology and biogeography in North America*. Lewis Publishers, Boca Raton, FL

- Fragoso C, Lavelle P, Blanchart E, Senapati B, Jimenez J, de los Angeles Martinez M, Decaens T, Tondoh J (1999) Earthworm communities of tropical agroecosystems: origin, structure and influences of management practices. In: Lavelle P, Brussaard L, Hendrix P (eds) Earthworm management in tropical agroecosystems. CABI Publishing, New York, pp 27–55
- Gates GE (1970) Miscellaneous Megadrilologica VII. Megadrilologica 1:1–6
- Ghilarov MS, Perel TS (1984) Transfer of earthworms (Lumbricidae, Oligochaeta) for soil amelioration in the USSR. *Pedobiologia* 27:107–113
- Gilot-Villenave C (1994) Determination of the origin of the different growing abilities of two populations of *Millsonia anomala* (Omodeo and Vaillaud), a tropical geophagous earthworm. *Eur J Soil Biol* 39:125–131
- González G, Zou X (1999) Earthworm influence on N availability and the growth of *Cecropia schreberiana* in tropical pasture and forest soils. *Pedobiologia* 43(6):824–829
- González G, Seastedt TR, Donato Z (2003) Earthworms, arthropods and plant litter decomposition in aspen (*Populus tremuloides*) and lodgepole pine (*Pinus contorta*) forests in Colorado, USA. *Pedobiologia* 47:863–869
- González G, Zou X, Borges S (1996) Earthworm abundance and species composition in abandoned tropical croplands: comparisons of tree plantations and secondary forests. *Pedobiologia* 40:385–391
- Graham RC, Wood HB (1991) Morphologic development and clay redistribution in lysimeter soils under chaparral and pine. *Soil Sci Soc Am J* 55:1638–1646
- Graham RC, Ervin JO, Wood HB (1995) Aggregate stability under oak and pine after four decades of soil development. *Soil Sci Soc Am J* 59:1740–1744
- Hale CM, Frelich LE, Reich PB (2005) Exotic European earthworm invasion dynamics in northern hardwood forests of Minnesota, USA. *Ecol Appl* 15:848–860
- Hendrix PF (ed) (1995) Earthworm ecology and biogeography in North America. Lewis Publishers, Boca Raton, FL
- Hendrix PF, Bohlen P (2002) Ecological assessment of exotic earthworm invasions in North America. *Bioscience* 52:801–811
- Hendrix PF, Callahan MA Jr, Lachnicht SL, Blair JM, James SW, Zou X (1999a) Stable isotopic studies of resource utilization by nearctic earthworms (*Diplocardia*, *Oligochaeta*) in subtropical savanna and forest ecosystems. *Pedobiologia* 43:818–823
- Hendrix PF, Lachnicht SL, Callahan MA Jr, Zou X (1999b) Stable isotopic studies of earthworm feeding ecology in tropical ecosystems of Puerto Rico. *Rapid Commun Mass Spectrom* 13:1295–1299
- Hendrix PF, Mueller BR, Bruce RR, Langdale GW, Parmelee RW (1992) Abundance and distribution of earthworms in relation to landscape factors on the Georgia Piedmont, USA. *Soil Biol Biochem* 24:1357–1361
- Hendrix PF, Peterson AC, Beare MH, Coleman DC (1998) Long-term effects of earthworms on microbial biomass nitrogen in coarse and fine textured soils. *Appl Soil Ecol* 9:375–380
- Hoogerkamp M, Rogaar H, Eijsackers HJP (1983) Effect of earthworms on grassland on recently reclaimed polder soils in the Netherlands. In: Satchell JE (ed) Earthworm ecology from Darwin to vermiculture. Chapman and Hall, London, pp 85–105
- James SW (1982) Effects of fire and soil type on earthworm populations in a tallgrass prairie. *Pedobiologia* 24:37–40
- James SW (1991) Soil, nitrogen, phosphorus, and organic matter processing by earthworms in tallgrass prairie. *Ecology* 72:2101–2109
- James SW, Hendrix PF (2004) Invasion of exotic earthworms into North America and other regions. In Edwards CA (ed) Earthworm ecology, 2nd edn. CRC Press, Boca Raton, FL, pp 75–88
- Jiménez JJ, Moreno AG, Decaens T, Lavelle P, Fisher MJ, Thomas RJ (1998) Earthworm communities in native savannas and man-made pastures of the Eastern Plains of Colombia. *Biol Fertil Soils* 28:101–110
- Jiménez JJ, Rossi JP, Lavelle P (2001) Spatial distribution of earthworms in acid-soil savannas of the eastern plains of Colombia. *Appl Soil Ecol* 17:267–278
- Kalisz P (1993) Native and exotic earthworms in deciduous forest soils of eastern North America. In: Knight BN (ed) Biological pollution: the control and impact of invasive exotic species. Indiana Academy of Science, Indianapolis, IN, pp 93–100
- Kalisz PJ, Dotson DB (1989) Land-use history and the occurrence of exotic earthworms in the mountains of eastern Kentucky. *Am Midl Nat* 122:288–297
- Kalisz PJ, Wood HB (1995) Native and exotic earthworms in wildland ecosystems. In: Hendrix P (ed) Earthworm ecology and biogeography in North America. Lewis Publishers, Boca Raton, FL, pp 117–126
- Lachnicht SL, Hendrix PF, Zou X (2002) Interactive effects of native and exotic earthworms on resource use and nutrient mineralization in a tropical wet forest soil of Puerto Rico. *Biol Fertil Soil* 36:43–52
- Langmaid KK (1964) Some effects of earthworm invasion in virgin podsols. *Can J Soil Sci* 44:34–37
- Lavelle P (1978) Les vers de terre de la savanne de Lamto (Côte d'Ivoire): peuplements, populations et fonctions de l'écosystème, vol 12. Publications Laboratoire Zoologie, ENS, Paris
- Lavelle P, Pashanasi B (1989) Soil macrofauna and land management in Peruvian Amazonia (Yurimaguas, Loreto). *Pedobiologia* 33:283–291
- Lavelle PC, Lattaud D, Barois TI (1995) Mutualism and biodiversity in soils. *Plant Soil* 170:23–33
- Lavelle P, Brussaard L, Hendrix P (eds) (1999) Earthworm management in tropical agroecosystems. CABI Publishing, London
- Lawson LM (nee Wheeler) (1993) The distribution and abundance of native and introduced earthworms in an area of pasture and native vegetation near Cape Jervis, South Australia. Ph.D. Dissertation, School of Biological Sciences, Flinders University of South Australia
- Lee KE (1961) Interactions between native and introduced earthworms. *Proc NZ Ecol Soc* 8:60–62

- Lee KE (1985) Earthworms, their ecology and relationships with soils and land use. Academic Press, Orlando, FL
- Liu ZG, Zou XM (2002) Exotic earthworms accelerate plant litter decomposition in a Puerto Rican pasture and a wet forest. *Ecol Appl* 12:1406–1417
- Ljungstrom PO (1972) Taxonomical and ecological notes on the earthworm genus *Udeina* and a requiem for the South African acanthodrilines. *Pedobiologia* 12:100–110
- Mack RN, Simberloff D, Lonsdale WM, Evans H, Clout M, Bazzaz FA (2000) Biotic invasions: causes, epidemiology, global consequences and control. *Ecol Appl* 10:689–710
- McLean MA, Parkinson D (2000) Field evidence of the effects of the epigeic earthworm *Dendrobaena octaedra* on the microfungus community in pine forest floor. *Soil Biol Biochem* 32:351–360
- Mele PM, Carter MR (1999a) Species abundance of earthworms in arable and pasture soils in south-eastern Australia. *Appl Soil Ecol* 12:129–137
- Mele PM, Carter MR (1999b) Impact of crop management factors in conservation tillage farming on earthworm density, age structure and species abundance in south-eastern Australia. *Soil Tillage Res* 50:1–10
- Parmelee RW, Beare MH, Cheng WX, Hendrix PF, Rider SJ, Crossley DA Jr, Coleman DC (1990) Earthworms and enchytraeids in conventional and no-tillage agroecosystems: a biocide approach to assess their role in organic matter breakdown. *Biol Fertil Soil* 10:1–10
- Peterson AC, Hendrix PF, Haydu C, Graham RC, Quideau SA (2001) Single-tree influence on earthworms and soil macroarthropods in the southern California chaparral. *Pedobiologia* 45:509–522
- Reynolds JW (1995) Status of exotic earthworm systematics and biogeography in North America. In: Hendrix P (ed) Earthworm ecology and biogeography in North America. Lewis Publishers, Boca Raton, FL, pp 1–28
- Sampedro L, Whalen JK, Shanmugam B, Waheed T (2003) Changes in soil microbial community structure during transit through the earthworm gut. *Abstr Soil Ecol Soc* 9:69
- Satchell JE (ed) (1983) Earthworm ecology, from Darwin to vermiculture. Chapman and Hall, London
- Scheu S, Parkinson D (1994) Effects of invasion of an aspen forest (Canada) by *Dendrobaena octaedra* (Lumbricidae) on plant growth. *Ecology* 75:2348–2361
- Simberloff D (1989) Introduced insects: a biogeographical and systematic perspective. In Mooney HA, Drake JA (eds) Ecology of biological invasions of North America and Hawaii ecological studies 58. Springer-Verlag, NY, pp 3–26
- Singleton DR, Hendrix PF, Coleman DC, Whitman WB (2003) Identification of uncultured bacteria tightly associated with the intestine of the earthworm *Lumbricus rubellus* (Lumbricidae; Oligochaeta). *Soil Biol Biochem* 35:1547–1555
- Smith F (1928) An account of changes in the earthworm fauna of Illinois and a description of one new species. *Ill Nat Hist Surv Bull* 17:347–362
- Stebbins JH (1962) Endemic-exotic earthworm competition in the American Midwest. *Nature* 196:905–906
- Steinberg DA, Pouyat RV, Parmelee RW, Groffman PM (1997) Earthworm abundance and nitrogen mineralization rates along an urban-rural gradient. *Soil Biol Biochem* 29:427–430
- Stockdill SMJ (1982) Effect of introduced earthworms on the productivity of New Zealand pastures. *Pedobiologia* 24:29–35
- Williamson M (1996) Biological invasions. Chapman and Hall, London
- Winsome T (2003) Native and exotic earthworms in a California oak savanna ecosystem. Ph.D. Dissertation, University of Georgia, USA, p 124
- Winsome T, Epstein L, Hendrix PF, Horwath WR (2006) Habitat quality and interspecific competition between native and exotic earthworm species in a California grassland. *Appl Soil Ecol* 32:38–53
- Wood HB, Oliver KL, James SW (1997) Relict Megascleidae and exclusion of Lumbricidae from basalt-derived soils in southern California. *Soil Biol Biochem* 29:241–244
- Zou XM, Bashkin M (1998) Soil carbon accretion and earthworm recovery following revegetation in abandoned sugarcane fields. *Soil Biol Biochem* 30:825–830
- Zou XM, Gonzalez G (1997) Changes in earthworm density and community structure during secondary succession in abandoned tropical pastures. *Soil Biol Biochem* 29:627–629