

# Relationship between stem CO<sub>2</sub> efflux, stem sap velocity and xylem CO<sub>2</sub> concentration in young loblolly pine trees

CHRIS A. MAIER<sup>1</sup> & BARTON D. CLINTON<sup>2</sup>

<sup>1</sup>U.S.D.A. Forest Service, Research Triangle Park, NC 27709, and <sup>2</sup>U.S.D.A. Forest Service, Coweeta Hydrologic Laboratory, Otto, NC 28763, USA

## ABSTRACT

We measured diel patterns of stem surface CO<sub>2</sub> efflux ( $E_s$ ,  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ), sap velocity ( $v_s$ ,  $\text{mm s}^{-1}$ ) and xylem CO<sub>2</sub> concentration ([CO<sub>2</sub>]) ( $X_s$ , %) in 8-year-old loblolly pine trees during the spring to determine how  $v_s$  and  $X_s$  influence  $E_s$ . All trees showed a strong diel hysteresis between  $E_s$  and stem temperature, where at a given temperature,  $E_s$  was lower during the day than at night. Diel variations in temperature-independent  $E_s$  were correlated with  $v_s$  ( $R^2 = 0.54$ ), such that at maximum  $v_s$ ,  $E_s$  was reduced between 18 and 40%. However, this correlation may not represent a cause-and-effect relationship. In a subset of trees,  $v_s$  was artificially reduced by progressively removing the tree canopy. Reducing  $v_s$  to near zero had no effect on  $E_s$  and did not change the diel hysteretic response to temperature. Diel  $X_s$  tended to decrease with  $v_s$  and increase with  $E_s$ , however, in defoliated trees, large increases in  $X_s$ , when  $v_s \approx 0$ , had no effect on  $E_s$ . We conclude that at this time of the year,  $E_s$  is driven primarily by respiration of cambium and phloem tissues and that sap flow and xylem transport of CO<sub>2</sub> had no direct influence on  $E_s$ .

**Key-words:** chambers; CO<sub>2</sub> microelectrode; Granier sensors; *Pinus taeda*; sap flow; stem respiration.

## INTRODUCTION

Respiration of above ground woody tissues (stem and branch) comprises 15–25% of forest ecosystem respiration (Ryan *et al.* 1994, 1996; Xu *et al.* 2001; Maier *et al.* 2004). These estimates are based on empirical data where CO<sub>2</sub> efflux from the stem (or branch) surface into a chamber is measured with an infrared gas analyzer. This approach assumes that CO<sub>2</sub> generated from metabolism of cambium and xylem parenchyma tissues enclosed within the chamber diffuses radially from the stem interior across the cambial sheath to the surface. However, at high transpiration rates, a portion of the respired CO<sub>2</sub> in sapwood may be carried upward by the transpiration stream instead of released horizontally through the bark, so that measured CO<sub>2</sub> efflux underestimate the actual respiration of the sample section.

*Correspondence:* Chris A. Maier. Fax: +1 919 549 4047; e-mail: cmaier@fs.fed.us

A number of studies (Negisi 1975, 1978, 1982; Lavigne 1987; Kabubari 1988) found that on warm sunny days, measured stem CO<sub>2</sub> efflux ( $E_s$ ) rates were much lower, 25–50%, compared with what would be expected based on temperature alone. Other studies found that the diel relationship between  $E_s$  and temperature exhibits a hysteresis, where CO<sub>2</sub> efflux measurements made at a similar temperature is higher in the late afternoon and evening, when transpiration is low, than in the morning, when transpiration is high (Martin, Teskey & Dougherty 1994; Ryan *et al.* 1995; Lavigne 1996; Stockfors 2000; Maier 2001; Bosc, De Grandcourt & Loustau 2003). These studies suggest that stem surface CO<sub>2</sub> efflux in forest ecosystems may be linked to canopy water use.

Stem CO<sub>2</sub> concentrations are high, ranging from 2 to 10% (Hari, Pekka & Korpilahti 1991; Eklund 1993; Teskey & McGuire 2002). Dissolved carbon in the xylem (CO<sub>2</sub>, H<sub>2</sub>CO<sub>3</sub> and HCO<sub>3</sub><sup>-</sup>) is a combination of CO<sub>2</sub> derived from respiration of nearby xylem parenchyma and cambium tissues, CO<sub>2</sub> imported from respiratory activity of stem and roots lower in the xylem stream, and CO<sub>2</sub> taken up in soil water. Given the high xylem CO<sub>2</sub> concentration ([CO<sub>2</sub>]) ( $X_s$ ), the aqueous transport of carbon in the xylem stream represents a potentially large and poorly understood carbon flux in forest ecosystems. If transport and storage of CO<sub>2</sub> in the xylem strongly affects  $E_s$  under normal field conditions, then the interpretation of stem gas exchange measured with chamber methods becomes equivocal (Martin *et al.* 1994; Teskey & McGuire 2002). A more complete evaluation of these relationships is needed to understand variation in  $E_s$  rates.

Recently, there has been a renewed interest in measuring the origin and fate of carbon in the xylem sap and determining what effect this carbon flux may have on the measurement of  $E_s$  (Stringer & Kimmerer 1993; Martin *et al.* 1994; Kaipainen *et al.* 1998; Edwards & Wullschleger 2000; Clinton, Maier & Sullivan 2001; Teskey & McGuire 2002, 2005; McGuire & Teskey 2002, 2004; Bowman *et al.* 2005). Teskey & McGuire (2002) measured sap flow rate and  $X_s$  in large trees of several species (*Quercus alba*, *Liriodendron tulipifera* and *Pinus taeda*) and found that diel patterns were opposed, suggesting that transpiration may significantly affect stem [CO<sub>2</sub>] and thus the driving force for radial diffusion of CO<sub>2</sub> in stem tissue. Teskey & McGuire (2005) further demonstrated in hardwood

saplings, by artificially manipulating  $X_s$ , that stem surface CO<sub>2</sub> efflux was directly related to internal [CO<sub>2</sub>]. McGuire & Teskey (2004) proposed a mass balance approach for estimating stem respiration that accounted for the rates of xylem CO<sub>2</sub> inputs, outputs and storage and surface CO<sub>2</sub> efflux. They found that the diel flux of respired CO<sub>2</sub> within the stem followed different pathways dependent on sap flow rate. At night, when sap flow rates were low, stem surface CO<sub>2</sub> efflux accounted for 74–93% of total stem respiration (i.e. the total from all sources); but during the day, when sap flow rates were high, surface CO<sub>2</sub> efflux accounted for only 23–72% of estimated total stem respiration. Bowman *et al.* (2005) found similar results in several *Dacrydium cupressinum* trees. However, a consistent and measurable relationship between  $E_s$  and stem sap flow is far from universal. Clinton *et al.* (2001) found a negative relationship between apparent stem respiration and sap velocity ( $v_s$ ) in large yellow poplar trees, while others found either no relationship (Carey, Delucia & Ball 1996; unpublished observations, Edwards & Wullschleger 2000) or a positive correlation (Levy *et al.* 1999). In the Levy *et al.* (1999) study, increases in apparent stem respiration with  $v_s$  were attributed to transport of CO<sub>2</sub> from the roots, which were assumed to be in equilibrium with high soil pCO<sub>2</sub>.

In this study, we examined the relationship between stem surface CO<sub>2</sub> efflux rate (i.e. apparent stem respiration),  $v_s$  and  $X_s$  in stems of 8-year-old loblolly pine trees over a 2 week period. Half of the experimental trees had received optimum nutrition from annual fertilization since planting, while the other half grew in the native nutrient-poor soil. Fertilization had significantly increased tree height, diameter and leaf area relative to non-fertilized controls. Fertilized trees are likely to have a different wood hydraulic architecture (Tyree & Ewers 1991), as well as differing patterns of water uptake (Ewers, Oren & Sperry 2000), canopy conductance, stand transpiration (Ewers *et al.* 2001) and rates of maintenance respiration (Maier *et al.* 1998). In some of the trees, we artificially altered  $v_s$  through a step-wise reduction in canopy leaf area (Pataki, Oren & Phillips 1998). The objectives were to: (1) determine if there is a diel relationship between stem  $v_s$  and stem surface CO<sub>2</sub> efflux; (2) determine if there is a diel relationship between stem  $v_s$  and  $X_s$ ; and (3) determine if there is a relationship between  $X_s$  and stem surface CO<sub>2</sub> efflux.

## MATERIALS AND METHODS

### Site description

The study was conducted in an 8-year-old loblolly pine plantation located at the SETRES II GxE-QTL study site in Scotland County, NC, U.S.A (McKeand *et al.* 2000). The soil is a Wakulla series characterized as a sandy, siliceous, thermic Psammentic Hapludult (sand to >43 m), which is very infertile, somewhat excessively drained, with a water holding capacity of 10–12 cm in a 2 m profile. The site receives an average annual precipitation of 1200 mm

distributed evenly throughout the year. Annual temperature averaged 17 °C, with a seasonal average of 26 °C in summer and 9 °C in winter. Greenhouse-grown seedlings were planted in November 1993 after the removal of the existing 10-year-old loblolly pine. Five full-sib families of Atlantic coastal plain and Texas origin were planted in 100 tree plots. Our measurements were confined to a non-fertilized and fertilized plot of one Atlantic coastal plain family (9–1046). The site average leaf area index for the non-fertilized and fertilized plots were 1.19 and 2.91 m<sup>2</sup> m<sup>-2</sup>, respectively, in October 2000 (Francisco Flores, North Carolina State University, personal communication).

### Measurements

We selected six trees in the non-fertilized and fertilized plots (12 trees total). Tree height, stem diameter and number of branches were measured (Table 1). Branch and canopy foliage biomass was estimated using site-specific regression equations (Tim Albaugh, North Carolina State University, personal communication). Average ( $\pm$  SE) stem diameter and estimated canopy biomass were 8.2 ± 0.5 cm and 1725 ± 704 kg in non-fertilized trees and 10.6 ± 0.2 cm and 2658 ± 129 kg in fertilized trees.  $E_s$ , stem temperature and  $v_s$  were monitored continuously for 11 d. On the last 3 d,  $X_s$  was measured in a subset of trees. Instantaneous photosynthesis and stomatal conductance (Licor 6400, Li-Cor, Inc., Lincoln, NE, USA) were measured at the beginning, middle and end of the experiments on upper canopy 1-year-old foliage on all trees during the morning hours (0900–1100 h).

$E_s$  (μmol CO<sub>2</sub> m<sup>-2</sup> s<sup>-1</sup>) measurements were made using an automated, multichamber sampling system (Butnor, Johnsen & Maier 2005) that consisted of stem chambers, an infrared gas analyzer (EGM-2, PP Systems, Amesbury, MA, USA) and a series of solenoids that sequentially measured stem chambers. The system had an open flow-through design where CO<sub>2</sub> efflux was estimated as the difference between the CO<sub>2</sub> concentration entering and exiting the chamber. Chambers were constructed of Teflon film that surrounded the tree stem 1 m above the ground. The Teflon film was fastened to the stem using collars of closed-cell foam and double-sided tape. Air was distributed to and sampled from the chamber using diffuser rings positioned at the top and bottom of the chamber. Chamber lengths were 25 cm and chamber volume ranged from 0.00179 to 0.00269 m<sup>3</sup>, depending on stem diameter. All chambers were leak tested prior to use. Airflow to the chambers was fixed at 0.00225 m<sup>3</sup> min<sup>-1</sup>. Each chamber was measured for 6 min to assure stable CO<sub>2</sub> measurements. The last minute of each cycle was retained for calculation of surface CO<sub>2</sub> efflux rates. A complete cycle through all of the chambers, including a null chamber, was completed in 42 min, which equals approximately 34 observations for each chamber per day. All chambers were continuously flushed with ambient air (0.00225 m<sup>3</sup> min<sup>-1</sup>) when chambers were not measured. Simultaneous measurements of chamber air and stem cambium temperature (3 mm) were made using copper/

**Table 1.** Tree characteristics in April 2001 for the non-fertilized (NF) and fertilized (F) trees located at SETRES II, Scotland County, NC, USA

Plot	Tree	Treatment	d.b.h. (cm)	Height (cm)	Number of branches	Foliage biomass (g)	
						Predicted	Measured
NF	1	UC	9.6	560	44	2017	—
	2	C	8.3	541	43	2006	1984
	3	UC	6.6	476	37	1239	—
	4	C	7.1	459	47	1531	1185
	5	C	8.3	516	39	1661	1299
	6	UC	9.1	546	45	1898	—
F	1	C	10.8	767	37	3063	2781
	2	C	11.2	709	42	2491	2912
	3	C	10.4	701	38	2387	2758
	4	UC	10.7	714	32	2913	—
	5	UC	10.9	691	36	2820	—
	6	UC	9.8	755	34	2278	—

The column labelled 'Treatment' refers to whether the canopies of trees were removed [cut (C)] or left intact [uncut (UC)]. d.b.h., diameter at breast height.


constantan thermocouples. Immediately after the experiments were completed, stem diameter at the top and bottom of the chambers were measured with digital calipers. Stem surface area inside the chamber was estimated from the average of the two diameter measurements and includes the bark.

Stem  $v_s$  ( $\text{mm s}^{-1}$ ) was measured using custom-made 30-mm-long thermal dissipation sap velocity probes (Granier 1985, 1987). Briefly, paired probes were inserted radially into the tree such that the probes were approximately 5 cm apart vertically. For each tree, two probes were installed on opposite sides (north and south) of the stem just below the stem chamber.  $v_s$  was measured every 10 s and these values were averaged every 15 min. In our trees, essentially all of the xylem was hydroactive; however, while we only measured the outer 3 cm, the probes measured the previous 2 years of growth and captured the majority of stem sap flow in these trees (Ewers & Oren 2000).

$X_s$  (%) was measured *in situ* on four trees using  $\text{CO}_2$  microelectrodes (Model MI-720; Microelectrodes, Inc., Bedford, NH, USA). We followed methods described by McGuire & Teskey (2002) and Teskey & McGuire (2002). Briefly, electrodes were calibrated with humidified compressed  $\text{CO}_2$  gas at 2, 5 and 10% concentrations. Because the electrodes are temperature sensitive, a temperature correction was applied (McGuire & Teskey 2002). To measure  $X_s$ , a small hole 10 mm in diameter and 7–10 mm deep was drilled through the bark into the xylem, 20–25 cm below the stem chamber. The tip of a 5-cm-long low-density polyethylene tube was inserted into the hole, and the outside edge sealed to the tree with putty adhesive. A microelectrode was then inserted into the polyethylene tube such that the electrode tip did not make contact with xylem. Adhesive putty was used to seal the body of the microelectrode to the polyethylene tube. Four probes were installed, one each in four trees (two non-fertilized and two fertilized).

## Experiments

We examined the relationship between  $E_s$ ,  $v_s$  and  $X_s$  using two different approaches. In the first experiment, we compared hourly average measurements of  $E_s$  and  $v_s$  for three trees in the non-fertilized (trees 1, 3 and 6) and fertilized (trees 4, 5 and 6) plots (Table 1). In young loblolly pine trees, the response of  $E_s$  to diel changes in stem temperature typically exhibits a hysteresis, where at a similar temperature  $E_s$  is higher at night than during the day (Maier 2001) (Fig. 1). We assumed *a priori* that this diel hysteresis was a function of  $v_s$ . Therefore, to remove potential effects of  $v_s$ , only night-time (2300–0500 h)  $E_s$  measurements were



**Figure 1.** An example of the diel pattern of hysteresis between measured stem surface  $\text{CO}_2$  efflux ( $E_s$ ) and cambium temperature ( $T_c$ ), where at a given temperature,  $E_s$  was lower during the day than at night. Response patterns from a non-fertilized tree are shown for day of year (DOY) 110 near the beginning of the experiment and DOY 117 after >90% of the canopy had been removed. Corresponding measurements for three times are shown.

used to model  $E_s$ .  $v_s$  during this time was always less than  $0.01 \text{ mm s}^{-1}$ . Night-time  $E_s$  was modelled as a function of temperature by fitting the data to the exponential equation:

$$E_s = \beta_0 e^{(kT_c)}, \quad (1)$$

where  $E_s$  is measured stem  $\text{CO}_2$  efflux ( $\mu\text{mol m}^{-2} \text{s}^{-1}$ ),  $\beta_0$  is  $\text{CO}_2$  efflux at  $0^\circ\text{C}$ ,  $k$  is the temperature coefficient and  $T_c$  is measured cambium temperature. Nonlinear regression (PROC NLIN, SAS Institute, Cary, NC, USA) was used to estimate  $\beta_0$  and  $k$  in Eqn 1. Model performance was examined graphically by comparing predicted and observed values of  $E_s$  and by calculating the percent root mean square error (%RMSE), a measure of model precision, and the percent absolute deviation (%AD), an estimate of model accuracy (Maier 2001). A  $t$ -statistic was used to test for differences between non-fertilized and fertilized plots. Equation 1 was then used to predict diel patterns of  $E_s$  ( $E_p$ ). Residual  $E_s$  ( $E_r$ ), the difference between  $E_s$  and  $E_p$ , represents variation assumed to a result of xylem transport of  $\text{CO}_2$ . We hypothesized that the diel pattern of  $E_r$  and the ratio  $E_r/E_p$  would be correlated with  $v_s$ .

In the second experiment, we examined how artificially reducing  $v_s$ , through a progressive removal of canopy leaf area, affected the diel patterns  $E_s$  and  $X_s$ . In trees with low leaf area, transpiration is proportional to leaf area (Cieniakala & Lindroth 1995; Sala, Smith & Devitt 1996), and abrupt reductions in leaf area can reduce canopy transpiration and  $v_s$  (Oren *et al.* 1999). In this experiment, the canopies of three of the six trees in the non-fertilized (trees 2, 4 and 5) and fertilized (trees 1, 2 and 3) plots (Table 1) were removed in thirds (Cut treatment). The canopy of the cut tree was divided vertically into three levels based on an equal number of branches. Canopy removal was done equally from each level based on branch foliage biomass (Table 1). After an initial period, to establish individual tree  $E_s$  and  $v_s$  behaviour ( $\approx 48 \text{ h}$ ), one-third of the canopy leaf area in the cut trees was removed by removing branches. Measurements continued for 3 d, and then another third of foliage biomass was removed followed by another 4 d of measurements after which the remaining foliage was removed except for a single 1-year-old branch at the top of the canopy. Branch removal was completed by 1000 h on the day of treatment.

Diel patterns of  $v_s$  and  $E_s$  were compared graphically between uncut and cut trees. To facilitate comparisons, maximum daily rates of  $v_s$  and  $E_s$  data were normalized to the maximum rates measured on DOY (day of year) 109 before the cutting treatments began. Changes in the normalized maximum rates of  $v_s$  and  $E_s$  resulting from the cutting treatment were compared using repeated measures analysis of variance (ANOVA) (PROC MIXED, SAS Institute, Cary, NC, USA), utilizing an autoregressive covariance structure. We hypothesized that during the daylight hours maximum  $v_s$  would decrease and maximum  $E_s$  would increase in cut trees relative to uncut trees. On the last 2 d of the experiment, we compared diel patterns of  $v_s$ ,  $E_s$  and  $X_s$ . We hypothesized that  $X_s$  would increase in cut trees relative to uncut trees because respiration  $\text{CO}_2$  in

the xylem would accumulate when  $v_s$  is low in cut trees, and that increases in  $X_s$  in cut trees would cause a concomitant increase in  $E_s$ .

## RESULTS

### Diel patterns


Stem cambium temperatures ranged from  $0\text{--}30^\circ\text{C}$  over the 11 d of measurements (Fig. 2a). Stem temperatures peaked in early afternoon on most days.  $E_s$  had a strong diel pattern that was well correlated with cambium temperature (Fig. 2b), however, maximum daily  $E_s$  always occurred after maximum stem temperature (35–210 min) (Fig. 2a & b) creating a diel hysteresis. For example on DOY 110,  $E_s$  was lower during the morning and early afternoon than at night when measured at a similar temperature (Fig. 1). The magnitude of the daily hysteretic response varied among trees and within a tree over time.

Stem  $v_s$  increased rapidly during the morning and reached maximum values between 1000 and 1200 h then rapidly declined in the afternoon (Fig. 2c and inset). An exception to this pattern was observed on DOY 115, which was rainy and cool, when maximum  $v_s$  was measured in late afternoon. During the night,  $v_s$  was less than  $0.01 \text{ mm s}^{-1}$ . Peak sap velocity generally occurred 5–8 h before maximum  $E_s$ .

### Experiment 1

To remove potential effects of  $v_s$  on  $E_s$ , temperature response curves were developed using only nighttime measurements when  $v_s$  was less than  $0.01 \text{ mm s}^{-1}$ . Night-time  $E_s$  was well correlated ( $R^2 = 0.90\text{--}0.96$ , Eqn 1) with stem cambium temperature (Table 2). Predictions based on the equation showed good agreement with observed values. The %RMSE averaged 12.3% of mean night-time  $E_s$ . All of the models exhibited a mean %AD of less than 14% and indicates that the models accurately predicted night-time  $E_s$  over the time period measured. There was no significant difference between non-fertilized and fertilized trees in basal  $\text{CO}_2$  efflux rate ( $\beta_0$ ; non-fertilized:  $1.22 \pm 0.08\text{SE}$ ; fertilized:  $1.37 \pm 0.04$ ;  $P = 0.12$ ) or the temperature coefficient ( $k$ ; non-fertilized:  $0.073 \pm 0.001$ ; fertilized:  $0.071 \pm 0.001$ ;  $P = 0.77$ ).

Using the parameters in Table 2 with Eqn 1, we compared predicted  $E_p$  with observed  $E_s$  (Fig. 2b and inset). During the day,  $E_p$  was always greater than measured  $E_s$ , with the largest differences occurring around midday during periods of rapid change in stem temperature and  $v_s$ .  $E_r$ , the difference between  $E_p$  and  $E_s$ , generally decreased with increasing  $v_s$  (Fig. 2b and inset). The negative ratio of  $E_r/E_p$ , a relative measure of reduced  $E_s$ , was negatively correlated with  $v_s$  (Fig. 3) and there was no difference between non-fertilized and fertilized trees in the slope of this relationship. These data suggest that during the day high  $v_s$  could potentially reduce  $E_s$  up to 40% of that predicted on temperature alone.



**Figure 2.** An example of the diel response patterns of (a) cambium temperature ( $T_c$ ), (b) measured ( $E_s$ ), predicted ( $E_p$ ) and residual ( $E_r$ ) stem surface  $\text{CO}_2$  efflux, and (c) sap velocity ( $v_s$ ) for a fertilized uncut tree. Residual respiration is the difference between  $E_s$  and  $E_p$ . Predicted stem surface  $\text{CO}_2$  efflux ( $E_p$ ) was estimated using temperature response curves developed from nighttime  $E_s$  measurements when  $v_s < 0.01 \text{ mm s}^{-1}$ . Inset: Diel patterns of the parameters for 1 d. DOY, day of year.

## Experiment 2

In this experiment, we examined how artificially changing  $v_s$  through a progressive removal of canopy leaf area affected  $E_s$ . Foliage was removed in thirds on DOY 110, 113 and 117 (arrows, Fig. 4). To aid in making comparisons between cut and uncut trees,  $v_s$  and  $E_s$  were normalized to the maximum rates measured on DOY 109 before the branch removal treatment began (Fig. 5). There were no significant differences in  $v_s$  or  $E_s$  between non-fertilized and fertilized trees (Table 3). Removal of one-third to two-thirds of the canopy leaf area had only small effects on  $v_s$  and  $E_s$ ; and non-fertilized and fertilized trees behaved differently (Figs 4 & 5, Table 3). In the non-fertilized trees, maximum daily  $v_s$  was reduced  $\approx 20\%$  following the first

cutting (DOY 110) when roughly one-third of the canopy was removed (Fig. 5). Removal of the second third of the canopy (DOY 113) had no further effect on the magnitude of this response. In fertilized trees, the first cutting treatment had no effect on  $v_s$ , but  $v_s$  was significantly less in the cut trees 3 d following the second cutting treatment (Fig. 5). After removal of most of the canopy (DOY 117),  $v_s$  was reduced 80–90% of that in uncut trees in both non-fertilized and fertilized trees. The lack of a large decrease in  $v_s$  following abrupt changes in leaf area was likely due to stomatal compensation. Stomatal conductance significantly increased in foliage of cut trees after the second cutting treatment (Table 4). Net photosynthesis tended to increase in cut trees following branch removal, but this difference was only significant for DOY 111.

**Table 2.** Parameter estimates and fit statistics for Equation 1

	Tree	a	k	$r^2$	n <sup>a</sup>	%RMSE <sup>b</sup>	%AD <sup>c</sup>
Non-fertilized	1	1.53	0.075	0.94	209	12.6	9.5
	2	1.35	0.074	0.93	209	13.9	11.0
	3	1.06	0.070	0.96	206	12.2	8.9
	4	1.14	0.072	0.92	208	14.4	11.9
	5	1.05	0.073	0.94	202	12.8	10.4
	6	1.18	0.070	0.96	206	9.9	7.6
Fertilized	1	1.39	0.070	0.88	208	15.2	12.5
	2	1.50	0.072	0.87	209	16.2	13.5
	3	1.25	0.073	0.95	208	11.3	8.7
	4	1.37	0.072	0.95	206	10.9	7.9
	5	1.25	0.073	0.96	207	8.4	6.8
	6	1.43	0.068	0.96	208	9.6	7.3

Equation 1 was fitted to stem surface CO<sub>2</sub> efflux ( $E_s$ ) measured at night, between 2300 and 0500 h, when sap velocity ( $v_s$ ) was less than 0.01 mm s<sup>-1</sup>.

<sup>a</sup>n is the number of observations.


<sup>b</sup>Percent root mean square error. %RMSE =  $\left[ \frac{1}{n} \sum_{i=1}^n \left( \frac{\hat{y}_i - y_i}{y_i} \right)^2 \right]^{1/2} \times 100$

<sup>c</sup>Percent absolute deviation. %AD =  $\frac{100}{n} \sum_{i=1}^n \left| \frac{\hat{y}_i - y_i}{y_i} \right|$

In contrast, cutting treatment had little effect on the diel patterns of  $E_s$  (Fig. 4). There was no significant fertilizer or fertilizer-by-cutting treatment interaction on normalized  $E_s$  (Table 3). However, in the fertilized trees  $E_s$  declined 7–15% two days following the first cutting treatment (Fig. 5). This difference was maintained throughout the experiment. There was no apparent response of  $E_s$  to large changes in  $v_s$ . The pattern of  $E_s$  in fertilized and non-fertilized trees was

similar throughout the experiment even after the final removal of branches when  $v_s$  in cut trees was ≈10% of that in uncut trees (Figs 4 & 5). Furthermore, large reductions in  $v_s$  following the final cutting treatment had little effect on the diel  $E_s$ -temperature hysteresis. For example, the magnitude of the hysteresis was similar on days having a similar range in temperature, but a large difference in maximum  $v_s$  (compare DOY 110 and 117, Fig. 1).


We measured  $X_s$  in four trees over the last 2.5 d of the experiment. During this time,  $X_s$  ranged from 1 to 8%.  $X_s$  changed diurnally reaching a maximum at night and a minimum near noon (Fig. 6). In uncut trees,  $X_s$  generally decreased during the day when  $v_s$  was high and increased at night when  $v_s$  was low suggesting that sap flow had a strong influence over  $X_s$ . A similar pattern was observed in cut trees; however,  $X_s$  increased more relative to uncut trees when  $v_s$  was reduced following the final cutting treatment. Large diel changes in  $X_s$  appeared to have little



**Figure 3.** The relationship between the negative ratio of residual and predicted stem surface CO<sub>2</sub> efflux ( $E_r/E_p$ ) and sap velocity ( $v_s$ ). Predicted stem surface CO<sub>2</sub> efflux ( $E_p$ ) was estimated using temperature response curves developed from nighttime  $E_s$  measurements when  $v_s < 0.01$  mm s<sup>-1</sup>. Residual respiration is the difference between  $E_s$  and  $E_p$ .  $E_r/E_p$  is expressed as a negative ratio to illustrate the potential reduction in  $E_s$  at high  $v_s$ . Each point is the average of three trees.  $y = -0.026 - 4.526x$   $R^2 = 0.54$ .

**Table 3.** Probability values for the effect of fertilization, cutting treatment and time and their interactions on the normalized maximum daily sap velocity ( $v_s$ ) and stem surface CO<sub>2</sub> efflux ( $E_s$ )

Effect	$v_s$	$E_s$
Fertilization (F)	0.3964	0.4608
Cut (C)	<0.0001	0.0914
F × C	0.2556	0.1258
Day (D)	<0.0001	<0.0001
F × D	0.2127	<0.0001
C × D	<0.0001	0.1538
F × C × D	0.1709	0.4096



**Figure 4.** Diel patterns of hourly sap velocity ( $v_s$ ) and stem surface  $\text{CO}_2$  efflux ( $E_s$ ) for non-fertilized and fertilized trees with intact canopies (uncut) and those where the canopy was progressively removed (cut) in thirds (arrows) over the course of the study. Each point is the average of three trees. DOY, day of year.


effect on  $E_s$ . For example in the fertilized trees,  $X_s$  in cut trees increased twofold after trees received the last canopy removal (DOY 117), but there was no apparent change in  $E_s$ . In the non-fertilized trees,  $E_s$  was similar between uncut and cut trees on day 118 despite a fourfold difference in  $X_s$ . These data suggest that  $E_s$  in these trees was not influenced by  $\text{CO}_2$  transported in the xylem stream.

## DISCUSSION

Our trees showed the typical diel counterclockwise hysteresis between  $E_s$  and stem temperature (Fig. 1) reported for trees in other studies (Ryan *et al.* 1995; Lavigne 1996; Stockfors 2000; Maier 2001; Damesin *et al.* 2002; Bosc *et al.* 2003). We used two different approaches to determine whether or not xylem transport and storage of  $\text{CO}_2$  was responsible for this hysteresis. These two approaches produced seemingly contradictory results. The first experiment assumed *a priori* that the diel hysteresis was a function xylem  $\text{CO}_2$  transport. Therefore, night-time  $E_s$  temperature response functions, when  $v_s \approx 0$ , were used to predict

daytime rates ( $E_p$ ). Predicted daytime rates were always greater than observed. The difference between observed and predicted stem  $\text{CO}_2$  efflux ( $E_r = E_p - E_s$ ) was correlated with  $v_s$  and suggest that during the day high  $v_s$  could reduce  $E_s$  by up to 40% below that measured when  $v_s$  is low. These data support the idea that midday suppression of stem respiration measured in other studies (Negisi 1975, 1978, 1982; Lavigne 1987; Kabubari 1988) is a function of transpiration rate. Negisi (1979) artificially alter  $v_s$  in detached *Pinus densiflora* stems and found that  $E_s$  was reduced by 70% at a  $v_s$  rate of  $0.15 \text{ mm s}^{-1}$ , a value that corresponds well with our data (Fig. 3). Levy *et al.* (1999) also found a correspondence between  $E_r$  and  $v_s$ ; however, in their study, residual efflux was positively correlated with  $v_s$ . They concluded that imported  $\text{CO}_2$  in the xylem stream contributed up to 12% of  $E_s$ .

The second experiment examined how artificially reducing  $v_s$  by eliminating canopy leaf area affected  $E_s$ . Our attempt to create varying levels of stem  $v_s$  by a stepwise defoliation of the canopy was only partially successful. Increased stomatal conductance largely compensated for a




**Figure 5.** Least squares mean ( $\pm$  SE) of maximum daily sap velocity ( $v_s$ ) and stem surface  $\text{CO}_2$  efflux ( $E_s$ ) for non-fertilized and fertilized trees. Data are normalized to the maximum values measured on day of year (DOY) 109. Comparisons are made between trees with intact canopies (uncut) and those where the canopy was progressively removed (cut) in thirds over the course of the study. Each point is the average of three trees. An asterisk denotes a significant difference between uncut and cut means at  $\alpha = 0.05$ .

**Table 4.** Stomatal conductance ( $g_L$ ) and net photosynthesis ( $P_n$ ) of upper canopy foliage in non-fertilized (NF) and fertilized (F) trees with intact canopies [uncut (UC)] and those where the canopies were progressively removed [cut (C)] over the course of the study

	Treatment	108		114		117	
		$g_L$	$P_n$	$g_L$	$P_n$	$g_L$	$P_n$
NF	UC	48.0 (13.0)	3.6 (0.8)	17.0 (1.8)	3.4 (0.2)	30.0 (4.4)	5.6 (0.4)
	C	39.0 (7.0)	3.9 (0.7)	29.0 (2.6)	5.0 (0.4)	47.6 (5.9)	6.5 (0.3)
	UC	37.4 (1.0)	4.3 (0.2)	8.7 (0.6)	2.1 (0.2)	31.0 (4.1)	4.7 (0.2)
	C	44.3 (4.0)	5.0 (0.1)	32.0 (1.0)	4.9 (0.4)	55.0 (9.5)	5.5 (0.5)
F							

Measurements on day of year (DOY) 108 represent pre-cutting values whereas values on DOY 114 and 117 are after two-thirds of the canopy was removed in cut trees. Values are the mean of three trees. Numbers within parentheses are the SEs of the mean.



**Figure 6.** Comparison of stem surface  $\text{CO}_2$  efflux ( $E_s$ ), cambium temperature ( $T_c$ ), sap velocity ( $v_s$ ) and xylem  $\text{CO}_2$  concentration ( $X_s$ ) for a tree with an intact canopy (uncut) and one with the canopy removed (cut) in the (a) non-fertilized and (b) fertilized plots. Measurements are for the last 2 d of the study. The final cutting treatment was completed by 1000 h on day of year (DOY) 117.

partial reduction ( $\approx 33$  and  $63\%$ ) in canopy leaf area maintaining  $v_s$  near pre-treatment rates as observed in Pataki *et al.* (1998). Only after removal of most of the canopy ( $> 90\%$ ) was  $v_s$  substantially reduced. Even though we were unable to create a progressive reduction in  $v_s$ , it was evident that reducing  $v_s$  to near zero by removing almost all canopy foliage had little effect on  $E_s$ . Diel patterns of  $E_s$  and the daily maximum values were similar between cut and uncut trees throughout the experiment indicating that xylem CO<sub>2</sub> transport in the sap had little effect on  $E_s$  in these trees. In addition, if  $v_s$  strongly affected  $E_s$  then the magnitude of the diel hysteresis between  $E_s$  and  $T_c$  should be smaller or eliminated in cut trees as  $v_s$  approached zero. However, in our trees neither the pattern nor magnitude of the diel hysteresis was affected by large changes in  $v_s$  (Fig. 1). These data suggest that  $v_s$  and  $E_s$  are uncoupled in these trees and the *a priori* assumption that the diel hysteresis is a function of xylem CO<sub>2</sub> transport and storage is incorrect. Thus, the correlation between  $E_r/E_p$  and  $v_s$  (Fig. 3) does not represent a causative response.

The apparent uncoupling of  $v_s$  (or  $X_s$ ) and  $E_s$  suggest that that the radial diffusion of CO<sub>2</sub> from the xylem to the stem surface is restricted in these trees. Conifers have few intercellular spaces and radial gas diffusion must occur in the liquid phase which is several magnitudes lower than gas phase diffusion (Hari *et al.* 1991). The large difference between  $X_s$  and the ambient air in our trees indicates a high resistance to CO<sub>2</sub> diffusion from the xylem through the bark (Eklund 1990, 1993; Hari *et al.* 1991). Eklund (1990) and Eklund & Lavigne (1996) found little diffusion of O<sub>2</sub> or argon gas from the atmosphere to the xylem or from the xylem to the atmosphere in conifer stems. However, despite a high resistance to gas diffusion through the bark, several studies have shown that artificially manipulating of sap flow (Negisi 1979) and/or  $X_s$  (Teskey & McGuire 2002, 2004) clearly influences  $E_s$ . So, why did large changes in  $v_s$  and  $X_s$  in our cut trees have no effect on  $E_s$ ? We conducted the experiment in the spring when stem respiration and growth were at a maximum for these stands (Maier 2001). At this time of year, the thin cambium and phloem meristems likely respire at a much higher rate than the xylem parenchyma and thus would be a major source of respiratory CO<sub>2</sub> in the stem. Goodwin & Goddard (1940) measured oxygen consumption in black ash stems and found that O<sub>2</sub> uptake was several magnitudes higher in the cambium and phloem compared to the xylem. Similarly, Pruyne, Gartner & Harmon (2002, 2003) showed that the respiratory potential of the inner bark of ponderosa and white pine trees was 3–15 times greater than that of the sapwood. The surface CO<sub>2</sub> efflux we measured was likely a result of growth related respiration associated with differentiating cells in the cambium and from energy expended in phloem transport. Thus, during periods of rapid stem growth, the CO<sub>2</sub> concentration in the stems would be much higher in the cambium and phloem regions than in the xylem. Under these conditions, the CO<sub>2</sub> concentration gradient (i.e. diffusion gradient) would likely decrease from the cambium layer to the xylem effectively isolating the xylem tissue as a source of CO<sub>2</sub>

evolved from the stem surface. Although our data is limited, measurements of  $X_s$  appear to support this hypothesis. We found in our cut trees that while  $X_s$  increased after defoliation ( $v_s < 0.01 \text{ mm s}^{-1}$ ), it was not followed by increases in  $E_s$  which would be expected if the CO<sub>2</sub> concentration gradient decreased radially from the xylem to the stem surface. In addition, the rate and magnitude of  $E_s$  was similar on consecutive days with large differences in  $X_s$  (Fig. 6) suggesting that  $X_s$  had no effect on  $E_s$ . It is interesting to note that atmospheric O<sub>2</sub> in the xylem of conifer stems decreases from near ambient to less than 5% of ambient during the growing season (Eklund 1993, 2000); apparently, the active cambial tissue consumes most of the O<sub>2</sub> that diffuses into the stem (Hook *et al.* 1972).

Our measurements of  $E_s$  probably reflect respiration associated with diameter growth and phloem transport; however, it underestimates this component of stem respiration as some of the respired CO<sub>2</sub> is expended into the xylem (Eklund & Lavigne 1996; McGuire & Teskey 2004). This is illustrated by the increase in  $X_s$  at night when  $v_s \approx 0$  and is a result of the metabolism of local xylem parenchyma and from cambial tissue internal to the xylem. This CO<sub>2</sub> is stored in the stem segment and is later transported up the stem as part of the internal CO<sub>2</sub> flux. McGuire & Teskey (2004) used a mass balance approach to estimate total stem respiration from stem segments of several hardwood species. They found that CO<sub>2</sub> respiration within the stems either diffused radially through the bark to the atmosphere, is transported upward in the xylem stream, or is temporarily stored in the xylem. For example, surface flux, xylem transport flux and storage account for 85, 15 and 8%, respectively, of stem respiration over a 24 h period in a beech tree. The relative proportion of each component varied during the day depending on sap flow rates. In their analysis, dissolved CO<sub>2</sub> in the xylem served as a sink or a source for CO<sub>2</sub> diffusion to the atmosphere. Our data suggests that internal stem CO<sub>2</sub> dynamics are more complex. It appears that there may be conditions, perhaps during periods of high cambial activity, when  $E_s$  is uncoupled from internal xylem CO<sub>2</sub> fluxes. The relationship between  $v_s$ ,  $E_s$  and  $X_s$  may be different at other times of the year. For example when growth ceases,  $X_s$  may exceed that in the cambium and phloem regions and be a source of CO<sub>2</sub> to  $E_s$ . Root absorption of dissolved CO<sub>2</sub> in the soil can potentially contribute large amounts of CO<sub>2</sub> to the xylem (Levy *et al.* 1999). However, because of the porous nature of the sandy soils at our site, soil pCO<sub>2</sub> rarely exceeds 5000  $\mu\text{mol m}^{-2}$  (0.5%) in the top 50 cm and generally is much lower during the spring ( $\approx 1000$ –1100  $\mu\text{mol m}^{-2}$ , Maier, unpublished results) hence, soil CO<sub>2</sub> would not likely contribute much CO<sub>2</sub> to the xylem stream in our trees.

There are other possible explanations for the hysteresis between  $E_s$  and  $T_c$ : (1) a lag between temperature and surface CO<sub>2</sub> efflux because of high resistance to diffusion (Hari *et al.* 1991; Eklund & Lavigne 1996; Stockfors 2000); (2) refixation of respired CO<sub>2</sub> during cortical photosynthesis (Sprugel & Benecke 1991; Cernusak & Marshall 2000); (3) diel differences in substrate supply (Edwards &

McLaughlin 1978; Martin *et al.* 1994); and (4) diel patterns of stem growth. Because of the high resistance to CO<sub>2</sub> diffusion,  $T_c$  measured at the time of flux measurements may not reflect surface CO<sub>2</sub> evolution. Modelling  $E_s$  using lagged temperatures, measured sometime earlier, can account for a substantial portion of the hysteresis (Ryan *et al.* 1995; Lavigne 1996; Stockfors 2000; Maier 2001; Bosc *et al.* 2003). We modelled  $E_s$  (Eqn 1) for our trees using all of the data (day and night) and found that,  $E_s$  was best correlated ( $R^2 = 0.91\text{--}0.95$ ) with  $T_c$  measured between 42 and 168 min earlier. We note, that residuals from these regressions were not correlated with  $v_s$  ( $R^2 = 0.23, P = 0.45$ ). If a lag in CO<sub>2</sub> production were a major factor responsible for the hysteresis, then the magnitude of the hysteresis should be a function of tree size (Bosc *et al.* 2003) or bark thickness (Ryan *et al.* 1995). Although the range in tree size was small in our study, we found no relationship between lag times and tree diameter or bark thickness.

Bark photosynthesis can refix a substantial amount of respired CO<sub>2</sub> in woody tissue (Sprugel & Benecke 1991) thus lowering the apparent stem respiration rate during the day. Cernusak & Marshall (2000) estimated that bark refixation rates were 70% of night-time respiration rates in western white pine. However, refixation is not an issue in our study because we used the main stem that had little or no chlorophyll present and we used opaque chambers so there would be no bark photosynthesis at least for the tissue inside the chamber.

Substrate supply can affect respiration rates (Amthor 1989). Edwards & McLaughlin (1978) found that the diel pattern of  $E_s$  in yellow poplar trees was correlated with the concentration of reducing sugars in the phloem, indicating that the diel pattern of  $E_s$  may in part be driven by transported photosynthate. This response may occur quickly once carbohydrate supply is compromised. Edwards, Tschaplinski & Norby (2002) showed that  $E_s$  in sweetgum trees increased in response to elevated CO<sub>2</sub> but decreased to rates measured at ambient CO<sub>2</sub> within several days after the elevated CO<sub>2</sub> treatment was turned off. Removal of photosynthetic surface area would likely affect canopy assimilation and reduce substrate supply to the stem. We could not assess the impact of the cutting treatment on stem carbohydrate supply; however, the cutting treatment slightly reduced  $E_s$  in the fertilized trees. Effects of reduced carbohydrate supply on  $E_s$  could potentially mask effects of reduced  $v_s$ . Martin *et al.* (1994) found in loblolly pine seedlings that  $E_s$  declined after girdling the phloem above the respiration chamber; although, they concluded that the response was too slow to account for diel hysteresis.

The hysteresis between  $E_s$  and  $T_c$  may reflect diel patterns of growth. Daudet *et al.* (2005) measured diel  $E_s$  and stem diameter in potted hybrid walnut (*Juglans nigra* × *Juglans regia*) saplings under constant temperature conditions. Diel patterns of  $E_s$  were highly correlated with changes in stem diameter. Maximum  $E_s$  occurred at night, suggesting that more energy was being expended in growth and maintenance processes at this time.

## SUMMARY

Our trees showed a strong diel hysteresis between  $E_s$  and stem temperature. The diel variation in temperature-independent  $E_s$  was correlated with  $v_s$ , such that at high  $v_s$ ,  $E_s$  could be reduced by up to 40%. However, this correlation may not represent a cause-and-effect relationship. Artificially reducing  $v_s$  through a progressive defoliation of the canopy had little effect on  $E_s$  and had no effect on the magnitude of the diel hysteresis. These data indicate that  $E_s$  is uncoupled from  $v_s$  in these trees. We suggest that high metabolic activity in the cambium during this time of year (spring) is likely a source of CO<sub>2</sub> to the xylem and thus, CO<sub>2</sub> transported in the xylem stream would not contribute to  $E_s$ . This hypothesis is supported by the observation that diel changes in  $X_s$  correspond with  $E_s$  but the large increase in  $X_s$  measured in the cut trees, when  $v_s \approx 0$ , had no effect on  $E_s$ . Increased resolution of measurements of stem [CO<sub>2</sub>] in cambium and xylem regions is needed to confirm this. Understanding diel and seasonal variation in surface CO<sub>2</sub> efflux and the relationship to  $v_s$  (or sap flow) and  $X_s$  will provide a more complete characterization of stem respiration and whole-plant carbon cycles.

## ACKNOWLEDGMENTS

This work was supported in part through a Challenge Cost Share Grant from the USDA Southern Research Station. We would like to thank Pete Anderson and Daniel McInnis for technical assistance in the field, Karen Sarsony for laboratory analysis and Chelsea R. Maier for help with branch harvest and processing. We also would like to thank Dr Steve McKeand, North Carolina State University for permission to use the SETRES II site. We gratefully acknowledge critical pre-submission reviews by Mary Anne McGuire, Chelsy Ford, Mike Ryan and Robert Teskey.

## REFERENCES

- Amthor J.S. (1989) *Respiration and Crop Productivity*. Springer-Verlag, New York, USA.
- Bosc A., De Grandcourt A. & Loustau D. (2003) Variability of stem and branch maintenance respiration in a *Pinus pinaster* tree. *Tree Physiology* **23**, 227–236.
- Bowman W.P., Barbour M.M., Turnbull M.H., Tissue D.T., Whitehead D. & Griffin K.L. (2005) Sapflow rates and sapwood density are critical factors in within- and between-tree variation in CO<sub>2</sub> efflux from stems of mature *Dacrydium cupressinum* trees. *New Phytologist* **167**, 815–828.
- Butnor J.R., Johnsen K.H. & Maier C.A. (2005) Soil properties differentially influenced estimates of soil CO<sub>2</sub> efflux from three chamber based measurement systems. *Biogeochemistry* **73**, 283–301.
- Carey E.V., DeLucia E.H. & Ball J.T. (1996) Stem maintenance and construction respiration in *Pinus ponderosa* grown in different concentrations of atmospheric CO<sub>2</sub>. *Tree Physiology* **16**, 125–130.
- Cernusak L.A. & Marshall J.D. (2000) Photosynthetic refixation in branches of Western White Pine. *Functional Ecology* **14**, 300–311.
- Cienciala E. & Lindroth A. (1995) Gas exchange and sap flow measurements of *Salix viminalis* trees in short rotation forest.

- II. Diurnal and seasonal variation of stomatal response and water use efficiency. *Trees* **9**, 295–301.
- Clinton B.D., Maier C.A. & Sullivan N.H. (2001) An examination of the relationship between sapflow rate and measured stem CO<sub>2</sub> efflux in trees. The Ecological Society of America 86th Annual Meeting, Madison, WI, August 5–10.
- Damesin C., Ceschia E., Le Goff N., Ottorini J.M. & Dufrene E. (2002) Stem and branch respiration of beech: from tree measurements to estimations at the stand level. *New Phytologist* **153**, 159–172.
- Daudet F.A., Améglio T., Cochard H., Archilla O. & Lacointe A. (2005) Experimental analysis of the role of water and carbon in tree stem diameter variations. *Journal of Experimental Botany* **56**, 135–144.
- Edwards N.T. & McLaughlin S.B. (1978) Temperature-independent diel variations of respiration rates in *Quercus alba* and *Liriodendron tulipifera*. *Oikos* **31**, 200–206.
- Edwards N.T. & Wullschleger S.D. (2000) Carbon dioxide efflux rates from stems of mature *Quercus prinus* L. & *Acer rubrum* L. trees do not appear to be affected by sapflow rates. *Journal of Sustainable Forestry* **10**, 125–131.
- Edwards N.T., Tschaplinski T.J. & Norby R.J. (2002) Stem respiration increases in CO<sub>2</sub>-enriched sweetgum trees. *New Phytologist* **155**, 239–248.
- Eklund L. (1990) Endogenous levels of oxygen, carbon dioxide and ethylene in stems of Norway spruce trees during one growing season. *Trees* **4**, 150–154.
- Eklund L. (1993) Seasonal variations of O<sub>2</sub>, CO<sub>2</sub>, and ethylene in oak and maple stems. *Canadian Journal of Forest Research* **23**, 2608–2610.
- Eklund L. (2000) Internal oxygen levels decrease during the growing season and with increasing stem height. *Tree* **14**, 177–180.
- Eklund L. & Lavigne M.B. (1996) Restricted lateral gas movement in *Pinus strobus* branches. *Trees* **10**, 83–85.
- Ewers B.E. & Oren R. (2000) Analysis of assumptions and errors in the calculation of stomatal conductance from sap flux measurements. *Tree Physiology* **20**, 579–589.
- Ewers B.E., Oren R. & Sperry J.S. (2000) Root hydraulic conductance: a reflection of water balance and a constraint on canopy stomatal conductance. *Plant, Cell & Environment* **23**, 1055–1066.
- Ewers B.E., Oren R., Johnsen K.H. & Landsberg J.J. (2001) Estimating maximum mean canopy stomatal conductance for use in models. *Canadian Journal of Forest Research* **31**, 198–207.
- Goodwin R.H. & Goddard D.R. (1940) The oxygen consumption of isolated woody tissues. *American Journal of Botany* **27**, 234–237.
- Granier A. (1985) Une nouvelle méthode pour la mesure du flux de sève brute dans le tronc des arbres. *Annales Des Sciences Forestières* **42**, 193–200.
- Granier A. (1987) Evaluation of transpiration in a Douglas-fir stand by means of sap flow measurements. *Tree Physiology* **3**, 309–320.
- Hari P., Pekka P. & Korpilahti E. (1991) Internal circulation of carbon within a tree. *Canadian Journal of Forest Research* **21**, 514–515.
- Hook D.D., Brown C.L. & Wetmore R.H. (1972) Aeration in trees. *Botanical Gazette* **133**, 443–454.
- Kabubari Y. (1988) Diurnal and seasonal fluctuations in the bark respiration of standing *Fagus sylvatica* trees at Solingen, West Germany. *Journal of the Japanese Forestry Society* **70**, 64–70.
- Kaiplainen L.K., Sofronova G.I., Hari P. & Yalinskaya E.E. (1998) The role of xylem in CO<sub>2</sub> exchange in *Pinus sylvestris* woody stems. *Russian Journal of Plant Physiology* **45**, 500–505.
- Lavigne M.B. (1987) Differences in stem respiration response to temperature between balsam firs trees in thinned and unthinned stands. *Tree Physiology* **3**, 225–233.
- Lavigne M.B. (1996) Comparing stem respiration and growth of jack pine provenances from northern and southern locations. *Tree Physiology* **16**, 847–852.
- Levy P.E., Meir P., Allen S.J. & Jarvis P.G. (1999) The effect of aqueous transport of CO<sub>2</sub> in xylem sap on gas exchange in woody plants. *Tree Physiology* **19**, 53–58.
- McGuire M.A. & Teskey R.O. (2002) Microelectrode technique for in situ measurement of carbon dioxide concentrations in xylem sap of trees. *Tree Physiology* **22**, 807–811.
- McGuire M.A. & Teskey R.O. (2004) A method for estimating stem respiration in trees using a mass balance approach that accounts for internal and external fluxes of CO<sub>2</sub>. *Tree Physiology* **24**, 571–578.
- McKeand S.E., Grissom J.E., Handest J.A., O'Malley D.M. & Allen H.L. (2000) Responsiveness of diverse provenances of loblolly pine to fertilization – age 4 results. *Journal of Sustainable Forestry* **10**, 87–94.
- Maier C.A. (2001) Stem growth and respiration in loblolly pine plantations differing in soil resource availability. *Tree Physiology* **21**, 1183–1193.
- Maier C.A., Albaugh T.J., Allen H.L. & Dougherty P.M. (2004) Respiratory carbon use and storage in mid-rotation loblolly pine (*Pinus taeda*) plantations: the effect of site resources on the stand carbon balance. *Global Change Biology* **10**, 1335–1350.
- Maier C.A., Zarnoch S.J. & Dougherty P.M. (1998) Effects of temperature and tissue nitrogen on dormant season stem and branch maintenance respiration in a young loblolly pine (*Pinus taeda*) plantation. *Tree Physiology* **18**, 11–20.
- Martin T.A., Teskey R.O. & Dougherty P.M. (1994) Movement of respiratory CO<sub>2</sub> in stems of loblolly pine (*Pinus taeda*) seedlings. *Tree Physiology* **14**, 481–495.
- Negisi K. (1975) Diurnal fluctuations of CO<sub>2</sub> released from the stem of standing young *Pinus densiflora* trees. *Journal of Japanese Forestry Society* **57**, 375–383.
- Negisi K. (1978) Daytime depression in bark respiration and radial shrinkage in stem of a standing young *Pinus densiflora* tree. *Journal of Japanese Forestry Society* **60**, 380–382.
- Negisi K. (1979) Bark respiration rate in stem segments detached from young *Pinus densiflora* trees in relation to velocity of artificial sap flow. *Journal of Japanese Forestry Society* **61**, 88–93.
- Negisi K. (1982) Diurnal fluctuations of the stem bark respiration in relationship to the wood temperature in standing young *Pinus densiflora*, *Chamaecyparis obtuse*, and *Quercus myrsinaefolia* trees. *Journal of Japanese Forestry Society* **64**, 315–319.
- Oren R., Phillips N., Ewers B.E., Pataki D.E. & Megonigal J.P. (1999) Sap-flux-scaled transpiration response to light, vapor pressure deficit, and leaf area reduction in a flooded *Taxodium distichum* forest. *Tree Physiology* **19**, 337–347.
- Pataki D.E., Oren R. & Phillips N. (1998) Responses of sap flux and stomatal conductance of *Pinus taeda* L. trees to stepwise reductions in leaf area. *Journal of Experimental Botany* **49**, 871–878.
- Pruyn M.L., Gartner B.L. & Harmon M.E. (2002) Respiratory potential in sapwood of old versus young ponderosa pine trees in the Pacific Northwest. *Tree Physiology* **22**, 105–116.
- Pruyn M.L., Harmon M.E. & Gartner B.L. (2003) Stem respiratory potential in six softwood and four hardwood tree species in the central cascades of Oregon. *Oecologia* **137**, 10–21.
- Ryan M.G., Linder S., Vose J.M. & Hubbard R.M. (1994) Dark respiration in pines. *Ecological Bulletins* **43**, 50–63.
- Ryan M.G., Gower S.T., Hubbard R.M., Waring R.H., Ghosh H.L., Cropper W.P. Jr. & Running S.W. (1995) Woody tissue maintenance respiration of four conifers in contrasting climate. *Oecologia* **101**, 133–140.
- Ryan M.G., Hubbard R.M., Pongracic S., Raison R.J. & McMurtrie R.E. (1996) Foliage, fine root, woody tissue and stand

- respiration in *Pinus radiata* in relation to nitrogen status. *Tree Physiology* **16**, 333–343.
- Sala A., Smith S.D. & Devitt D.A. (1996) Water use by *Tamarix ramosissima* and associated phreatophytes in a Mojave Desert floodplain. *Ecological Applications* **6**, 888–898.
- Sprugel D.G. & Benecke U. (1991) Measuring woody-tissue respiration and photosynthesis. In *Techniques & Approaches in Forest Tree Ecophysiology* (eds J.P. Lassoie & T.M. Hinckley), pp. 329–355. CRC Press, Boca Raton, FL, USA.
- Stockfors J. (2000) Temperature variations and distribution of living cells within tree stems: implications for stem respiration modeling and scale-up. *Tree Physiology* **20**, 1057–1062.
- Stringer J.W. & Kimmerer T.W. (1993) Refixation of xylem sap CO<sub>2</sub> in *Populus deltoids*. *Physiologia Plantarum* **89**, 243–251.
- Teskey R.O. & McGuire M.A. (2002) Carbon dioxide transport in xylem causes errors in estimation of rates of respiration in stems and branches of trees. *Plant, Cell & Environment* **25**, 1571–1577.
- Teskey R.O. & McGuire M.A. (2005) CO<sub>2</sub> transported in xylem sap affects CO<sub>2</sub> efflux from *Liquidambar styraciflua* and *Platanus occidentalis* stems, and contributes to observed wound respiration phenomena. *Trees* **19**, 357–362.
- Tyree M.T. & Ewers F.W. (1991) The hydraulic architecture of trees and other woody plants. Tansley Review No. 34. *New Phytologist* **119**, 345–390.
- Xu M., DeBiase T., Qi Y., Goldstein A. & Liu Z. (2001) Ecosystem respiration in a young Ponderosa pine plantation in the Sierra Nevada Mountains, California. *Tree Physiology* **21**, 309–318.

Received 3 January 2006; accepted for publication 8 March 2006