

#78

PHORETIC CARRYING CAPACITY OF FLYING SOUTHERN PINE BEETLES (COLEOPTERA: SCOLYTIDAE)

JOHN C. MOSER

Southern Forest Experiment Station, USDA Forest Service, Pineville, Louisiana

Mites do not have wings, but in their course of evolution many species have developed an association with insects, using them as a vehicle of distribution. Occasionally they cover the host so completely that the insect cannot fly. The literature is replete with these observations. Except for a single speculation (Fronk 1947), there are no reports as to how many mites *Dendroctonus frontalis* Zimmerman can support and still fly to its intended destination.

Of 2539 southern pine beetles trapped in flight with Stickem Special^{®1} (Moser 1976), 23.3% possessed at least one deutonymph of *Trichouropoda australis* Hirschmann (Fig. 1), one of the larger mites associated with *D. frontalis*. The mean number of *T. australis* per infested beetle was 4.9 for male beetles and 5.1 for females. The greatest number of these mites found on a single infested beetle was 64 deutonymphs on a male beetle and 41 on a female.

When 51 live *T. australis* deutonymphs were extracted from laboratory-reared beetles and immediately weighed on a Cahn[®] electrobalance, they averaged 0.0064 mg each. Barras and Hodges (1974) showed that emerging males and females of *D. frontalis* weighed about 2.07 and 2.14 mg each, respectively. Hence, an average




FIG. 1. Adult southern pine beetle with deutonymph of *Trichouropoda australis* attached by anal pedicel (A).

¹Mention of trade name is solely to identify material used and does not imply endorsement by the U.S. Department of Agriculture.

male carrying 64 deutonymphs is supporting about a 20% additional load, and a female with 41 deutonymphs, a 12% additional load. Average additional loads supported by both sexes were ~~1.5%~~ 15%.

The critical weight that *D. frontalis* can sustain in flight is still unknown, but Fronk's (1947) contention that *D. frontalis* cannot fly with over 40 uropodids attached seems wrong. Although Fronk's specimens were taken from Virginia, we can reasonably assume that he was dealing with the same uropodid. *T. australis* not only occurs there but is the only uropodid associated with *D. frontalis* that attaches in such large numbers.

References

- Barras, S. J. and J. D. Hodges. 1974. Weight, moisture, and lipid changes during life cycle of the southern pine beetle. *U.S. Dep. Agric. For. Serv. Res. Note* SO-178. South. For. Exp. Stn., New Orleans, La. 5 pp.
- Fronk, W. D. 1947. The southern pine beetle, its life history. *Tech. Bull. Va agric. Exp. Stn* 108. Blacksburg, Va. 12 pp.
- Moser, J. C. 1976. Surveying mites (Acarina) phoretic on the southern pine beetle (Coleoptera: Scolytidae) with sticky traps. *Can. Ent.* **108**: 809-813.

(Received 6 May 1975)