

Reprinted from

# Forest Ecology and Management

---

Forest Ecology and Management 73 ( 1995 ) 2 1 1-2 19

## Basal area growth for 15 tropical tree species in Puerto Rico

**Bernard R. Parresol<sup>1</sup>**

*Institute for Quantitative Studies, USDA Forest Service, Southern Forest Experiment Station, Room T-10210 USPS Bldg.,  
701 Loyola Ave., New Orleans, LA 70113, USA*

Accepted 6 September 1994



# Forest Ecology and Management

---

**Aims and scope.** Forest *Ecology and Management* publishes scientific articles concerned with forest management and conservation, and in particular the application of biological, ecological and social knowledge to the management of man-made and natural forests. The scope of the journal includes all forest ecosystems of the world. A refereeing process ensures the quality and international interest of the manuscripts accepted for publication. The journal aims to encourage communication between scientists in disparate fields who share a common interest in ecology and natural-resource management, and to bridge the gap between research workers and forest managers in the field to the benefit of both. The journal should be of interest to research workers, managers and policy makers in forestry, natural resources, ecological conservation and related fields.

## FOUNDING EDITOR

Laurence L. Roche, Murroe, Ireland

## EDITORS-IN-CHIEF

For the Americas, Australia,  
New Zealand and the Pacific:

R.F. Fisher  
Department of Forest Science, Texas A&M University  
College Station, TX 77843-2135, USA

For the rest of the world:

E.P. Farrell  
Environmental Resource Management  
University College Dublin, Belfield  
Dublin 4, Ireland

## BOOK REVIEW EDITOR

D. Binkley, Forest and Wood Science, Colorado State University, Fort Collins, CO 80523, USA

## EDITORIAL ADVISORY BOARD

G. Abrahamsen, Agricultural University of Norway, Ås, Norway

M.R. Ahuja, Institute of Forest Genetics & Forest,

Grosshansdorf, Germany

F. Andersson, Swedish University of Agricultural Sciences,

Uppsala, Sweden

M.G.R. Cannell, Institute of Terrestrial Ecology, Penicuik,  
Midlothian, UK

J.D. Deans, Institute of Terrestrial Ecology, Penicuik,  
Midlothian, UK

J.S. de Queiroz, Utah State University, Logan, UT, USA

D.P. Dykstra, CIFOR, Jakarta, Indonesia

M.J. Foxe, National Centre for Agriculture & Veterinary  
Biotechnology, Dublin, Ireland

P.H. Freer-Smith, Forestry Commission, Farnham, UK

R.A. Goyer, Louisiana State University, Baton Rouge, LA, USA

J.B. Hall, University College of North Wales, Bangor, UK

J.J. Havel, Havel Land Consultants, Wanneroo, W.A.,  
Australia

P. Höglberg, University of Umeå, Umeå, Sweden

E. Klimo, University of Agriculture, Brno-Sobesice, Czech  
Republic

B.M. Kumar, Kerala Agricultural University, Kerala, India

J.P. Lanly, F.A.O., Forest Resources Division, Rome, Italy

J.N. Long, Utah State University, Logan, UT, USA

A.E. Lugo, International Institute of Tropical Forestry,  
Rio Piedras, PR, USA

J.A. Maghembe, SADCC/ICRAF Agroforestry Project,  
Zomba, Malawi

F. Makeschin, Martin Luther Universität, Halle, Germany

D.C. Malcolm, University of Edinburgh, Edinburgh, UK

E. Mäikönen, Finnish Forest Research Institute, Vantaa,  
Finland

J.A. McLean, University of British Columbia, Vancouver,  
B.C., Canada

G.M.J. Mohren, Institute for Forestry and Nature Research  
(IBN-DLO), Wageningen, Netherlands

R. Morandini, Istituto Sperimentale per la Selvicoltura,  
Arezzo, Italy

M.A.R. Nahuz, Instituto de Pesquisas Tecnológicas,  
São Paulo, SP, Brazil

R.F. Powers, Pacific Southwest Research Station, Redding,  
CA, USA

P.S. Ramakrishnan, Jawaharlal Nehru University,  
New Delhi, India

L. Rasmussen, Danish Forest and Landscape Research  
Institute, Lyngby, Denmark

R. Sands, School of Forestry, Creswick, Vic., Australia

J.A. Stanturf, Stoneville, MS, USA

O. Sziklai, University of British Columbia, Vancouver,  
B.C., Canada

K. von Weissenberg, University of Joensuu, Joensuu,  
Finland

---

**Publication information:** Forest *Ecology and Management* (ISSN 0378-1127). For 1995 volumes 71-79 are scheduled for publication. Subscription prices are available upon request from the Publisher. Subscriptions are accepted on a prepaid basis only and are entered on a calendar year basis. Issues are sent by surface mail except to the following countries where Air delivery via SAL mail is ensured: Argentina, Australia, Brazil, Canada, Hong Kong, India, Israel, Japan, Malaysia, Mexico, New Zealand, Pakistan, PR China, Singapore, South Africa, South Korea, Taiwan, Thailand, USA. For all other countries airmail rates are available on request. Claims for missing issues should be made within six months of our publication (mailing) date. Please address all your requests regarding orders and subscription queries to: **Elsevier Science B.V., Journal Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, tel. (+31-20) 4853642, fax (+31-20) 4853598.**

*In the USA and Canada:* For further information on this and other Elsevier journals please contact: **Elsevier Science Inc., Journal Information Center, 655 Avenue of the Americas, New York, NY 10010, USA. Tel. (212) 6333750; fax (212) 6333764; telex 420-643 AEP UI.**

**Back volumes:** Please contact the Publisher.

## Basal area growth for 15 tropical tree species in Puerto Rico

Bernard R. Parresol<sup>1</sup>

Institute for Quantitative Studies, USDA Forest Service, Southern Forest Experiment Station, Room T-10210 USPS Bldg.,  
701 Loyola Ave., New Orleans, LA 70113, USA

Accepted 6 September 1994

### Abstract

The tabonuco forests of Puerto Rico support a diverse population of tree species valued for timber, fuel, food, wildlife food and cover, and erosion control among other uses. Tree basal area growth data spanning 39 years are available on 15 species from eight permanent plots in the Luquillo Experimental Forest. The complexity of the rain forest challenges current forest stand modeling techniques. As a starting point individual tree basal area growth is modeled using the Chapman-Richards function constrained for hypothetical maximum tree size. In addition to initial tree diameter or basal area, significant explanatory variables are crown class, topographic position and degree of ground incline. Plots illustrate the differing growth patterns of the 15 tropical mixed/moist forest species. Two species exhibit exceptional growth. *Buchenavia capitata* (Vahl) Eichl. has basal area growth peaking at 87 cm<sup>2</sup> year<sup>-1</sup>. The *Manilkara bidentata* (A. DC.) A. Chev. data show growth rates in excess of 60 cm<sup>2</sup> year<sup>-1</sup> and the Chapman-Richards function indicates growth potential to a peak of 122 cm<sup>2</sup> year<sup>-1</sup>.

**Keywords:** Tabonuco; Chapman-Richards function; Periodic annual increment; Crown class; Topographic position; Degree of ground incline

### 1. Introduction

Stand dynamics within tropical mixed/moist forests are the most complex in the world. Studies in the tropics of the eastern hemisphere have led to diverse and generally discouraging conclusions concerning growth prediction. Lowe ( 1966, 1973 ) concluded that differences in growth rates are only partly due to contemporary influences because trees develop a 'pecking order' while still young. Gerrard ( 1968 ) saw little practical potential for the individual tree approach to growth prediction in stands containing a diversity of species and age classes. Mervart ( 1969, 1972 )

concluded that the process of growth was sufficiently random to frustrate any attempt to interpret it in deterministic terms, and that the possibility of reducing the error of increment estimates via regression analysis of the girth-increment relationship was quite limited.

In the neo-tropics an early study by Wadsworth ( 1953 ), relating tree growth to prior tree size, crown vigor, and surrounding stand density, found only a weak correlation. However, a more recent study by Wadsworth et al. (1989) points to a number of tree increment indicators with potential for accurately predicting individual tree breast height diameter (dbh) or basal area (*ba*) growth. The best indicators were crown diameter (*cd*) and *cd/dbh* ratio in conjunction

<sup>1</sup> Tel. 504-589-4547; Fax. 504-589-3961.

with initial dbh or *ba*. Unfortunately, most tropical tree **datasets** do not contain crown diameter, but based on the work of Wadsworth et al. this variable should be included in future inventory and measurement work. In the present work crown diameter at the beginning of the growth period was not available. The objective of this study was to investigate other explanatory variables and construct equations for predicting tree

basal area growth for 15 of the more commercially and ecologically important species of **ta**-bonuco forests in Puerto Rico. These equations should provide a starting point for more detailed investigations of stand and forest growth. It is believed that these 15 species should display similar, though not exact, growth patterns elsewhere in the **tropics**.




Fig. 1. Location of selected long term growth plots within the Luquillo Experimental Forest in northeastern Puerto Rico. Dashed line in inset represents forest boundary.

Table 1  
Descriptive statistics on study sites within the Luquillo Experimental Forest, Puerto Rico

Site"	Average elevation (m)	Prevailing aspect	Mean slope (deg.)
1	440	S80E	31
2	500	N75E	37
3	550	N55E	35
5	450	N80E	22
6	640	N70E	16
7	480	N65E	25
8	470	S65E	39
9	200	N60E	33

"Site 4 was destroyed during road construction.

## 2. Study sites

The availability of many hundreds of repeatedly measured trees from eight sites (nine originally established, site 4 destroyed during road construction) within a 20 km<sup>2</sup> area of the Luquillo Experimental Forest in Puerto Rico (Fig. 1) presented an opportunity to model individual tree growth. Sites are briefly described in Table 1. Latitude is 18 ° N and the elevation of the forests studied range from 200 to 640 m above sea level. The climate, frost-free with rainfall ranging from 250 to 450 cm annually, is classified as subtropical wet and rain forest life zones (sensu Holdridge, 1967). Soils are acid clays and loams, mostly ultisols and inceptisols. The tabonuco forests (so-called because the tabonuco tree (*Dacryodes excelsa* Vahl) is the most prominent species) used are both primary and late secondary, reaching 30 m in height.

## 3. Data

Permanent remeasurement plots were established in the Luquillo Experimental Forest in 1943. Diameter and crown class (dominant, co-dominant, intermediate, and suppressed) of all trees were measured (as well as other variables) and for each plot topography (ridge, slope, bottom), aspect (recorded as azimuth), and incline (greatest downward slant, degrees from horizon-

tal) were noted. Trees were remeasured for dbh in 1947, 1952, 1958, 1965, 1976, and 1982, resulting in 39 years of growth measurement. From the pool of data 15 species had sufficient numbers and size range for modeling tree basal area growth. These species were: ( 1) *Dacryodes excelsa* Vahl; ( 2) *Tabebuia heterophylla* (DC. ) Britton; ( 3) *Ormosia krugii* Urban; ( 4) *Buchenavia capitata* (Vahl) Eichl.; ( 5) *Manilkara bidentata* (A. DC. ) A. Chev.; ( 6) *Matayba dominicensis* (DC. ) Radlk.; ( 7 ) *Didymopanax morototoni* (Aubl.) Decne. & Planch.; ( 8) *Cecropia peltata* L.; ( 9) *Inga fastuosa* (L.) Willd.; ( 10) *Homalium rucemosum* Jacq.; ( 11) *Sloanea berteriana* Choisy; ( 12) *Linociera dominicensis* (Lam. ) Knobl.; ( 13) *Tetragastris balsamifera* (Sw. ) Kuntze; ( 14) *Byrsonima coriacea* (Sw.) DC.; ( 15) *Guarea guidonia* (L. ) Sleumer. The tree data are summarized in Table 2.

## 4. Methods

If one wishes to obtain an accurate assessment of growth pattern over time, the tree should be measured at short intervals. Therefore I computed periodic annual increments (PAI) for each time interval in the data. That is, I started at interval 1943– 1947 and for each tree with a diameter at both time periods I computed a periodic annual basal area increment (*ba* growth divided by number of years). I then computed PAI for the intervals 1947–1952, 1952–1958, 1958–1965, 1965–1976, and 1976–1982. If for a tree a measurement was not made at a measurement period then the PAI was coded as missing for the intervals involved.

Because of its widespread use and familiarity, I settled on the Chapman-Richards growth function to express periodic annual basal area growth. Under this model (Pienaar and Turnbull, 1973 ) the growth rate of an organism is expressed as the difference between constructive metabolism (anabolic growth) and destructive metabolism (catabolic growth). This function, constrained to equal zero at theoretical maximum tree size

Table 2

Mean sample tree diameters (cm) ( $\pm$  SD) of 15 species in a moist tropical forest in Puerto Rico for seven time periods

Species	1943			1947			1952			1958			1965			1976			1982		
	$\bar{x}$	SD	<i>n</i>																		
<i>Dacryodes excelsa</i> Vahl	19.2	12.6	101	27.2	19.3	214	29.4	19.8	112	25.2	13.1	329	33.6	19.4	211	35.1	16.0	138	37.7	17.3	359
<i>Tabebuia heterophylla</i> (DC.) Britton	6.7	1.3	5	11.3	3.3	37	15.0	4.0	35	15.8	4.9	280	17.3	5.0	170	22.1	6.7	231	23.4	7.4	287
<i>Ormosia krugii</i> Urban	11.5	6.7	8	14.9	8.7	37	18.9	9.3	37	20.7	8.5	107	24.2	9.2	61	29.2	9.2	63	31.6	9.7	110
<i>Buchenavia capitata</i> (Vahl) Eichl.	22.2	12.9	6	28.7	15.1	16	35.9	15.3	13	28.5	14.7	45	33.9	15.3	20	41.4	19.9	24	44.1	19.2	50
<i>Manilkara bidentata</i> (A. DC.) A. Chev.	11.5	4.6	47	13.1	5.9	58	15.4	6.3	56	17.9	6.7	115	22.3	8.3	54	30.0	9.6	35	29.1	10.3	119
<i>Matayba domingensis</i> (DC.) Radlk.			0	12.0	5.7	58	15.1	6.0	57	17.2	5.9	142	20.4	7.3	30	24.2	6.8	63	26.0	7.1	142
<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	18.7	7.9	7	13.0	6.0	30	14.9	6.2	27	17.3	5.7	121	20.3	5.9	75	23.5	6.9	76	25.3	8.3	123
<i>Cecropia peltata</i> L.	17.8	14.5	3	18.2	17.4	11	20.7	16.0	12	20.0	7.4	60	28.9	12.2	22	30.6	10.8	25	32.0	12.5	61
<i>Inga fastuosa</i> (L.) Willd.	17.0	5.3	3	12.8	3.3	8	15.6	4.2	5	17.5	7.1	62	18.3	6.2	39	26.6	9.3	53	28.4	9.9	65
<i>Homalium racemosum</i> Jacq.	6.1	-	1	17.9	12.5	10	19.4	12.3	10	17.6	9.3	38	18.7	6.1	10	26.2	9.4	26	31.8	15.5	42
<i>Sloanea berteriana</i> Choisy	12.6	7.9	14	15.3	9.8	20	18.4	10.1	17	20.0	10.5	40	19.7	7.8	16	24.7	9.6	17	27.6	15.1	44
<i>Linociera domingensis</i> (Lam.) Knobl.	7.3	4.6	4	9.5	2.8	12	11.6	2.9	9	15.2	5.3	41	18.0	-	1	23.2	8.5	19	23.5	7.5	43
<i>Tetragastris balsamifera</i> (Sw.) Kuntze	19.7	12.7	15	21.4	13.1	15	23.7	13.0	15	25.9	13.4	16	27.5	15.0	13	37.8	-	1	33.0	14.7	16
<i>Byrsonima coriacea</i> (Sw.) DC.	5.9	1.0	4	6.9	1.9	4	-	-	0	13.0	3.0	19	16.1	2.9	14	25.0	8.5	15	29.9	10.3	22
<i>Guarea guidonia</i> (L.) Sleumer			0-			0	12.9	2.3	2	18.1	5.9	10	22.7	6.4	10	32.7	7.9	8	33.2	10.7	10

Table 3

Asymptotic tree basal area, regression coefficients and associated fit statistics from growth models for 15 species in a moist tropical forest in Puerto Rico

Species	A	b <sub>1</sub>	b <sub>2</sub>	b <sub>3</sub>	b <sub>4</sub>	b <sub>5</sub>	R <sup>2</sup>	RMSE
<b>Eq. (1)</b>								
<i>Tabebuia heterophylla</i>	3849	0.31983	0.89421				0.13	8.42
<i>Manilkara bidentata</i>	31416	0.30372	0.93193				0.64	8.53
<i>L. fastuosa</i>	3849	0.645	14	0.95038			0.12	13.70
<i>L. domingensis</i>	5027	0.27013	0.89684				0.40	5.74
<b>Eq. (2)</b>								
<i>Dacryodes excelsa</i>	31416	5.94762	0.28608	-0.28976			0.35	12.30
<i>O. krugii</i>	5027	0.45568	0.81041	-0.06587			0.34	10.27
<i>Matayba domingensis</i>	2828	1.78329	0.57596	-0.20575			0.36	7.31
<i>Tetragastris balsamifera</i>	7854	4.51702	0.43270	-0.45533			0.69	8.17
<b>Eq. (3)</b>								
<i>Buchanavia capitata</i>	17672	1.29601	0.72459	-3.01103	0.54071	0.50262	0.51	22.18
<i>Didymopanax morototoni</i>	2828	0.53916	0.67774	0.23697	-0.76755	-0.99593	0.13	8.92
<i>C. peltata</i>	5027	3.78875	0.28512	1.98154	-2.45468	0.46728	0.30	17.91
<i>H. racemosum</i>	7854	0.37467	0.72026	-1.31511	2.22638	2.93882	0.50	10.20
<i>S. berteriana</i>	11310	0.13564	0.77037	2.32871	-0.60188	0.46750	0.31	10.46
<i>Byrsonima coriacea</i>	2828	0.70093	0.56782	0.90109	0.60853	0.0	0.44	5.25
<i>G. guidonia</i>	13274	0.69385	0.65432	-0.24530	0.30781	1.14116	0.32	19.84

Note: A is asymptote value (cm<sup>2</sup>); b<sub>i</sub>'s are regression coefficients; R<sup>2</sup> is the variation explained in the dependent variable, calculated as 1 - Σ (y<sub>i</sub> - ŷ) / Σ (y<sub>i</sub> - ȳ); and RMSE is root mean square error (cm<sup>2</sup>).

\*There were no trees occurring in a bottom type topography for this species.

(Shifley and Brand, 1984), is

$$\Delta X / \Delta t = b_1 (X^{b_2} - X/A^{(1-b_2)}) \tag{1}$$

where ΔX/Δt is PAI, X is tree basal area ( cm<sup>2</sup> ), t is time (for this study Δt is always 1 year), A is asymptotic tree basal area-the point where basal area growth equals zero, and b<sub>1</sub> and b<sub>2</sub> are equation coefficients. This function is usually presented as a differential equation; however, the discrete nature of the data makes a difference equation more appropriate.

Additional variables available for modeling were crown class (CC) (Smith, 1962 ), degree of incline (INCL), topography (TOPO), and aspect. I ran a series of multiple nonlinear regressions using initial basal area and various combinations of the above variables to predict PAI. I found that CC (coded as: 1, dominant; 2, co-dominant; 3, intermediate; 4, suppressed) and the interaction of INCL with the class variable TOPO improved the prediction for some spe-

cies. Therefore, two additional models were formulated

$$\Delta X / \Delta t = b_1 (X^{b_2} - X/A^{(1-b_2)}) \exp(b_3 CC) \tag{2}$$

$$\Delta X / \Delta t = b_1 (X^{b_2} - X/A^{(1-b_2)}) \exp [ (b_3 + b_4 I_1 + b_5 I_2) INCL / 100 ] \tag{3}$$

where INCL is degrees from the horizontal (i.e. in the range 0-90) and I<sub>1</sub> = 1 if TOPO is classified as slope else I<sub>1</sub> = 0 and I<sub>2</sub> = 1 if TOPO is classified as bottom else I<sub>2</sub> = 0. If both I<sub>1</sub> and I<sub>2</sub> equal zero then we are left with b<sub>3</sub> which is the effect of ridge topography. The coefficient b<sub>4</sub> measures the differential effect between ridge and slope topography and b<sub>5</sub> measures the differential effect between ridge and bottom topography. If inclinkis measured in percent, then values can be converted to degrees for use in Eq. (3) with the formula: degrees = cotan (percent/ 100).




Fig. 2. Comparative growth rate curves of 15 species in a mixed/moist tropical forest in Puerto Rico. Species were grouped into (A) small, (B) medium, (C) large, and (D) very large tree sizes. Solid lines show extent of actual data, dashed lines are extrapolations.

## 5. Results and discussion

The data from each species were fitted to the three equations via nonlinear least squares and residuals were examined for departures from regression assumptions. The most appropriate equation for a given species was determined from the residual analyses as well as the variation explained in the dependent variable ( $R^2$ ). The equation giving the highest  $R^2$  was not necessarily judged as best though in most cases (13 of 15) this held true. Table 3 lists asymptote values (F.H. Wadsworth, International Institute of

Tropical Forestry, USDA Forest Service, Rio Piedras, Puerto Rico, personal communication, 1994), equation coefficients, and lit statistics. Four species (*I. fastuosa*, *L. domingensis*, *Manilkara bidentata*, and *Tabebuia heterophylla*) did best under Eq. (1), the standard Chapman-Richards function. Additional variables did not improve predictions. For four species (*Dacryodes excelsa*, *M. domingensis*, *O. krugii*, *Tetragastris balsamifera*), the addition of CC provided sufficient improvement to warrant inclusion. Finally, for the remaining seven species, the TOPO by INCL interaction contributed




Fig. 3. Illustration of growth rate response across crown class. Species is *Tetragastris balsamifera* in a moist tropical forest in Puerto Rico. Solid lines show extent of actual data, dashed lines are extrapolations.

enough explanatory power to justify its use.

Plots illustrating comparative growth rates of these species are given in Fig. 2. Species were grouped by their asymptote values as (A) small, (B) medium, (C) large, and (D) very large. For species under Eq. (2) the dominant crown class was used. This shows the best response for those species. For the eight sites the overall average degree of incline was close to 30 so this value was used for INCL in Eq. (3) coupled with slope topography ( $I_1 = 1, I_2 = 0$ ). Keep in mind that for the seven species under Eq. (3) plotted in Fig. 2, the curves shown are one of many possible responses. The growth rate of each species is shown by a solid line followed by a dashed line which shows the extent of the actual data and extrapolation to the asymptote, respectively. One general trend that is immediately obvious is that the maximum growth rate range from group A to group D steadily increases. For group A maximum rates range from 14 to 31  $\text{cm}^2 \text{ year}^{-1}$ , for group B from 16 to 42  $\text{cm}^2 \text{ year}^{-1}$ , for group C from 29 to 87  $\text{cm}^2 \text{ year}^{-1}$ , and for group D from 37 to 122  $\text{cm}^2 \text{ year}^{-1}$ . All species exhibit unique

growth rate patterns, none of the 15 species overlap (except for *O. krugii* and *H. racemosum* for tree basal area size less than 800  $\text{cm}^2$ ). For five of the species, namely *H. racemosum*, *L. dominicensis*, *G. guidonia*, *S. berteriana*, and *Manilkara bidentata*, the growth rates from the available data had not peaked. The dashed line portions of the growth rate curves of these five species in Fig. 2 are an extensive extrapolation beyond the range of the data and represent the least squares best 'guess' of the growth rate trend to the asymptote. The fitted equations for these five species should be used with caution.

The species *Buchenavia capitata* had the best growth from actual data, peaking at 87  $\text{cm}^2 \text{ year}^{-1}$ . *Buchenavia capitata* is a large, spreading timber tree that occurs throughout the Caribbean islands, Panama and South America from Venezuela to French Guiana, Brazil, and Bolivia. The wood is reported to be of good quality, moderately heavy (specific gravity 0.61), and strong (Little and Wadsworth, 1964). The tree is also used for shade. Based on this growth analysis and its ecological and commercial impor-




Fig. 4. Illustration of growth rate response surfaces by topographic position. Species is *Cecropia peltata* in a moist tropical forest in Puerto Rico. Maximum observed  $ba$  values were  $1700 \text{ cm}^2$ ,  $3200 \text{ cm}^2$ , and  $2800 \text{ cm}^2$  for ridge, slope, and bottom topography, respectively. Observed ground incline ranged from  $12$  to  $42^\circ$ ,  $16$ – $64^\circ$ , and  $25$ – $51^\circ$  for ridge, slope, and bottom topography, respectively.

tance, this species has much to recommend it.

The species *Manilkara bidentata* shows the highest growth potential. Actual data exhibit growth rates in excess of  $60 \text{ cm}^2 \text{ year}^{-1}$  and the fitted function extrapolates growth to a peak of  $122 \text{ cm}^2 \text{ year}^{-1}$ . Additional research is needed, with data from older and larger trees, to see if this growth rate can actually be obtained. However, these initial results are very encouraging. Little and Wadsworth (1964) report this species as slow growing, which seems contrary to the results of this study. *Manilkara bidentata* occurs throughout the Caribbean, Panama, and northern South America from the Guianas and Venezuela to Peru and Northern Brazil. The wood is very hard, heavy (specific gravity 0.82), and extremely strong. The milky sap of this species is the source of balata gum, and the species is also used for shade.

As mentioned, four species showed significant trends with crown class. *Tetragastris balsamifera* is plotted in Fig. 3 to demonstrate the response curve by crown class. The family of curves show the substantial effect on growth going from one canopy position to another. The response surfaces of *C. peltata* are shown in Fig. 4 to depict the wide range of responses possible for those species sensitive to topographic position and the amount of ground incline. In ridge and bottom topography increasing ground incline greatly improves PAI. However, other species have different response surface shapes molded by the occurrence of positive and negative  $b_3$  through  $b_5$  coefficients. For example, *H. racemosum* has a negative  $b_3$  coefficient so increasing degree of ground incline in ridge topography would result in decreasing PAI. Each of the seven species under Eq. (3) has a unique set of response surfaces.

## 6. Conclusions

From this study of the growth rate trends of 15 tropical species within a mixed/moist forest, one can see the effects on growth from the complex mix of physiography and stand dynamics. I will leave it to tropical ecologists and foresters to investigate why some species are sensitive to

ground inclination and topography and others more to canopy position. Ultimately crown diameter may be the best indicator of growth, according to Wadsworth et al. (1989), and this should be investigated more fully. The Chapman-Richards model provides a flexible framework for investigating species growth, capable of incorporating many independent variables.

### Acknowledgments

The author gratefully acknowledges the International Institute of Tropical Forestry in Rio Piedras, Puerto Rico for providing data used in this study. The author would also like to thank the reviewers for their helpful comments and suggestions. This research was supported by USDA Forest Service forest management research funds.

### References

- Gerrard, D.J., 1968. Competition quotient: an index of the competitive stress affecting individual forest trees. *Dissertation Abstr. B*, 29: 426-427.
- Holdridge, L.R., 1967. *Life Zone Ecology*. Tropical Science Center, San Jose, Costa Rica, 206 pp.
- Little, Jr., E.L. and Wadsworth, F.H., 1964. Common trees of Puerto Rico and the Virgin Islands. *Agriculture Handbook No. 249*. USDA Forest Service, Washington, DC, 548 pp.
- Lowe, R.G., 1966. The effect of competition on tree growth (preliminary results). In: *Papers of the 2nd Nigerian Forestry Conference*, Enugu, Nigeria, pp. 115-149.
- Lowe, R.G., 1973. **Silvicultural** characteristics of trees in growth by pattern analysis and standcurve analysis on the electronic computer. *Res. Pap. 13*, Federal Department of Forest Research, Lagos, Nigeria, 14 pp.
- Mervart, J., 1969. Growth studies in the natural tropical forest for forest management purposes. *Obeche*, 1: 48-59.
- Mervart, J., 1972. Growth and mortality rates in the natural high forest of Western Nigeria. *Nig. For. Inf. Bull.* 22, 46 PP.
- Pienaar, L.V. and Turnbull, K.J., 1973. The Chapman-Richards generalization of von Bertalanffy's growth model for basal area growth and yield in even-aged stands. *For. Sci.*, 19: 2-22.
- Shifley, S.R. and Brand, G.J., 1984. Chapman-Richards growth function constrained for maximum tree size. *For. Sci.*, 30: 1066-1070.
- Smith, D.M., 1962. *The Practice of Silviculture*, 7th edn. John Wiley, New York.
- Wadsworth, F.H., 1953. New observations of tree growth in tabonuco forest. *Caribb. For.*, 14: 106-111.
- Wadsworth, F.H., Parresol, B.R. and Colon, J.C.F., 1989. Tree increment indicators in a subtropical wet forest. In: W.R. Wan Mohd, H.T. Chan and S. Appanah (Editors), **Proc. Seminar on Growth and Yield in Tropical Mixed/Moist Forests**. Forest Research Institute Malaysia, Kuala Lumpur, Malaysia, pp. 205-213.



# Forest Ecology and Management

---

## Submission of manuscripts.

Manuscripts should be submitted in triplicate. Authors from the Americas, Australia, New Zealand and the Pacific are requested to send their manuscripts to Forest Ecology and **Management**, Dr. Richard F. Fisher, Department of Forest Science, Texas A&M University, College Station, TX77843-2135, USA; all other authors are requested to send their manuscripts to the Editorial Office, Forest Ecology and Management, P.O. Box 181, 1000 AD Amsterdam, The Netherlands.

**Authors in Japan please note:** Upon request, Elsevier Science Japan will provide authors with a list of people who can check and improve the English of their paper (*before submission*). Please contact our Tokyo office: Elsevier Science Japan, 20-12 Yushima 3-chome, Bunkyo-ku, Tokyo 113; tel. (03)-3833-3821; fax (03)-3836-3064.

**Electronic manuscripts:** Electronic manuscripts have the advantage that there is no need for the rekeying of text, thereby avoiding the possibility of introducing errors and resulting in reliable and fast delivery of proofs. For the initial submission of manuscripts for consideration, hardcopies are sufficient. For the processing of *accepted papers*, electronic versions are preferred. After *final acceptance*, your disk plus two, final and exactly matching printed versions should be submitted together. Double density (DD) or high density (HD) diskettes (3.5 or 5.25 inch) are acceptable. It is important that the file saved is in the native format of the wordprocessor program used. Label the disk with the name of the computer and word-processing package used, your name, and the name of the file on the disk. Further information may be obtained from the Publisher.

All questions arising after acceptance of the manuscript, especially those relating to proofs, should be directed to Elsevier Editorial Services, Mayfield House, 256 Banbury Road, Oxford OX2 7DH, UK, tel. (+44)(0)1865 314900, fax (+44)(0)1865 314990.

**Advertising information:** Advertising orders and enquiries may be sent to: Elsevier Science B.V., Advertising Department, P.O. Box 211, 1000 AE Amsterdam, The Netherlands, tel. (+31-20) 4853795, fax (+31-20) 4853810. Courier shipments to street address: Molenwerf 1, 1014 AG Amsterdam, The Netherlands. In the UK: TG Scott & Son Ltd., attn. Vanessa Bird, Portland House, 21 Narborough Rd., Cosby, Leicestershire, LE9 5TA, UK, tel. (+44)(0)116 2750-521/2753-333, fax (+44)(0)116 2750-522. In the USA and Canada: Weston Media Associates, attn. Daniel Lipner, P.O. Box 1110, Greens Farms, CT 06436-1110, USA, tel. (203) 261-2500, fax (203) 261-0101.

**US mailing notice-forest Ecology and Management (0378-I 127)** is published monthly by Elsevier Science B.V. (Molenwerf 1, Postbus 211, 1000 AE Amsterdam). Annual subscription price in the USA is US\$ 1492.00 (valid in North, Central and South America), including air speed delivery. Second class postage rate is paid at Jamaica, NY 11431.

USA POSTMASTERS: Send address changes to *Forest Ecology and Management* Publications Expediting, Inc., 200 Meacham Avenue, Elmont, NY 11003. Airfreight and mailing in the USA by Publications Expediting.

**Forest Ecology and Management has no page charges**

For a full and complete Guide for Authors, please refer to *Forest Ecology and Management*, Vol. 72, No. 1, pp. 93-96

© 1995, Elsevier Science B.V. All rights reserved.

0378-I 127/95/\$09.50

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying or otherwise, without the prior written permission of the publisher, Elsevier Science B.V., Copyright and Permissions Department, P.O. Box 521, 1000 AM Amsterdam, The Netherlands.

No responsibility is assumed by the Publisher for any injury and/or damage to persons or property as a matter of products liability, negligence or otherwise, or from any use or operation of any methods, products, instructions or ideas contained in the material herein.

Although all advertising material is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement of the quality or value of such product or of the claims made of it by its manufacturer.

@The paper used in this publication meets the requirements of ANSI/NISO Z39.48-1992 (Permanence of Paper).

Printed in The Netherlands

# LANDSCAPE AND URBAN PLANNING

An International Journal of Landscape Ecology,  
Landscape Planning and Landscape design

## Editor-in-Chief:

J.E. Rodiek, *College of Architecture, Texas A & M University, College Station, TX 778433137, USA*

## AIMS AND SCOPE

A journal concerned with conceptual, scientific and design approaches to land use. By emphasizing ecological understanding and a multi-disciplinary approach to analysis and planning and design, it attempts to draw attention to the interrelated nature of problems posed by nature and human use of land. In addition, papers dealing with ecological processes and interactions within urban areas, and between these areas and the surrounding natural systems which support them, will be considered. Papers in which specific problems are examined are welcome. Topics might include but are not limited to landscape ecology, landscape planning and landscape design. Landscape ecology examines how heterogeneous combinations of ecosystems are structured, how they function and how they change. Landscape planning examines the various ways humans structure their land use changes. Landscape design involves the physical strategies and forms by which land use change is actually directed. *Landscape and Urban Planning* is based on

the premise that research linked to practice will ultimately improve the human made landscape.

## Editorial Advisory Board:

I.D. Bishop, *Parkville, Vic., Australia*,  
E.G. Bolen, *Wilmington, NC, USA*,  
I.D. Bruns, *Schorndorf, Germany*,  
J.B. Byrom, *Edinburgh, UK*,  
T.C. Daniel, *Tucson, AZ, USA*,  
R.M. DeGraaf, *Amherst, MA, USA*,  
J.G. Fabos, *Amherst, MA, USA*,  
S. Gonzalez Alonso, *Madrid, Spain*,  
M. Hough, *Etobicoke, ON, Canada*,  
P. Jacobs, *Montreal, PQ, Canada*,  
D.S. Jones, *Melbourne, Vic., Australia*,  
H. Lavety, *Milton, Qld., Australia*,  
W.M. Marsh, *Flint, MI, USA*,  
D.L. Mitchell, *Dallas, TX, USA*,  
D.G. Morrison, *Athens, GA, USA*,  
J.I. Nassauer, *St. Paul, MN, USA*,  
M. Nelischer, *Guelph, ON, Canada*,  
D.D. Paterson, *Vancouver, BC, Canada*,  
A. Ramos, *Madrid, Spain*,  
P. Shepard, *Claremont, CA, USA*,  
O.R. Skage, *Alnarp, Sweden*,  
R.C. Smardon, *Syracuse, NY, USA*,  
G. Sorte, *Alnarp, Sweden*,  
F. Stearns, *Rhineland, WI, USA*.

R.C. Szaro, *Washington, DC, USA*,  
J.W. Thomas, *La Grande, OR, USA*,  
P.J. Trowbridge, *Ithaca, NY, USA*,  
T.H.D. Turner, *London, UK*,  
M.J. Vroom, *Wageningen, The Netherlands*,  
W.V. Wendler, *College Station, TX, USA*,  
B.-E. Yang, *Seoul, Korea*,  
E.H. Zube, *Tucson, AZ, USA*

ABSTRACTED/INDEXED IN  
Applied Ecology Abstracts,  
Biological Abstracts,  
Current Contents B & S,  
Environmental Periodicals  
Bibliography, Geobase,  
Geographical Abstracts,  
LandSearch.

## 1994 SUBSCRIPTION DATA

Volumes **28-30** (in 9 issues)  
Subscription price:  
Dfl. 1080.00 (US \$564.00)  
incl. Postage  
**ISSN 0169-2046**



**ELSEVIER**  
SCIENCE PUBLISHERS

Elsevier Science Publishers,  
P.O. Box 211, 1000 AE Amsterdam,  
The Netherlands  
Fax: (020) 5803-203

Customers in the USA and Canada:  
Elsevier Science Publishers  
P.O. Box 945, Madison Square Station,  
New York, NY 101 W-0757, USA  
Fax: (212) 633-3680

The Dutch Guilder price quoted applies worldwide, except in the Americas (North, Central and South America). The US Dollar price quoted applies in the Americas only. Non VAT registered customers in the European Community should add the appropriate VAT rate applicable in their country to the price.