

# Response of beetles (Coleoptera) at three heights to the experimental removal of an invasive shrub, Chinese privet (*Ligustrum sinense*), from floodplain forests

Michael D. Ulyshen · Scott Horn ·  
James L. Hanula

Received: 29 June 2009 / Accepted: 19 August 2009 / Published online: 3 September 2009  
© U.S. Government 2009

**Abstract** Chinese privet (*Ligustrum sinense* Lour.), an invasive shrub from Asia, is well established in the southeastern United States where it dominates many floodplain forests. We used flight intercept traps to sample beetles at three heights (0.5, 5 and 15 m) in ~2 ha plots in which *L. sinense* had (by chainsaws or mulching machine) or had not been removed from forests in northeastern Georgia. Removing *L. sinense*, particularly by machine, increased the richness and diversity of beetles and affected composition near the ground (0.5 m) but not in the forest canopy (15 m). There were no differences among treatments above the *L. sinense* canopy (5 m) aside from *Xylosandrus crassiusculus*, an exotic ambrosia beetle from Asia, dominating the beetle community at that height in control plots. Removing the *L. sinense* layer greatly increased beetle richness near the ground, resulting in vertical distribution patterns more similar to those observed in areas of forest devoid of *L. sinense*. We suspect that even organisms in the canopy will benefit from privet removal in the long term given that tree regeneration is nearly impossible in forests in which *L. sinense* has become well established.

**Keywords** Arthropods · Biodiversity · Bottomland hardwood forests · Insects ·

Non-native · Restoration · Temperate deciduous forests · Vertical distribution · Vertical stratification · Wetlands

## Introduction

Since its introduction in the mid nineteenth century, Chinese privet (*Ligustrum sinense* Lour.), an evergreen shrub native to Asia, has become one of the most abundant non-native invasive plants in the southeastern United States. The species currently occupies about 3.5% of the total land area covered by forests in the region (Rudis et al. 2006) and dominates many riparian areas. For example, 59% of the Upper Oconee River floodplain in north Georgia was covered by *L. sinense* by 1999 (Ward 2002). The expansion of *L. sinense* into new areas can be largely attributed to seed dispersal by animals (Miller 2003) and floodwaters (Ward 2002) and rapid establishment is aided by high growth rates, vegetative reproduction, shade tolerance, and prolific seed production (Langeland and Burkes 1998). Once established, *L. sinense* grows in dense thickets up to 9 m in height (Miller 2003). The species competes strongly with native plants for light and nutrients (Fig. 1), displacing many species from the understory and suppressing tree regeneration (Miller 1997; Brown and Pezeshki 2000; Kittell 2001; Morris et al. 2002; Merriam and Feil 2002; Wilcox and Beck 2007;

M. D. Ulyshen (✉) · S. Horn · J. L. Hanula  
USDA Forest Service, Southern Research Station,  
320 Green Street, Athens, GA 30602, USA  
e-mail: mulyshen@hotmail.com



**Fig. 1** Forest floor beneath Chinese privet in Clarke County, Georgia. Almost all plants visible here are Chinese privet

Hanula et al. 2009). Although *L. sinense* may provide suitable forage for some generalist vertebrate herbivores (Stromayer et al. 1998; Kittell 2001), it is doubtful that many species have benefited from its introduction. Forest floors lacking native plants will likely lack other species as well, including many insects. Insects active above the level of *L. sinense* may also be affected if *L. sinense* disrupts their seasonal (e.g., Schaefer 1991) or daily (e.g., Costa and Crossley 1991) movements between the forest canopy and ground. Previous studies have shown that invasive plants can affect the abundance, diversity and composition of native insect communities (e.g., Greenwood et al. 2004; de Groot et al. 2007; Wiezik et al. 2007; Remsburg et al. 2008). However, to our knowledge, nothing is known about how insect faunas are affected above the reach of invasive plants, for example, in the forest canopy. The objective of this study was to determine how removing *L. sinense* would affect beetle diversity and composition at three heights in southeastern U.S. floodplain forests.

## Methods

### Study area


Study sites were located within the Oconee River watershed in northeast Georgia (Fig. 2). The four sites (i.e., blocks) selected were Sandy Creek Nature Center, Clarke Co.; Georgia State Botanical Gardens, Clarke Co.; Scull Shoals Experimental Forest, Oconee National Forest, Oglethorpe Co.; and Watson Springs Forest, University of Georgia, Greene Co.

### Experimental design

Three ~2 ha plots were established in floodplain forests dominated by *L. sinense* at each of the four sites. Each plot at each site received one of the following treatments in October/November 2005: (1) *Mulch*, mechanical removal of all *L. sinense* using a machine (Klepac et al. 2007), leaving a layer of mulch on the soil surface; (2) *Chainsaw*, hand removal of all *L. sinense* <1.27 cm using gas-powered saws and machetes, leaving piles of cut *L. sinense* <1 m high; and (3) *Control*, no removal of *L. sinense*. Treatments were not randomly assigned to plots for logistical reasons; the most accessible plot at each location received the mulch treatment. Following removal, all *L. sinense* stumps were treated with either 30% triclopyr (Garlon®4) or 30% glyphosate (Foresters'®).

### Sampling

In total, 72 flight intercept traps identical to those used by Ulyshen and Hanula (2007) were used to sample beetles at three heights (0.5, 5 and 15 m) above the forest floor. Because *L. sinense* reached 4–5 m tall on average, the traps at 5 m in control plots were usually situated just above the *L. sinense* canopy. However, *L. sinense* canopies are not even surfaces with some individuals exceeding 10 m in height (personal obs.). The traps at 15 m were generally above the lowest leaf bearing branches of the tallest trees and were therefore considered to be within the lower forest canopy. The lower (0.5 m) traps were suspended from metal poles (Ulyshen and Hanula 2007). The middle and upper traps were suspended above the lower traps from ropes pulled over limbs in the canopy (Ulyshen and Hanula 2007). These traps were attached to rectangular hangers made from 1.6 cm diam PVC tubing and rope. To construct each hanger, two 46 cm lengths of PVC tubing were attached from their ends by rope to create a rectangular frame 1 m long. Each trap was suspended from the middle of the upper section of PVC tubing. Sections of rope cut to length were attached to the upper and lower sections of tubing to complete the assembly. Trap assemblies were installed at two locations near the center of each plot. When in use, a 1% formaldehyde and saturated salt solution with a small amount of dish soap was added to the traps to kill and preserve the catch. We



**Fig. 2** Research plots in northeastern Georgia

sampled on the following dates in 2006: 7–14 March, 4–11 April, 2–9 May, 2–9 June, 13–20 July and 17–24 August. The two traps at each height in each plot were combined before analysis.

#### Statistical analysis

The complete dataset, consisting of 550 morphospecies and 10,412 individuals, was used to perform rarefaction using PAST (Hammer et al. 2001) to compare the number of beetle morphospecies among treatments and trap heights. After excluding several sampling periods (March, July and August) and locations (botanical garden control 5 m and sandy creek control 0.5 m) due to missing samples, the dataset used for all other analyses (but further

modified for PC-ORD and ANOSIM, see below) consisted of 427 morphospecies and 7,237 individuals. Separate ANOVAs for fixed effects were performed on morphospecies richness, Shannon's diversity and evenness (i.e., the response variables) using SAS. The blocking variable, location, was removed from the model after preliminary analyses found it to be insignificant for all response variables. Nonmetric multidimensional scaling (NMS) was performed on a dataset consisting of 191 morphospecies captured in  $\geq 3$  samples using PC-ORD (McCune and Mefford 2006). The same modified data set was used in PAST (Hammer et al. 2001) to perform ANOSIM with 10,000 permutations using a Bray–Curtis distance measure to quantitatively compare beetle community similarity among treatments at each trap height.


## Results

### Rarefaction

For a given number of individuals, there were more morphospecies at 15 and 5 m than at 0.5 m above the ground and there were fewer morphospecies in control plots than mulch and chainsaw plots (Fig. 3).

### Richness

Richness varied significantly among treatments ( $F_{2,25} = 8.78$ ,  $P < 0.01$ ) and there was a significant interaction between treatment and trap height ( $F_{4,25} = 9.60$ ,  $P < 0.0001$ ). Richness was similar among treatments for traps suspended 5 and 15 m, but varied greatly 0.5 m above the ground (Fig. 4). On average, there were nearly twice as many morphospecies near the ground in mulch plots than in control plots and



**Fig. 3** Rarefaction analyses ( $\pm$ SD) for beetles collected at three heights (0.5, 5 and 15 m) above the ground and in plots in which *L. sinense* was removed by chainsaw, by machine (i.e., “mulch”), or was not removed (i.e., “control”) in northeastern Georgia


about 11 more morphospecies near the ground in mulch plots than in chainsaw plots (Fig. 4).

### Evenness

Evenness varied significantly among both treatments ( $F_{2,25} = 3.39$ ,  $P = 0.05$ ) and trap heights ( $F_{2,25} = 6.19$ ,  $P < 0.01$ ) and there was a significant interaction between treatment and trap height ( $F_{4,25} = 3.93$ ,  $P < 0.05$ ). Evenness was similar among treatments for traps suspended 0.5 and 15 m, but varied greatly 5 m above the ground (Fig. 5). Evenness was considerably lower in control plots at 5 m because  $>51\%$  of the beetles collected there belonged to a single species, *Xylosandrus crassiusculus*. When *X. crassiusculus* was removed from the dataset there were no differences in evenness among treatments or trap heights (Fig. 5).



### Diversity

For the complete dataset, diversity also varied significantly among both treatments ( $F_{2,25} = 6.44$ ,  $P \leq 0.01$ ) and trap heights ( $F_{2,25} = 7.06$ ,  $P < 0.01$ ) and there was a significant interaction between treatment and trap height ( $F_{4,25} = 2.90$ ,  $P < 0.05$ ). However, after removing *X. crassiusculus* from the dataset, diversity differed only among treatments ( $F_{2,25} = 3.66$ ,  $P \leq 0.05$ ), being lowest in control plots (Fig. 5).



**Fig. 4** Mean  $\pm$  SE ( $n = 3$  or 4) morphospecies richness of beetles collected at three heights above the ground in deciduous floodplain forests in which *L. sinense* had (i.e., Chainsaw, Mulch) or had not (i.e., Control) been removed in northeast Georgia

**Fig. 5** Mean  $\pm$  SE ( $n = 3$  or 4) evenness and diversity of beetles collected at three heights above the ground in deciduous floodplain forests in which *L. sinense* had (i.e., Chainsaw, Mulch) or had not (i.e., Control) been removed in northeast Georgia



**Fig. 6** Nonmetric multidimensional scaling plot with lines separating the three trap heights (0.5, 5 and 15 m)

### Composition

The ordination from NMS (final stress 16.1) shows that the beetle faunas at 0.5 and 15 m were distinctly different (i.e., the points representing the trap locations

at the two heights are widely separated) and that the fauna at 5 m was intermediate in composition (Fig. 6). Although treatments did not form obvious groupings at 0.5 m or at 15 m, control plots formed a distinct grouping at 5 m (Fig. 6). The results from ANOSIM support these conclusions: there were no significant differences in community similarity among treatments at 0.5 or 15 m but the fauna in control plots was significantly dissimilar from that in chainsaw or mulch plots at 5 m (Table 1). However, when *X. crassiusculus* was removed from the dataset, there were significant differences among treatments at 0.5 m but not at 5 or 15 m (Table 1).

### Discussion

Removing *L. sinense*, particularly by machine, increased the richness and diversity of beetles and affected composition near the ground (0.5 m) but not in the forest canopy (15 m). More species were captured at 0.5 m in mulch plots than in chainsaw plots, possibly because the piles of debris left in chainsaw plots reduced the numbers of insects reaching the traps. There were no differences among treatments above the *L. sinense* canopy (5 m) aside from *X. crassiusculus*, an exotic ambrosia beetle from

**Table 1** Results of ANOSIM (analysis of similarity) among treatments at each trap height (0.5, 5 and 15 m) for the entire dataset (top) and excluding *X. crassiusculus* (bottom)

	Chainsaw	Control	Mulch
<i>Entire data set</i>			
0.5 m: $R = 0.21$ , $P = 0.07$			
Chainsaw	–	0.18	0.34
Control	–	–	0.06
5 m: $R = 0.47$ , $P = 0.01$			
Chainsaw	–	0.03	0.80
Control	–	–	0.03
15 m: $R = -0.14$ , $P = 0.87$			
Chainsaw	–	0.68	1
Control	–	–	0.72
<i>Excluding X. crassiusculus</i>			
0.5 m: $R = 0.53$ , $P = < 0.001$			
Chainsaw	–	0.03	0.26
Control	–	–	0.03
5 m: $R = -0.02$ , $P = 0.57$			
Chainsaw	–	0.3	0.77
Control	–	–	0.44
15 m: $R = -0.14$ , $P = 0.87$			
Chainsaw	–	0.72	1
Control	–	–	0.72

$P$ -values for pairwise comparisons are also given in matrices for each height. Large positive values of  $R$  (up to 1) and low values of  $P$  (e.g.,  $< 0.05$ ) signify dissimilarity among or between beetle communities

Asia, dominating the beetle community there in control plots. *X. crassiusculus* is not known to attack *L. sinense*, however it seems possible given the wide host range of the species and the fact that both species are originally from Asia.

Previous studies on the vertical distribution patterns of beetles in temperate deciduous forests have shown either no differences between the ground and canopy, or higher richness or diversity near the ground (Nielsen 1987; Preisser et al. 1998; Ulyshen and Hanula 2007). For example, Ulyshen and Hanula (2007) sampled beetles at 0.5 and  $\geq 15$  m in the same Scull Shoals forest used in this project, only slightly higher on the floodplain where *L. sinense* grew sparsely or not at all. In that study, richness was similar between the two heights and diversity was significantly higher near the ground. In this study, there were significantly fewer beetle species near the ground in control plots (i.e., beneath a canopy of

*L. sinense*) than at 5 or 15 m. Removing the *L. sinense* layer greatly increased beetle richness near the ground, resulting in vertical distribution patterns more similar to those observed in areas of forest devoid of *L. sinense*.

The results of this study speak only to the effects of *L. sinense* removal on beetles within the first year. Understanding the long-term effects will require further research. Because tree regeneration is nearly impossible beneath *L. sinense*, even organisms in the canopy will be at risk as canopies become increasingly sparse with each dying tree.

**Acknowledgments** We thank Stephanie Cahill for constructing the traps, Mike Cody for assisting with field work and Jared Swain for sorting samples. We also thank two anonymous reviewers for suggestions that greatly improved the manuscript. This research was supported by the USDA Forest Service Special Development Technology Program (Project Number R8-2005-01).

## References

- Brown CE, Pezeshki SR (2000) A study on waterlogging as a potential tool to control *Ligustrum sinense* populations in western Tennessee. *Wetlands* 20:429–437
- Costa JT III, Crossley DA Jr (1991) Diel patterns of canopy arthropods associated with three tree species. *Environ Entomol* 20:1542–1548
- De Groot M, Kleijn D, Jogan N (2007) Species groups occupying different trophic levels respond differently to the invasion of semi-natural vegetation by *Solidago canadensis*. *Biol Conserv* 136:612–617
- Greenwood H, O'Dowd DJ, Lake PS (2004) Willow (*Salix x rubens*) invasion of the riparian zone in south-eastern Australia: reduced abundance and altered composition of terrestrial arthropods. *Divers Distrib* 10:485–492
- Hammer Ø, Harper DAT, Ryan PD (2001) PAST: paleontological statistics software package for education and data analysis. *Palaeontologia Electronica* 4(1):9
- Hanula JL, Horn S, Taylor JW (2009) Chinese privet (*Ligustrum sinense*) removal and its effect on native plant communities of riparian forests. *Invasive Plant Science and Management* (in press)
- Kittell MM (2001) Relationships among invasive Chinese privet, plant diversity, and small mammal captures in southeastern deciduous forests. MS Thesis, Clemson University, 35 pp
- Klepac J, Rummer RB, Hanula JL, Horn S (2007) Mechanical Removal of Chinese Privet. USDA Forest Service, Southern Research Station, Research Paper SRS-43
- Langeland KA, Burkes KC (1998) Identification and biology of non-native plants in Florida's natural areas. University of Florida, Gainesville, 165 pp
- McCune B, Mefford MJ (2006) PC-ORD. multivariate analysis of ecological data. Version 5. MJM Software. Gleneden Beach, Oregon

- Merriam RW, Feil E (2002) The potential impact of an introduced shrub on native plant diversity and forest regeneration. *Biol Invasions* 4:369–373
- Miller JH (1997) Exotic invasive plants in southeastern forests. In: Britton KO (ed) *Proceedings of exotic pests of eastern forests*. Tennessee Exotic Pest Plant Council, Nashville, TN, pp 97–106
- Miller J (2003) *Nonnative Invasive Plants of Southern Forests*. USDA Forest Service, General Technical Report, SRS-62. Asheville, N.C.: USDA, Forest Service, Southern Research Station, p 93
- Morris LL, Walck JL, Hidayati SN (2002) Growth and reproduction of the invasive *Ligustrum sinense* and native *Forestiera ligustrina* (Oleaceae): implications for the invasion and persistence of a nonnative shrub. *Int J Plant Sci* 163: 1001–1010
- Nielsen BO (1987) Vertical distribution of insect populations in the free air space of beech woodland. *Entomol Meddelelser* 54:169–178
- Preisser E, Smith DC, Lowman MD (1998) Canopy and ground level insect distribution in a temperate forest. *Selbyana* 19: 141–146
- Rensburg AJ, Olson AC, Samways MJ (2008) Shade alone reduces adult dragonfly (Odonata: Libellulidae) abundance. *J Insect Behav* 21:460–468
- Rudis VA, Gray A, McWilliams W, O'Brien R, Olson C, Oswalt S, Schulz B (2006) Regional monitoring of nonnative plant invasions with the Forest Inventory and Analysis program. In: McRoberts RE, Reams GA, Van Duesen PC, McWilliams WH (eds) *Proceedings of the sixth annual FIA Symposium 2004 Sept 21–24*, pp 49–64. Denver, CO. Gen. Tech. Rep. WO-70 U.S. Department of Agriculture, Forest Service, Washington, DC
- Schaefer M (1991) The animal community: diversity and resources. In: Röhrig E, Ulrich B (eds) *Temperate deciduous forests. Ecosystems of the world*. Elsevier, New York, pp 51–120
- Stromayer KAK, Warren RJ, Harrington TB (1998) Managing Chinese privet for white-tailed deer. *South J Appl For* 22:227–230
- Ulyshen MD, Hanula JL (2007) A comparison of the beetle (Coleoptera) fauna captured at two heights above the ground in a North American temperate deciduous forest. *Am Midl Nat* 158:260–278
- Ward RW (2002) Extent and dispersal rates of Chinese privet (*Ligustrum sinense*) invasion on the upper Oconee River floodplain, North Georgia. *Southeastern Geo* 1:29–48
- Wiezik M, Svitok M, Dovčiak M (2007) Conifer introductions decrease richness and alter composition of litter-dwelling beetles (Coleoptera) in Carpathian oak forests. *Forest Ecol Manag* 247:61–71
- Wilcox J, Beck CW (2007) Effects of *Ligustrum sinense* Lour. (Chinese Privet) on abundance and diversity of songbirds and native plants in a southeastern nature preserve. *Southeast Nat* 6:535–550