

Growth predictions for tree species planted on marginal soybean lands in the Lower Mississippi Valley

J.W. Groninger, W.M. Aust, M. Miwa, and J.A. Stanturf

ABSTRACT: The establishment of bottomland hardwood forest stands and riparian buffers on frequently-flooded soybean (*Glycine max.*) lands in the Lower Mississippi Valley represents a tremendous opportunity to provide both economic and environmental benefits to the region. Selecting appropriate sites for reestablishing tree cover, accurately predicting the productivity of planted trees and optimally matching species to site are critical for the economic justification and implementation of tree planting in conservation programs.

This study tests a low-cost methodology that incorporates the expert system developed by Baker and Broadfoot (1979) to predict tree growth rates calibrated with soils data from recently published Natural Resources Conservation Service (NRCS) soil surveys specific for combinations of tree species and soil series. This information is used to make site index projections for economically-marginal soybean lands. Site index estimates ranged from 28.0 m (92 ft) for cottonwood (base age 30) on Mhoun soils to 18.0 m (61 ft) for sycamore (base age 50) on Forestdale soils. Use of this method results in tree growth predictions that are both more mechanistically-based and often more conservative than the site index projections published in soil surveys, particularly for species intolerant of flooding during the growing season.

Key words: Bottomland hardwoods, CRP, *Glycine max.*, Lower Mississippi Valley, marginal soybean lands, site index, soil survey, tree growth, tree planting, tree species-site relations.

The Lower Mississippi Valley contains bottomland hardwood-dominated forests that are among the most productive in the United States. Widespread conversion of forest land to agricultural uses occurred throughout the region over the past 200 years. Presently, forests cover less than one-quarter of the area of forestland prior to European settlement. Most recently, widescale loss of forest cover occurred during the early 1970s in response to increases in soybean prices (Sternitzke 1976). In many cases, these last forests to be cleared occupied sites which were too frequently flooded to support row-crop agriculture on a sustainable basis due to poor internal drainage, backwater flooding, or both. Changes in both soybean and forest products markets have rendered some of these lands economically marginal for soybean production. Changing economics, in combination with an

increasing recognition of the environmental benefits of forested wetlands, have kindled widespread interest in converting marginal soybean land back to bottomland hardwood forests (Amacher et al. 1998). Accordingly, many of these lands are being enrolled in the Conservation Reserve Program (CRP), Wetlands Reserve Program and other cost-share programs involving forest establishment.

CRP landowners who elect to plant trees do so for a number of reasons, including enhancement of recreation, aesthetics, and wildlife values, as well as the eventual sale of forest products (Olmstead and McCurdy 1989). These benefits are realized soonest, and to the greatest extent, when the conversion from open cropland to closed forest canopy occurs rapidly (Hubbard and Lowrance 1997; Tweed and Portwood 1997). On- and off-site environmental benefits, such as nitrate and pesticide removal, streambank stabilization, and water temperature moderation also are accelerated through rapid reforestation. Maximizing tree and stand growth rates without increasing management costs generates stronger financial incentives to plant trees and may motivate landowners to keep cropland or riparian buffers in forest cover beyond the expiration date of current cost

share programs. Realization of these objectives requires an understanding of the physical, chemical, and hydrological characteristics of the soil and site (Hodges 1997), and knowing the tree species most appropriate for those conditions.

Soil series-specific tree planting recommendations and growth rate predictions are given in county and parish soil surveys produced by the U.S. Department of Agriculture's Natural Resources Conservation Service (NRCS). This information is generally based on research conducted on cutover forest lands and does not reflect soil and site changes brought about by 20 or more years of soybean culture, including increased soil bulk density, lower organic matter content, reduced rooting volume, and altered nutrient availability (Francis 1984). Unfortunately, species-to-plant recommendations and predictions of potential tree growth rate based on data from former soybean lands are not yet available. It will be some time before the trees which were only recently planted on these lands are old enough to provide empirical growth and productivity estimates that overcome some of the confounding factors associated with stand establishment. A mechanistically-sound basis for making tree species-to-plant recommendations and growth estimates would aid tree planting efforts in the interim.

Baker and Broadfoot (1978, 1979) developed a species-specific method for predicting tree height attained at a given base age (site index) for several bottomland hardwood species by evaluating soil and site attributes important to tree growth. Previous research with cherrybark oak (*Quercus pagoda* Raf.) demonstrated the accuracy of this method (Aust and Hodges 1988; Belli et al. 1998). In the past, use of this system required intensive sampling of the proposed planting site. However, recently published soil surveys now include the necessary information on the physical and chemical properties of soil series. With these data, along with assumptions regarding soil and site characteristics of former soybean lands, height growth can be predicted for several tree species planted on any soil series within the southern hardwood region (Broadfoot 1976).

The objective of this study is to evaluate the use of Baker and Broadfoot's (1979) field-based method for marginal soybean lands using published soils data and generic assumptions regarding soil and site conditions. Comparisons are then made with species-to-plant and site index estimates published in soil surveys.

John W. Groninger is assistant professor in the Department of Forestry, Southern Illinois University, Carbondale, Ill.; W. Michael Aust is associate professor, and Masuo Miwa is research scientist in the Department of Forestry, Virginia Polytechnic Institute and State University, Blacksburg, Va.; and John A. Stanturf is project leader at the U.S. Department of Agriculture Forest Service Southern Research Station, Stoneville, Miss.

Table 1. Baker and Broadfoot (1979) site index evaluation for cottonwood using assumptions of long-term soybean cultivation and information from a published soil survey. Described here is an Amagon soil with data from the USDA/NRCS Soil Survey Division Official Soil Series Descriptions (www.statlab.iastate.edu/soils/osd/). The sum of underlined point values equals predicted site index in feet at a base age of 30 years.

Soil-site property	Soil-site condition and relative quality			Source of Information
	Best	Medium	Poor	
Factor 1. Physical condition				
Soil depth and presence of artificial or inherent pan	Deep soil (> 4 ft; without pan) <u>(16)²</u>	Medium depth (2-4 ft), or a soil with a plowpan <u>(11)</u>	Shallow soil (< 2 ft), or a soil with an inherent pan <u>(-11)</u>	Soil Survey, plowpan assumed to be disrupted by deep plowing
Texture (in rooting zone)	Medium-textured; silty or loamy <u>(11)</u>	Coarse-textured; sandy <u>(8)</u>	Fine-textured; clayey <u>(-4)</u>	Soil Survey
Compaction (in surface foot)	No compaction; loose, porous, friable, bulk density < 1.4 g/cc <u>(9)</u>	Moderately compacted; firm moderately tight, bulk density 1.4-1.7 g/cc <u>(6)</u>	Strongly compacted; tight, bulk density > 1.7 g/cc <u>(-3)</u>	Soil Survey
Structure (in rooting zone)	Granular; structureless, single-grained if sandy, massive if loamy or silty <u>(5)</u>	Prismatic blocky <u>(-3)</u>	Massive (if clayey); platy <u>(3)</u>	Soil Survey
Past use and present cover	Undisturbed; near-virgin forest cover <u>(5)</u>	Moderate cultivation; cultivated < 20 yr, or open with grass cover <u>(2)</u>	Intensive cultivation; cultivated > 20 yr, or open and bare <u>(-2)</u>	Assume intensive cultivation for > 20 yr
Factor 2. Moisture availability during growing season				
Water table depth	2 - 6' <u>(10)</u>	1-2'; 7-10' <u>(7)</u>	1' (unsuitable); 10' (0) ² <u>(0)</u>	Soil Survey
Artificial or inherent pans	No pans <u>(9)</u>	Plowpan <u>(6)</u>	Inherent pan <u>(-6)</u>	Soil Survey, plowpan assumed to be disrupted by deep plowing
Topographic position	Floodplain or stream bottom <u>(7)</u>	Stream terraces or lower slopes <u>(5)</u>	Upland <u>(-5)</u>	Soil Survey
Microsite	Concave; depression, pocket, trough <u>(7)</u>	Level; flat <u>(5)</u>	Convex; ridge, mound <u>(-2)</u>	Assumed to be flat due to leveling associated with cultivation
Structure (in rooting zone)	Granular; structureless, massive (if silty, loamy, or clayey); stratified <u>(5)</u>	Prismatic; blocky (if sandy); platy <u>(9)</u>	Structureless, single-grained <u>(-3)</u>	Soil Survey
Texture (in rooting zone)	Silty or loamy (or stratified) <u>(4)</u>	Clayey <u>(2)</u>	Sandy <u>(-2)</u>	Soil Survey
Flooding	Winter through Spring <u>(2)</u>	Winter only <u>(1)</u>	None (-1); continuous (unsuitable) <u>(-1)</u>	Soil Survey
Past use and present cover	Undisturbed; near-virgin forest cover <u>(2)</u>	Moderate cultivation cultivated < 10 yr or open with grass cover <u>(1)</u>	Intensive cultivation cultivated > 10 years or open and bare <u>(-1)</u>	Assume intensive cultivation > 10 years

Table 1 cont.

Soil-site property	Soil-site condition and relative quality			Source of Information
	Best	Medium	Poor	
Factor 3. Nutrient availability				
Geologic source	Mississippi River, Loess, Blackland (8)	Mixed Coastal Plain and other (5)	Coastal Plain (-5)	Soil Survey
Past use and present cover	Undisturbed; near-virgin forest cover, cultivated < 5 yr (5)	Moderate cultivation cultivated 5-10 yr, or open with grass (3)*	Intensive cultivation cultivated > 10 yr or open and bare (4 if fertilized annually) (-3)*	Assume annual fertilization associated with soybean production
Organic matter (A-horizon)	> 2% (4)	1 - 2% (3)	< 1% (-3)	Assume < 1% due to long-term cultivation
Depth of topsoil (A-horizon)	> 6" or no profile development (3)	3 - 6" (2)	< 3" (-2)	Soil Survey
Soil age	Young, no profile development (Entisols) (3)	Medium, moderate profile development (Inceptisols) (2)	Old, well-developed, profile leached (Alfisols, Ultisols) (-2)	Soil Survey
pH (in rooting zone)	5.5 - 7.5 (3)	4.5 - 5.5 or 7.6 - 8.5 (2)	< 4.5 or > 8.5 (-2)	Soil Survey
Factor 4. Aeration				
Soil structure (in rooting zone)	Granular, porous; structureless, single-grained if sandy, massive if loamy or silty (3)	Prismatic; blocky (2)	Massive (if clayey); platy (-2)	Soil Survey
Swampiness	Wet in winter only (3)	Wet January-July (2)	Waterlogged all year (Unsuitable)	Soil Survey
Mottling	None to 18" depth (3)	None to 8" depth (2)	Mottled to surface surface or gray mineral soil (-2)	Soil Survey
Soil color (in rooting zone)	Black, brown, red (3)	Yellow, brownish-gray (2)	Gray (-2)	Soil Survey

Methods

Selection of soils and species. The criteria for selection of the 10 Lower Mississippi Valley soil series were:

1. Soybean farming is commonly practiced on the series.
2. At least some phases of the series are classified as poorly or somewhat poorly drained.
3. Flooding frequency ranges from occasional to frequent.
4. The soil series is classified as hydric (Soil Conservation Service 1987).

All series used in this study met each of these criteria except Dundee, which is not hydric, but commonly occurs in association with several of the other selected soil series. Tree species were selected for evaluation on their adaptation to bottomland

sites and potential merchantability, and included the following: Eastern cottonwood (*Populus deltoides* Barr. ex Marsh.), green ash (*Fraxinus pennsylvanica* Marsh.), Nuttall oak (*Quercus nuttalli* Palmer), swamp chestnut oak (*Q. michauxii* Nutt.), water oak (*Q. nigra* L.), sweetgum (*Liquidambar styraciflua* L.), and sycamore (*Platanus occidentalis* L.).

Site index estimates. Soil series and species-specific site index estimates were made using the method developed by Baker and Broadfoot (1979). For each species, 23 soil and site characteristics were assigned one of three point values representing conditions thought to be good, fair, or poor for tree growth (Table 1). Species specific estimates are made by weighting point values to reflect the relative importance of each criterion to the growth of a particular species, as

well as that species maximum attainable site index. The sum of these weighted point values provides an estimate of site index (tree height at a given base age) for a tree species under the specified soil series and site conditions. Site index values use a base age of 50 years for all species presented here, except cottonwood where the conventional base age is 30 years.

In this study, site index estimates for seven southern bottomland hardwood tree species on each of 10 Lower Mississippi Valley soil series were made by calibrating the models of Baker and Broadfoot (1979) with soil chemical, physical, and hydrological data drawn from the general soil series descriptions at the NRCS web site (18 variables) (Table 1). Values for the remaining five variables were assigned on the basis of

Table 2. Site index predictions (m) based on Baker and Broadfoot (1979) for seven hardwood tree species for soil series likely to support marginal soybean agriculture in the Lower Mississippi Valley. Note that point values differ by species due to inherently differing resource needs, tolerances, and growth rates.

Soil series	Cottonwood ^a	Green ash	Nuttall oak	Swamp chestnut oak	Sweetgum	Water oak	Sycamore
Alligator	23.2	22.6	24.4	20.1	23.8	21.9	22.9
Amagon	27.1	23.2	25.3	22.3	25.0	23.5	25.0
Bowdre	22.3	22.6	23.5	19.8	23.8	21.6	21.6
Dundee	27.7	24.1	25.3	21.9	26.8	24.4	25.0
Forestdale	24.1	21.6	23.2	19.5	21.6	19.8	18.0
Mhoon	28.0	23.8	25.0	20.7	23.2	21.0	21.6
Newellton	24.4	23.5	24.7	20.1	24.7	21.9	23.2
Sharkey	22.9	21.9	23.8	20.1	22.9	21.3	21.3
Tensas	22.6	22.3	24.1	20.1	23.2	21.9	21.3
Tunica	22.9	21.6	23.5	18.6	22.3	19.5	18.6
Standard deviation	2.11	0.85	0.74	1.03	1.41	1.41	2.21

^a Base age for cottonwood is 30 years, 50 years for all other species.

Table 3. Percent difference between Baker and Broadfoot (1979) site index estimates for marginal soybean lands in the Lower Mississippi Valley and those included in NRCS Soil Survey species-to-plant recommendations for five bottomland hardwood tree species. P-values were obtained using a paired t-test comparing site index values from the two aforementioned predictors.

Soil series	Cottonwood	Green ash	Nuttall oak	Sweetgum	Water oak
Alligator	-16	+6	—	-3	-20
Amagon	-11	-5	-17	-18	-23
Bowdre	-34	—	—	-18	-25
Dundee	-9	—	—	-12	-16
Forestdale	-21	-9	-20	-25	-28
Mhoon	-16	-13	—	-24	—
Newellton	-20	+3	-5	-15	-20
Sharkey	—	—	—	-17	-22
Tensas	—	—	—	-24	-24
Tunica	—	—	—	-19	—
p value	0.001	0.01	0.04	0.0001	0.0001

assumptions made for economically-marginal soybean lands of the region (Table 1). Species-to-plant recommendations and projected site indices were collected from published USDA Soil Conservation Service/NRCS county and parish soil surveys within a 62-county/parish area of Arkansas, Louisiana, and Mississippi in the lower Mississippi Alluvial Plain (Groninger et al., In Press).

Results and discussion

Site index estimates ranged from 28.0 m (92 ft) for cottonwood (base age 30) on Mhoon soils to 18.0 m (61 ft) for sycamore (base age 50) on Forestdale soils (Table 2). Silt loam soils (Amagon and Dundee series) typically showed higher site indices than the other series, which were predominantly of clay texture. Among the tree species used in this study, green ash and Nuttall oak site index estimates varied least between soil types. The relatively rapid growth across all soils is consistent with the recognition among foresters that these species are well adapt-

ed to a wide range of frequently-flooded soil types. Sycamore and cottonwood varied most among soil series consistent with their sensitivity to soil and site conditions. Therefore, identification of suitable soil and site conditions is especially important when considering these species for reforestation.

Site index values calculated using Baker and Broadfoot (1979) were significantly lower than values given for species recommended for planting in soil surveys (Table 3) with differences typically in the 10 to 20% range. Green ash was an exception on Alligator and Newellton soils where estimates of site index were lower than published soil survey values, reflecting the recognized tolerance of green ash to poor soil conditions and highly-disturbed sites throughout the region. Cottonwood site index estimates varied less for soils of silt loam than for clay textures, suggesting that row crop agriculture may have a lesser impact on these soils. Maintenance of better internal drainage and consequently, a deeper root-

ing zone following disturbance is likely an important consideration in the predicted success of cottonwood on the silt loam soils.

Differences between soil survey values and our estimates based on Baker and Broadfoot (1979) likely reflect differences in assumptions made regarding soil and site conditions. Estimates included in soil surveys were derived from naturally-regenerated, well-stocked forest stands with no evidence of recent cutting or burning (Francis 1984). Predictions using the Baker and Broadfoot (1979) guide incorporated likely changes in soil and site conditions associated with long-term row crop cultivation. For instance, agricultural soils are characterized by higher bulk density, lower soil organic matter content, altered fertility, and sometimes smaller available rooting volume caused by restrictive plowpans (Francis 1985; Stanturf et al. 1998). However, the shrink-swell clay mineralogy and previous agricultural amendments may reduce the impact of these impediments. Site preparation and post-establishment management practices, including weed control treatments, fertilizer applications, fallowing, and deep plowing, have been recommended to ameliorate the adverse effects of long-term agronomic practices (Allen 1990; Baker and Blackmon 1978; Blackmon and White 1972; Francis 1985).

Perhaps the greatest value of this adaptation of Baker and Broadfoot (1979) for land managers is the incorporation of more specific tree growth estimates into the species selection process. Failure to correctly match species and site has consequences ranging from sub-optimal stand productivity to complete planting failure. While site index estimates ob-

rained through the use of the Baker and Broadfoot (1979) method appear to be an improvement over those published in soil surveys, they cannot be used in isolation to make recommendations for the appropriateness of a certain tree species on a particular site. Baker and Broadfoot (1979) provide guidance on profitability for timber management that goes beyond site adaptation. Their recommendations reflect silvicultural practices and hardwood markets 20 years ago, and probably are no longer valid. For example, their productivity criteria suggest that cottonwood may be managed on Amagon, Dundee, Mhoon, and Newellton soils formerly in soybean production, but not on any of the other series included in this study. However, recent improvements in cottonwood pulpwood markets and advances in competition control permit the successful management of this species on a 10-year rotation on formerly row-cropped Sharkey soils (Stanturf et al. 1998).

The dynamic nature of hardwood markets and continual advances in bottomland hardwood silviculture indicate a need for flexible management decision systems. New techniques are emerging, such as using cottonwood as a nurse crop for interplanted Nuttall oak (Schweitzer et al. 1997). Monitoring the growth and maturation of these and other recent plantings on former agricultural lands can serve as the basis for empirically-derived models of stand growth rate and productivity that will provide better species planting recommendations. Further work should focus on determining site index using younger base ages to reflect the increasing utilization of smaller trees.

Conclusions and recommendations

Until empirically-based guides are available, the method of Baker and Broadfoot (1979) represents the best available tool for tree species selection and growth prediction. Ideally, application of this method to a particular site should include an on-site evaluation by a soil scientist or silviculturist, as well as an understanding of site flooding regimes. If this is not possible, we suggest application of the approach described here using soil survey soil data in appropriate regions of the southern United States.

Although the Baker and Broadfoot method was originally developed as a site-specific silvicultural tool, it may be useful to regional land use and economic planners wherever southern bottomland hard-

woods are planted. Overlaying these site index estimates and soil series data in a geographic information system can provide estimates of the contribution of reforestation to environmental remediation and long-term forest-based economic development. Further, reforestation efforts may be focused on marginal agricultural lands where forest productivity and associated economic and environmental benefits are optimized (Lovejoy et al. 1985). Modification of the Baker and Broadfoot (1979) method to conditions in other regions could facilitate the broader use of this tool in land use policy and program decisions.

REFERENCES CITED

- Allen, J.A. 1990. Establishment of bottomland oak plantations on the Yazoo National Wildlife Refuge Complex. *Southern Journal of Applied Forestry* 14:206-210.
- Amacher, G.S., J. Sullivan, L. Shabman, L. Zepp, and D. Grebner. 1998. Reforestation of flooded farmland: Policy implications from the Mississippi River Delta. *Journal of Forestry* 96(5):10-17.
- Aust, W.M., and J.D. Hodges. 1988. A comparison of methods for estimating cherrybark oak site index. *Southern Journal of Applied Forestry* 12:115-116.
- Baker, J.B., and B.G. Blackmon. 1978. Summer fallowing: a simple technique for improving old-field sites for cottonwood. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station Research Paper SO-142.
- Baker, J.B., and W.M. Broadfoot. 1978. A new technique of site selection for hardwoods. *Southern Journal of Applied Forestry* 2:42-43.
- Baker, J.B., and W.M. Broadfoot. 1979. A practical field method of site evaluation for commercially important southern hardwoods. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station General Technical Report SO-26.
- Belli, K.L., J.D. Hodges, and J.C.G. Goetz. 1998. Site productivity assessment for cherrybark oak on minor stream bottoms in Mississippi: A comparison of methods. *Southern Journal of Applied Forestry* 22:7-10.
- Blackmon, B.G., and E.H. White. 1972. Nitrogen fertilization increases cottonwood growth in old-field soil. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station Research Note SO-143.
- Broadfoot, W.M. 1976. Hardwood suitability for and properties of important midsouth soils. U.S. Department of Agriculture Forest Service Research Paper SO-127.
- Francis, J.K. 1984. Soil-site classification for bottomland hardwoods. In: *Proceedings of the 12th Annual Hardwood Symposium* Hardwood Research Council: Asheville, NC.
- Francis, J.K. 1985. Bottomland hardwood fertilization, the Stoneville experience. In: *Proceedings of the Third Biennial Southern Silvicultural Research Conference*. Atlanta, GA. U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station General Technical Report SO-54.
- Groninger, J.W., W.M. Aust, M. Miwa, and J.A. Stanturf. Tree species-soil relationships on marginal soybean lands in the Mississippi Delta. In: *Proceedings of the Tenth Biennial Southern Silviculture Research Conference*, Shreveport, LA. U.S. Department of Agriculture, Forest Service, Southern Research Station General Technical Report SRS. In Press.
- Hodges, J.D. 1997. Development and ecology of bottomland hardwood sites. *Forest Ecology and Management* 90:117-125.
- Hubbard, R.K. and R. Lawrence. 1997. Assessment of forest management effects on nitrate removal by riparian buffer systems. *Transactions of the American Society of Agricultural Engineers* 40:383-391.
- Lovejoy, S.B., J.G. Lee, and D.B. Beasley. 1985. Muddy water and American agriculture: How to best control sedimentation from agricultural land. *Water Resources Research* 21:1065-1068.
- Olmstead, C.J., and D.R. McCurdy. 1989. Factors affecting tree planting by landowners under the CRP, Southern Illinois, 1986-1987. *Journal of Soil and Water Conservation* 44(5):498-500.
- Schweitzer, C.J., J.A. Stanturf, J.P. Shepard, T.M. Wilkins, C.J. Portwood, and L.C. Dorris, Jr. 1997. Large scale comparison of reforestation techniques commonly used in the Lower Mississippi Alluvial Valley: First year results. In: *Proceedings of the 11th Central Hardwood Forest Conference*, Columbia, MO. U.S. Department of Agriculture, Forest Service, North Central Forest Experiment Station General Technical Report NC-188.
- Soil Conservation Service. 1987. Hydric soils of the United States. USDA Soil Conservation Service: Washington, DC.
- Stanturf, J.A., C.J. Schweitzer, and E.S. Gardiner. 1998. Afforestation of marginal agricultural land in the Lower Mississippi River Alluvial Valley. *U.S.A. Silva Fennica* 32:281-297.
- Sternitzke, H.S. 1976. Impact of changing land use on delta hardwood forests. *Journal of Forestry* 74(5):25-27.
- Twedt, D.J., and J. Portwood. 1997. Bottomland hardwood reforestation for neotropical migratory birds: Are we missing the forest for the trees? *Wildlife Society Bulletin* 25:647-652.