

EFFECT OF APICAL MERISTEM CLIPPING ON CARBON ALLOCATION AND MORPHOLOGICAL DEVELOPMENT OF WHITE OAK SEEDLINGS¹

P.P. Kormanik, S.S. Sung, T.L. Kormanik and S.J. Zarnoch²

Abstract—Seedlings from three open-pollinated half-sib white oak seedlots were clipped in mid-July and their development was compared to nonclipped controls after one growing season. In general when data were analyzed by family clipped seedlings were significantly less desirable in three to six of the eight variables tested. Numerically, in all families seedlots, the clipped seedlings were inferior to the controls. Regardless of clipping treatment, the growth variable that affected seedling development the most was the innate ability to develop lammas shoots. Seedlings that produced late season lammas shoots also produced the largest numbers for each of the eight variables tested.

INTRODUCTION

The range of white oak (*Quercus alba*) includes the entire eastern half of the United States because it can adapt to many climatic, edaphic and topographic conditions. It is one of the most commercially valuable oak timber species. White oak is also an important mast producer for wildlife such as squirrels, bluejays, crows, red-headed woodpeckers, deer, turkey, quail, mice, chipmunks, ducks, and raccoons (Rodgers 1990). Currently, white oak plays a major role in multiple use management and is usually included in other biodiversity and restoration ecology efforts.

Unfortunately, artificial regeneration of this valuable and adaptive species has been generally unsuccessful. Even though State nurseries have been providing white oak seedlings for over 50 years, successful plantings or plantations are difficult to find (Boyette 1980; Hill 1986; Kormanik and others 1989). In many instances, planting white oak is not recommended because attempts to artificially regenerate the species consistently fail (Russell 1971; Boyette 1980; Clausen 1983; Hill 1986). Most plantings fail because seedlings show poor growth and survival rates when they compete with herbaceous weeds and other woody, perennial plants. To assess the competitive ability of these seedlings, grading standards are needed. However, developing grading standards for the small white oak seedlings, which range from 25-30 cm in size, has been difficult.

At the Institute of Tree Root Biology (ITRB), we are developing a nursery fertility/management protocol for growing various species of oak and other hardwoods similar to the protocol for loblolly pine (Kormanik and others 1992). In one of the hardwood nursery trials at the Whitehall Experimental Nursery, deer browsed some of the white oak seedlings in mid-July. By mid-August we noted that browsing had affected the morphological development of the seedlings. We found a large variation in growth among the browsed seedlings; however, some of the browsed seedlings were 20 to 30 cm taller than adjacent nonbrowsed seedlings.

Normally, few axillary buds of white oak elongate in the nursery during the year of initiation except perhaps those at the terminal end of the initial flush. This strong apical control on axillary buds (Brown and others 1967) is evident even when cultural treatments result in five flushes instead of the three flushes typical in white oak seedlings. Browsing, however, resulted in rapid axillary bud elongation and lammas-like shoot development in some seedlings in late July. This shoot development is not usually observed on white oak seedlings until mid-August. We were unable to obtain reliable estimates of root and stem development of the browsed seedlings because (1) the number of seedlings affected was limited and (2) the seedlings were from a mixed seedlot.

¹Paper presented at the Eighth Biennial Southern Silvicultural Research Conference, Auburn, AL, Nov. 1-3, 1994. This research was funded in part by Department of Energy Grant DE-AI09-76SR00870. The use of trade or firm names in this article is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.

²Research Forester and Plant Physiologist, USDA Forest Service, Athens, GA; Graduate Student, Soil Sciences Department, University of Georgia, Athens; and Mathematical Statistician, USDA Forest Service, Athens, GA (respectively)

and obvious seed source differences were apparent within the nursery beds.

The objectives of this study were to determine the effect of clipping on (1) seedling height, root collar diameter (RCD) and number of elongated axillary buds, (2) total weight and weight of component parts of seedlings; and (3) root/stem morphology of seedlings.

METHODS

White oak acorns were collected daily from three different mother trees from seedlots 2, 3, and 4 in Clarke County, Georgia, during October-November 1992. The acorns were immediately placed in water and those that floated were discarded. Acorns that sank and showed no evidence of weevil or other damage were placed in plastic bags and stored at 2 to 4 °C until sown.

In late December 1992, acorns were sown in a single raised nursery bed described in detail earlier (Kormanik 1986, Kormanik and others 1990). Based on soil sample analyses from A&L Laboratories, Memphis, TN, the soils were amended with the appropriate fertilizers to obtain the following levels of P, K, Ca, Mg, Cu, Zn, and B, respectively 80, 80, 500, 50, 2.5, 5, and 0.8 ppm, and a pH of 5.8. In other nursery studies these levels had been shown to be appropriate for many species of hardwood including white oak (Kormanik and others 1993; 1994).

Nitrogen, applied as NH_4NO_3 at rates equivalent to 1345 kg/ha (1200 lb/acre) began in mid-May and continued at 10 day intervals until mid-September. Four different rates were used in the 11 applications. The first two applications were at a rate equivalent to 28 kg/ha (25 lb/acre), the third at 56 kg/ha (50 lb/acre), the next six applications were at 168 kg/ha (150 lb/acre) and the final two at 112 kg/ha (100 lb/acre). If needed, seedlings were watered daily until the first growth flush was essentially complete for all seedlots and some seedlings were beginning their second flush. At this point, seedlings received water when moisture tensionmeters registered 30-50 centibars at a depth of 15 to 20 cm. During excessively hot periods, watering occurred at 30 centibars, while during cooler periods, watering was delayed until 50 centibars.

Sixteen continuous rows were assigned to each half-sib mother tree seedlot with 13 acorns placed in each row. This resulted in effective seedling density of 54 to 57/m². Within each half-sib seedlot one row of each of

the seven pairs of rows was randomly chosen to be clipped. The remaining row in each pair became the nonclipped control row. Clipping was completed on July 13, 1993, and consisted of cutting approximately 3 to 5 cm off the current flush. The clipping was internodal and just above either a developed or immature axillary bud. In mid-December 1993, the seedlings were lifted and heights, RCD, first-order lateral root (FOLR) numbers, stem, and root green weights, lammas shoot development and seedling survival rates were obtained.

In this study, each seedlot was treated as a separate experiment. The experiments were analyzed as randomized block designs where the treatments were nonclipped and clipped. Since there were only two treatments, this analysis was equivalent to a paired t-test. Blocks were defined as adjacent pairs of rows in the nursery bed to which the two treatments were assigned at random, one to each row. The experimental unit was a row and the response variables were defined as row means based on thirteen seedlings. Similarly, survival percent and lammas shoot percent were computed on a row basis. Multiple embryos in about 10 to 15 percent of the acorns complicated survival data. Usually the largest seedling was maintained and measured when multiple embryo clusters occurred. Treatment differences were tested at the 0.05 probability level using PROC GLM (SAS Institute 1988).

RESULTS AND DISCUSSION

In this study, the clipping treatment and the innate ability of individual seedlings to develop lammas shoots affected seedling development and survival. Lammas shoot development was the most significant biological response and may be the most important factor in the artificial regeneration of white oak.

Effects of Clipping on Seedling Development

Clipping significantly affected seedling development in some half-sib mother tree seedlots for specific measured variables. However, because the overall responses were sometimes marginal and variable, they may not be of practical significance. Table 1 shows the sample means for clipped and nonclipped white oak seedlings for the eight variables and the associated P values from the analysis of variance. Half-sib seedlot differences in how clipping affected the variables were significant. However, the number of variables significantly different varied by family; the progeny from mother tree 4 were the most consistent. Numbers and

Table 1—Effects of clipping on growth, survival, and P- values for white oak seedlings from three open-pollinated half-sib seedlots

	RCD mm	Hgt cm	FOLR ^a No.	Top wgt g	Root wgt g	Total wgt g	Survival pct	Lammas shoot pct
----- Family 2 ^b -----								
Nonclipped	5.48	30.8	3.02	18.7	43.0	61.8	81.3	37.7
Clipped	5.17	27.7	2.00	14.4	33.1	47.5	73.6	15.6
P-value ^c	0.257	0.185	0.123	0.084	0.042	0.051	0.086	0.009
----- Family 3 ^b -----								
Nonclipped	5.68	29.2	4.21	16.4	37.2	53.6	71.4	37.5
Clipped	4.61	27.0	1.94	12.1	26.4	38.5	63.7	8.9
P-value ^c	0.090	0.465	0.043	0.150	0.086	0.101	0.449	0.001
----- Family 4 ^d -----								
Nonclipped	7.34	53.6	5.51	29.9	49.1	79.0	96.2	69.1
Clipped	6.15	47.4	2.95	19.5	32.6	52.1	91.0	54.1
P-value ^c	0.005	0.059	0.013	0.006	0.002	0.002	0.235	0.179

^a FOLR = first-order lateral root.

^b Based on 7 blocks.

^c The p-value obtained from testing the hypothesis that nonclipped and clipped means are equal.

^d Based on 6 blocks.

proximal diameters of FOLR ≥ 1 mm were used to assess potential competitiveness of white oak seedlings (Kormanik and others 1989). Clipping reduced numbers of FOLR exceeding this minimum diameter which in turn, reduced root weights by approximately 30 percent.

Like deer browsing, clipping resulted in the release of axillary buds, usually immediately below the severed internode. The degree of elongation of these stimulated shoots varied considerably, both within and among half-sib mother tree seedlots. Several weeks passed before the axillary buds were released from their quiescent condition. During this period, the nonclipped seedlings continued to undergo sequential flushes so commonly observed in oak seedlings in nurseries. Even with continued flushing, the overall mean sizes of seedlings (table 1), except those from half-sib seedlot 4 were no larger than normally reported (Boyette 1980; Hill 1986). Past and ongoing research at this location indicates that the time period when FOLR development of nonclipped seedlings accelerates is between flushes 3 and 4 and 4 and 5. Although the study design did not allow for periodic

seedling excavation to follow seasonal FOLR development, the lag period between clipping and axillary bud elongation suggests a wound response in terms of carbon sink relationships. We believe that root growth was restricted while carbon was shunted to repair clipping damage to the stem.

When the study was terminated, the clipped seedlings with no significant elongation had few, if any, FOLR that were ≥ 1 mm in diameter and had poorly developed root system. However, clipping was not the cause of poor root development. We found that nonclipped seedlings exhibiting poor growth were similar to the smaller clipped seedlings in both stem size and root system morphological development. Seedlings with undesirable growth and development had a significant impact on the mean growth variable shown in table 1. Unsatisfactory nursery development of open-pollinated half-sib progeny of white oak below a variables mean value has been shown to be accompanied by poor development of FOLR. Unfortunately, a high percentage of white oak seedlings from specific mother trees may be so affected (Kormanik and others 1989). Many of these inconsistencies were observed in the

past and can now be explained, in part, by lammas shoot development recorded in this study.

Effect of Lammas Shoots on Seedling Development

The response of white oak seedlings to the clipping treatment was of minor significance compared to normal lammas shoot development. It is well known that white oak seedlings tend to develop lammas shoots in late summer and early fall (Busgen and Lutch 1929; Zimmermann and Brown 1971; Kramer and Kozlowski 1979). In our nurseries studies, with white oak when either or both fourth and fifth flushes occur, these flushes can readily be classified as "lammas" rather than "proleptic" shoots (Kramer and Kozlowski 1979). Characteristically, these lammas shoots produce leaves that may be 3- to 7-fold larger in surface area than leaves produced earlier in the growing season. In the past, we have paid little attention to lammas shoots of white oak except to note their presence (Kormanik and others 1989; 1993).

Lammas shoot development is believed to be under genetic control (Cannell and Last 1976; Kramer and Kozlowski 1979). Our study supports this theory because 38 to 69 percent of the nonclipped seedlings produced lammas shoots. Moreover, in seedlots 2 and

3 clipping significantly reduced lammas shoot development. We found lammas shoots always developed from the terminal bud and the shoots from axillary buds released by clipping rarely exhibited typical lammas growth characteristics.

Mean values of the growth variables for clipped and nonclipped seedlings by lammas shoot development show the impact of lammas shoots on stem and root development (table 2). Because lammas shoot growth was a dichotomous response variable (yes or no) and was not an independent experimental variable, we felt that formal statistical tests for differences were inappropriate and might be misleading.

All individuals with lammas shoots were superior in all seedling development characteristics (table 2). Most pronounced differences occurred in FOLR numbers and weight of component seedling parts. These growth responses related to lammas shoot development help to explain early FOLR development patterns in white oak (Kormanik and others 1989; 1993). Normally, diameter growth of FOLR accelerates in late summer or early fall. This acceleration coincides closely with lammas shoot development with white oak seedlings in this study.

Table 2—Effect of lammas shoot development on growth of clipped and nonclipped white oak seedlings

Family	Treat- ment ^a	Shoot ^b growth	RCD mm	Hgt cm	FOLR ^c No.	Top wgt g	Root wgt g	Total wgt g
2	1	0	4.7	24.7	1.4	11.7	28.1	39.8
	1	1	7.7	43.9	5.2	28.0	58.2	86.2
2	0	0	3.9	19.6	0.7	6.5	21.5	28.0
	0	1	8.0	48.1	6.5	37.4	76.2	113.5
3	1	0	4.4	24.9	1.5	10.2	23.6	33.8
	1	1	7.4	48.4	6.8	32.4	56.5	88.9
3	0	0	4.6	20.2	1.9	8.7	24.2	32.8
	0	1	7.3	43.2	7.6	29.0	57.1	86.0
4	1	0	4.5	33.6	0.7	8.9	18.8	27.7
	1	1	7.5	58.8	4.9	28.2	44.3	72.5
4	0	0	5.6	34.0	1.8	9.8	30.0	39.9
	0	1	8.0	62.1	7.1	38.7	57.1	95.9

^a 1 = clipped; 0 = control.

^b 1 = lammas shoot; 0 = NO lammas shoot.

^c FOLR = first-order lateral root.

Table 3—Characteristics for leaves grown on normal shoot and lammas shoot of white oak seedlings.

Leaf type	Leaf area cm ²	Photosynthesis rate umol/m ² .sec	Total leaf photosynthesis nmol/leaf.sec mg/cm ²	Leaf specific weight
Mature normal	23	14.1	32.4	8.6
Mature normal	51	14.5	74.0	9.2
Mature lammas	132	14.6	192.7	11.6
Mature lammas	140	13.5	189.0	9.3

Seedlings exhibiting lammas shoot development were taller and had higher numbers of FOLR regardless of clipping treatment. This observation parallels earlier observations where seedlings with six or more FOLR were two times taller and had RCDs twice as large as those with less than six FOLR. Earlier research had shown even greater variation among root and stem weights where 4- to 7-fold differences in root and stem weight were associated with greater numbers of FOLR (Kormanik and others 1989; 1990). The largest seedlings in all studies frequently had two consecutive lammas shoot episodes.

Based on these data (tables 1 and 2), a direct genetic correlation between FOLR development and lammas shoots with this species appears to exist. Our research supports the hypothesis that the most dominant and codominant oak trees in the forest were those that were able to make rapid juvenile growth (Korstain 1927). This permitted them to get ahead of the herbaceous competition that is encountered after overstory harvesting. Therefore, the innate ability to produce lammas shoots may be significant in white oak seedling establishment. How long seedlings have the potential to produce lammas shoots is unknown. Even a lammas shoot development period of only 3 to 4 years, may contribute to seedling success.

We also measured photosynthesis and found that photosynthetic rates per unit leaf area are the same for the large lammas shoots/leaves and the smaller leaves produced earlier in the summer. The advantage of lammas leaves lies in their greater total leaf area that contributes significantly more carbon for seedling development (table 3).

We have also observed in several ongoing research studies at the ITRB that seedlings that develop lammas shoots the first year may also develop lammas shoots in the second growing season. When grown under proper nursery cultural conditions, transplanted white oak that developed lammas shoots the first and second years, attained heights from 1.7 to 1.8 m after the second growing season.

CONCLUSION

Clipping white oak seedlings does not improve seedling quality and is not recommended for improving seedling size or quality. Although lammas shoot development in nursery stock production has not been studied extensively, this innate trait may be important in early selection of competitive individual seedlings for outplanting. Clipping can significantly reduce the development of lammas shoots. When clipped seedlings develop lammas shoots, total height and RCD are comparable, but weight of other component parts of seedlings may be significantly reduced.

LITERATURE CITED

- Boyette, Warren G. 1980. Performance of ten species of 1-2 oak transplants in North Carolina after six and seven growing seasons. Forest Note 45. Raleigh, NC: North Carolina Forest Service. 4 pp.
- Brown, C.L.; McAlpine, R.G.; Kormanik, P.P. 1967. Apical dominance and form in woody plants: a reappraisal. American Journal of Botany 54: 153-162.

- Buogen, M.; Munch, E.** 1929. The structure and life of forest trees. New York: John Wiley & Sons. 436 pp.
- Carnell, M.G.R.; Last, F.T.** 1976. Tree physiology and yield improvement. London: Academic Press. 567 pp.
- Clausen, K.E.** 1983. English oak grow better than white oak of comparable seedling size. *Tree Planters' Notes* 34:17-19.
- Mill, Joseph A.** 1986. Survival of Pennsylvania State Nursery seedlings 1971-81. pp. 1-4. In: *Proceedings Northeastern Area Nurseryman's Conference; 1986 July 14-17; State College, PA. State College, PA: Northeastern State, Federal and Provincial Nursery Association.*
- Kormanik, P.P.** 1986. Lateral root morphology as an expression of sweetgum seedling quality. *Forest Science* 32:595-604.
- Kormanik, P.P.; Ruehle, J.L.; Muse, H.D.** 1989. Frequency distribution of lateral roots of 1-0 bare-root white oak seedlings. Res. Note SE-353. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest Experiment Station. 5 pp.
- Kormanik, P.P.; Ruehle, J.L.; Muse, H.D.** 1990. Frequency distribution and heritability of first-order lateral roots in loblolly pine seedlings. *Forest Science* 36:802-816.
- Kormanik, P.P.; Sung, S.S.; Kormanik, T.L.** 1992. Controlling loblolly pine seedling growth through carbon metabolism regulation rather than mechanical procedures. pp. 6-11. In: *Proceedings Southern Forest Nursery Association Conference; 1992 July 20-23; Pine Mountain, GA. Macon, GA: Georgia Forestry Commission.*
- Kormanik, P.P.; Sung, S.S.; Kormanik, T.L.** 1993. Toward a single nursery protocol for oak seedlings. pp. 89-98. In: *Proceedings 22nd Southern Forest Tree Improvement Conference; 1993 July 14-17; Atlanta, GA. Atlanta, GA: Southern Forest Tree Improvement Committee.*
- Kormanik, P.P.; Sung, S.S.; Black, C.C.** 1994. Irrigating and fertilizing to grow better nursery seedlings. pp. 115-121. In: *Proceedings Northeastern and Intermountain Forest and Conservation Nursery Associations; 1993 Aug 2-5; St. Louis, MO. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station.*
- Korstain, Clarence F.** 1927. Factors controlling germination and early survival in oaks. *Yale University Bulletin* No. 19. New Haven, CT: School of Forestry.
- Kramer, P.J.; Kozlowski, T.T.** 1979. Physiology of wood plants. London: Academic Press. 567 pp.
- Rodgers, Robert.** 1990. White oak. pp. 605-613. In: *Burns, Russell M.; Honkala, Barbara H., tech. coord. Silvics of North American, Vol. 2, Hardwoods. Washington, DC: U.S. Department of Agriculture, Forest Service. Agriculture Handbook No. 654.*
- Russell, T.E.** 1971. Seeding and planting upland oaks. pp. 49-54. In: *Proceedings of the Oak Symposium; 1971 Aug 16-20; Morgantown, WV. Broomall, PA: U.S. Department of Agriculture, Forest Service, Northeastern Forest Experiment Station.*
- SAS Institute Inc.** 1988. *SAS/STAT User's Guide, Release 6.03 Edition.* Cary, NC: SAS Institute Inc. 1028 pp.
- Zimmermann, M.H.; Brown, C.L.** 1971. *Trees: structure and function.* New York: Springer-Verlag. 336 pp.