VERTICAL DISPLACEMENT OF *IPS LATIDENS* AND *IPS PINI* (COLEOPTERA: SCOLYTIDAE) BY SEMIOCHEMICAL INTERRUPTION

DANIEL R MILLER[1]

Centre for Pest Management, Department of Biological Sciences, Simon Fraser University, Burnaby, British Columbia, Canada V5A 1S6

Abstract

The Canadian Entomologist **132**: 789 – 797 (2000)

The effect of semiochemical interruptants was examined for *Ips latidens* (LeConte) and *Ips pini* (Say) using artificial trees (tall-traps) consisting of an array of seven Lindgren multiple-funnel traps suspended vertically on a rope ladder. *S*-(+)-Ipsdienol reduced the numbers of *I. latidens* captured in (±)-ipsenol-baited funnel traps relative to funnel traps immediately above them and to control tall-traps. The combination of (±)-*exo*-brevicomin and *cis*- and *trans*-verbenol reduced the numbers of *I. pini* captured in (±)-ipsdienol-baited funnel traps relative to funnel traps immediately above them and to control tall-traps. These results provide support for semiochemical-based pest-management tactics that utilize a point source of interruptant at the base of a tree while noting limitations in behavioural effects and efficacy.

Miller DR. 2000. Déplacement vertical d'*Ips latidens* et *Ips pini* (Coleoptera : Scolytidae) par interruption sémiochimique. *The Canadian Entomologist* **132** : 789–797.

Résumé

Nous avons testé les effets d'inhibiteurs sémiochimiques chez *Ips latidens* (LeConte) et *Ips pini* (Say) en utilisant des arbres artificiels (arbres-pièges) composés de sept pièges Lindgren à plusieurs entonnoirs suspendus verticalement sur une échelle de corde. Le *S*-(+)-ipsdiénol a réduit le nombre de *I. latidens* capturés dans des pièges à entonnoirs garnis de (±)-ipsénol comparativement au nombre capturé dans les pièges à entonnoirs installés immédiatement au-dessus et dans les arbres-pièges témoins. La combinaison de (±)-*exo*-brévicomine et de *cis*- et *trans*-verbénol a eu pour effet de réduire le nombre d'*I. pini* capturés dans les pièges à entonnoirs garnis de (±)-ipsdiénol comparativement au nombre capturé dans les pièges témoins situés immédiatement au-dessus et dans les arbres-pièges témoins. Ces résultats justifient le recours à des tactiques d'aménagement sémiochimiques qui utilisent une substance inhibitrice ponctuelle placée à la base d'un arbre, mais soulignent l'importance de tenir compte des limites de l'efficacité de la méthode et de ses effets sur le comportement.

[Traduit par la Rédaction]

Introduction

Ips pini (Say) and *Ips latidens* LeConte (Coleoptera: Scolytidae) are common species of bark beetles that breed in the phloem tissue of dead, dying, or downed lodgepole pine, *Pinus contorta* var. *latifolia* Engelmann (Pinaceae), in western North America (Furniss and Carolin 1980). These species are often present in large numbers during epidemics of the mountain pine beetle, *Dendroctonus ponderosae* Hopkins (Coleoptera: Scolytidae) (Furniss and Carolin 1980). *Ips latidens* generally breeds in the tops and limbs of trees, occasionally infesting dry portions of the main bole in association with either *I. pini* or *D. ponderosae* (Furniss and Carolin 1980; Miller and Borden 1985).

[1] Present address: Forestry Sciences Laboratory, Southern Research Station, USDA Forest Service, 320 Green Street, Athens, Georgia 30602-2044, United States (E-mail: dmiller03@fs.fed.us).

Both *I. pini* and *I. latidens* assemble on hosts in large, single-species aggregations, facilitated by aggregation pheromones (Borden 1982; Birch 1984; Byers 1989). Male *I. latidens* produce predominantly (*R*)-(+)-ipsenol (2-methyl-6-methylene-7-octen-4-ol), and both male and female *I. latidens* are attracted to racemic blends (Miller *et al.* 1991). *Ips pini* uses ipsdienol (2-methyl-6-methylene-2,7-octadien-4-ol) as an aggregation pheromone (Stewart 1975; Birch *et al.* 1980*a*; Lanier *et al.* 1980), preferring a racemic blend in southwestern British Columbia (Miller *et al.* 1996).

Both species exhibit mutual interruption of attraction through their use of pheromones, a mechanism that could ensure ecological and reproductive isolation (Birch 1984; Miller 1991). Ipsdienol, produced by male *I. pini*, interrupts the attraction of *I. latidens* to its pheromone (Miller and Borden 1992). Ipsenol, produced by male *I. latidens*, interrupts the attraction of *I. pini* to its pheromone, ipsdienol (Borden *et al.* 1992).

The present study examines the effect of vertical distance from a point source at the base of a tree on the interruption of pheromone attraction with *I. latidens* and *I. pini*. The concentration of a semiochemical interruptant from a point source at the base of a tree should diminish with increasing vertical distance along a tree bole due to dilution in a three-dimensional air space. This phenomenon should lead to a reduction in pheromone interruption because both species exhibit a dose-dependent response to interruptants (Miller 1991; Miller and Borden 1992).

The experimental design employed artificial "trees" rather than live trees to avoid the possibility of natural beetle attacks and confoundment from additional pheromone sources. Specifically, this study assessed the effect of semiochemical interruptants in reducing the numbers of *I. latidens* and *I. pini* arriving on an artificial "tree," particularly with respect to a source of interruptant located at the base of the tree. A point release source of interruptant(s) at the base of an artificial tree should (1) reduce numbers of landing beetles relative to the upper portions of the tree, and (2) reduce numbers of landing beetles in the upper portions of the treated tree relative to the same portions in untreated controls.

Materials and Methods

In 1986, two experiments were conducted in mature stands of lodgepole pine near Princeton (49°27′N, 120°31′W), British Columbia. In each experiment, eight tall-traps were set in four replicates of two tall-traps per replicate. Each tall-trap consisted of seven eight-unit multiple-funnel traps (Lindgren 1983) (Phero-Tech Inc., Delta, British Columbia) suspended in a vertical array on rungs of a ladder constructed of polypropylene rope (diameter 6 mm) and metal conduit pipe for rungs (length 70 cm, diameter 16 mm), spaced at intervals of about 1 m. Each tall-trap was positioned between trees by a rope-and-pulley suspension system such that no tall-trap was within 5 m of any tree and the collection cup of the bottom funnel trap was about 0.5 m above ground. Tall-traps were set about 25 m apart within a replicate with a spacing of >5 km between replicates.

The design in Experiment 1 approximated a hypothetical scenario involving attacks on trees by *I. latidens* (Fig. 1). The four lowest funnel traps of each tall-trap were baited with the pheromone for *I. latidens*, (±)-ipsenol, on 24 June. Within each replicate, the two lowest funnel traps of one tall-trap were baited on the same date with the interruptant, (+)-ipsdienol, and the tall-trap was designated as "treated." The three uppermost funnel traps were left blank to assess the accuracy of attraction relative to baited funnel traps. Trap catches were collected at intervals of 2–3 weeks until termination of the experiment on 4 August.



FIGURE 1. Design and baiting scheme for artificial trees (tall-traps) consisting of seven eight-unit Lindgren multiple-funnel traps each. Baits included (±)-ipsenol (Is), (±)-ipsdienol (Id), (+)-ipsdienol (+Id), and (±)-*exo*-brevicomin plus *cis*- and *trans*-verbenol (B + V).

Experiment 2 was designed for *I. pini*, which generally infest entire tree boles (Fig. 1). Because few *I. latidens* were captured in the blank funnel traps in Experiment 1, all funnel traps were baited with the pheromone for *I. pini*, (±)-ipsdienol, on 4 August. Within each replicate, the lowest funnel trap of one tall-trap was baited on the same date with the pheromones for *D. ponderosae*, *exo*-brevicomin and *cis*- and *trans*-verbenol (B + V), and the tall-trap was designated as treated. *exo*-Brevicomin and *cis*- and *trans*-verbenol, produced by *D. ponderosae*, interrupt the attraction of *I. pini* to its pheromone (Miller 1991). Trap catches were collected at intervals of 2–3 weeks until termination of the experiment on 2 September.

(±)-Ipsenol and (±)-ipsdienol (chemical purities > 98%) were obtained from Bedoukian Research Inc. (Danbury, Connecticut). BJ Johnston (Department of Chemistry, Simon Fraser University, Burnaby, British Columbia) supplied *S*-(+)-ipsdienol (optical purity 98%, chemical purity > 98%). Glass capillary (±)-*exo*-brevicomin lures (chemical purity > 98%) and polyethylene bubble-cap lures containing a 13:87 mixture of *cis*- and *trans*-verbenol [chemical purities 98%, enantiomeric compositions 83:17 (–):(+)] were supplied by Phero-Tech Inc.

Each ipsenol and ipsdienol lure consisted of eight Microcap® disposable pipettes (2 µL) (Drummond Scientific Co., Broomall, Pennsylvania), each sealed at one end and filled with (±)-ipsenol, (±)-ipsdienol, or (+)-ipsdienol and placed in an open polyethylene, microcentrifuge tube (1.8 mL) (Evergreen Scientific, Los Angeles, California). The release rate of ipsenol and ipsdienol from lures was about 80 µg/d at 24°C. The release rates of verbenols from bubble-cap lures were about 128 µg/d at 25°C, while the release rate of *exo*-brevicomin was 12–31 µg/d at 27°C. All release rates were determined by gravimetric analysis.

Sexes of *I. pini* were determined using declivital characters (Lanier and Cameron 1969), whereas those of *I. latidens* were determined by dissection and examination of

genitalia. Voucher specimens were deposited at the Entomology Museum, Simon Fraser University.

Paired *t* tests were used to compare total catches in selected traps between control and treated tall-traps. For each experiment, trap catch data [transformed by $\ln(Y + 1)$] and sex ratio data (for catches > 5) [transformed by arcsine(Y)] for control and treated tall-traps were subjected separately to two-way analysis of variance (ANOVA), using replicate and trap height as model factors. Fisher's least significant difference (LSD) multiple range test was performed when $P \leq 0.05$. The data were analyzed with the SYSTAT statistical package version 8.0 (SPSS Inc., Chicago, Illinois).

Results

The interruptants significantly reduced trap catches of *I. latidens* and *I. pini*. In Experiment 1, the mean ± SE total catch of 3 ± 1 *I. latidens* in the two bottom traps of the treated tall-traps was less than the mean total catch of 93 ± 38 in the two bottom traps of the control tall-trap ($t_3 = 7.70$, $P = 0.005$). In Experiment 2, the mean ± SE total catch of 8 ± 5 *I. pini* in the bottom trap of the treated tall-traps was less than the mean total catch of 107 ± 50 in the bottom trap of the control tall-trap ($t_3 = 6.52$, $P = 0.007$).

Interruptants in the lower funnel traps also reduced total catches of *I. latidens* and *I. pini* in traps above those treated with interruptants. In Experiment 1, combined total catches of *I. latidens* in ipsenol-baited funnel traps at heights of 3 and 4 m above ground were lower in treated tall-traps than in control tall-traps (mean ± SE total catches of 35 ± 9 and 103 ± 53, respectively) ($t_3 = 3.89$, $P = 0.030$). In Experiment 2, combined total catches of *I. pini* in ipsdienol-baited funnel traps at all heights ≥2 m were lower in treated tall-traps then in control tall-traps (mean ± SE total catches of 338 ± 155 and 642 ± 245, respectively) ($t_3 = 10.19$, $P = 0.002$).

Ips latidens and *I. pini* discriminated between individual pheromone sources within a vertical array of traps. In Experiment 1, the response of *I. latidens* was affected by trap height in both control ($F_{6,18} = 44.71$, $P < 0.001$) and treated ($F_{6,18} = 44.71$, $P < 0.001$) tall-traps, due largely to the unbaited traps at heights of 5–7 m (Fig. 2). As well, catches of beetles in the two lower ipsdienol- and ipsenol-baited funnel traps of the treated tall-trap were lower than catches in the two ipsenol-baited traps immediately above them. Catches in the lower traps were comparable to those in the three uppermost, blank traps. In contrast, there was no difference in trap catches of *I. latidens* among the four ipsenol-baited traps of the control tall-trap. Few *I. latidens* were captured in blank traps at heights of ≥5 m, even though the trap located at a height of 4 m was baited with its pheromone ipsenol.

In Experiment 2, the response of *I. pini* was affected by trap height in both control ($F_{6,18} = 5.00$, $P = 0.004$) and treated ($F_{6,18} = 34.74$, $P < 0.001$) tall-traps (Fig. 3). Catches of *I. pini* in the lower (B + V)- and ipsdienol-baited funnel trap of the treated tall-trap were less than catches in traps baited with ipsdienol alone. There was no difference in mean catches among the traps of the control tall-trap.

In Experiment 1, the sex ratio of *I. latidens* captured in funnel traps was unaffected by trap height in control ($F_{3,8} = 0.56$, $P = 0.655$) or treated ($F_{1,2} = 0.80$, $P = 0.465$) tall-traps. The mean ± SE proportion of male *I. latidens* in trap catches was 0.18 ± 0.02. In Experiment 2, the sex ratio of *I. pini* captured in funnel traps was unaffected by trap height in control ($F_{6,18} = 0.42$, $P = 0.856$) and treated ($F_{6,16} = 1.12$, $P = 0.393$) tall-traps. The mean ± SE proportion of male *I. pini* in trap catches was 0.40 ± 0.01.



FIGURE 2. Interruptive effect of ipsdienol (Id) on catches of *Ips latidens* in ipsenol-baited (1–4) and blank (5–7) funnel traps within tall-traps. Heights of traps determined at middle of each trap. Means, within a figure of seven trap means, followed by a different letter are significantly different (Fisher's LSD test, $P = 0.05$).



FIGURE 3. Interruptive effect of *exo*-brevicomin and verbenols (B + V) on catches of *Ips pini* in ipsdienol-baited funnel traps within tall-traps. Heights of traps determined at middle of each trap. Means, within a figure of seven trap means, followed by a different letter are significantly different (Fisher's LSD test, $P = 0.05$).

Discussion

These results with *I. latidens* and *I. pini* support the contention that variation in bark beetle behaviours, relative to vertical stratification of attacks and pheromone dispersal, is important in the dynamics of bark beetle populations. The behavioural

phenomena involved in vertical segregation of host material by bark beetles have been clearly described for the bark beetle guild in southern pine stands (Smith *et al.* 1993) and likely apply to other pine systems as well (Borden 1982; Birch 1984; Byers 1989). The southern pine beetle, *Dendroctonus frontalis* Zimmerman, generally attacks loblolly pines at heights of 2–6 m above ground (Dixon and Payne 1979; Coster *et al.* 1977). The mid-bole is generally occupied by *Ips calligraphus* (Germar), whereas *Ips grandicollis* (Eichhoff) dominates in lower branches and parts of the mid-bole (Birch and Svihra 1979; Paine *et al.* 1981). *Ips avulsus* (Eichhoff) is found throughout the trunk, dominating in the upper bole, where it stays separate from *I. grandicollis* (Birch and Svihra 1979; Svihra *et al.* 1980; Paine *et al.* 1981).

Host separation of bark beetle species in southern pine forests seems to be facilitated by semiochemicals (Birch *et al.* 1980*b*; Svihra *et al.* 1980; Smith *et al.* 1990). *Dendroctonus frontalis* is attracted by the pheromone frontalin (Vité 1971; Payne *et al.* 1978, 1982; Billings 1985). *Ips avulsus* and *I. calligraphus* are attracted by the pheromone ipsdienol (Vité *et al.* 1978; Smith *et al.* 1990; Kohnle *et al.* 1994), with attraction of *I. calligraphus* enhanced by *cis*-verbenol and host compounds (Renwick and Vité 1972). Attraction of *I. avulsus* is enhanced by (*S*)-(–)-ipsenol (Hedden *et al.* 1976; Kohnle *et al.* 1994) and mixtures of *endo*- and *exo*-brevicomin (Richerson and Payne 1979). *Ips grandicollis* is attracted by (*S*)-(–)-ipsenol and host monoterpenes (Vité *et al.* 1976; Billings 1985; Smith *et al.* 1990; Kohnle *et al.* 1994).

Niche separation is ensured by mutual interruption of pheromone attraction (Birch *et al.* 1980*b*; Svihra *et al.* 1980). The attraction of *I. calligraphus* is interrupted by the pheromone of *I. avulsus*, whereas the attraction of *I. grandicollis* is interrupted by the pheromones of *D. frontalis* and *I. calligraphus* (Birch *et al.* 1980*b*; Svihra *et al.* 1980; Kohnle *et al.* 1994). Attraction of *D. frontalis* is interrupted by the pheromone of *I. grandicollis* (Birch *et al.* 1980*b*).

Vertical stratification in host utilization by semiochemicals has implications for the development of semiochemical-based management tactics. A point source of interruptant at the base of an artificial tree reduced catches of both *I. latidens* and *I. pini* in vertically adjacent traps baited only with their respective pheromones, relative to the same traps in control trees (Figs. 2, 3). These results support the current application of interruptants at the base of a tree in operational tactics designed to prevent attacks by bark beetles over the length of a tree bole. Research trials to reduce infestation levels of *D. ponderosae* have utilized devices releasing the antiaggregation pheromone, verbenone, at heights of 1.5–2 m above ground on boles of lodgepole and ponderosa pine (Amman *et al.* 1989; Lindgren *et al.* 1989; Bentz *et al.* 1989; Gibson *et al.* 1991). A similar approach has been used with interruptants for *Dendroctonus brevicomis* LeConte (Paine and Bertram 1995), *Dendroctonus pseudotsugae* Hopkins (Ross *et al.* 1996), *Dendroctonus rufipennis* (Kirby) (Holsten and Werner 1987), *D. frontalis* (Billings *et al.* 1995), and *Ips paraconfusus* Lanier (Shea and Neustein 1995).

However, such tactics have not gained widespread acceptance due to inconsistent efficacy as exemplified by trials with verbenone on *D. ponderosae* (Borden 1994; Amman and Lindgren 1995). Verbenone significantly reduced infestation levels of *D. ponderosae* in some stands of lodgepole pine but not in others (Lindgren *et al.* 1989; Amman *et al.* 1991; Gibson *et al.* 1991; Shore *et al.* 1992). No significant effect of verbenone on populations of *D. ponderosae* was ever observed in stands of ponderosa pine (Bentz *et al.* 1989; Lister *et al.* 1990; Gibson *et al.* 1991).

The semiochemical ecology of bark beetles is complex, and sophisticated pest-management tactics that specifically address critical factors underlying the variation in beetle behaviour will be needed (Vité and Francke 1992; Borden 1995). Climate, stand structure, and technological limitations affected past trials (Shea *et al.* 1992; Borden 1994; Amman and Lindgren 1995). Vertical stratification in the effect of interruptants

may have played a role in past trials with verbenone as well. Catches of *I. latidens* and *I. pini* in traps above those baited with interruptants were significantly higher than those in traps baited with interruptants (Figs. 2, 3). The interruptants reduced catches but did not preclude significant numbers of landings by beetles along the bole of the tree. Similarly, the use of interruptants in operational trials may have been negated by beetle attacks above the location of the interruptants.

A thorough understanding of bark beetle behaviours associated with a range of critical factors should reveal vulnerable life history points to exploit in pest management (Borden 1989). Probability estimates for the various behaviours associated with semiochemical dispersion patterns, particularly as they fluctuate over time and space, will allow us to refine integrated pest-management tactics. Issues such as release rates from semiochemical devices and the density of release points must be considered and evaluated to address the variation in bark beetle behaviours. Additional classes of semiochemicals such as nonhost odours and pheromones from competing species may yield successes in the future by overcoming the constraints of beetle biology relative to spatial parameters (Borden 1994). These may include combinations of antiaggregative pheromones with pheromones of competing species (Borden *et al.* 1992; Paine and Bertram 1995) or the addition of nonhost compounds such as 4-allylanisole (Hayes *et al.* 1994) and green leaf volatiles (Poland *et al.* 1998).

Acknowledgements

I thank John H Borden for his unconditional support and effusive enthusiasm during the course of this and numerous other research efforts. B Sullivan, GL DeBarr, and P de Groot provided valuable reviews of the manuscript. LJ Chong, C Matteau, and L Wheeler assisted with technical support. This research was supported in part by an HR MacMillan Family Fund Fellowship to DRM, the Natural Sciences and Engineering Research Council of Canada, and the Science Council of British Columbia.

References

Amman GD, Lindgren BS. 1995. Semiochemicals for management of mountain pine beetle: status of research and application. pp. 14–22 *in* SM Salom, KR Hobson (Tech Eds), Application of Semiochemicals for Management of Bark Beetle Infestations, Proceedings of an Informal Conference. *United States Department of Agriculture Forest Service General Technical Report* **INT-GTR-318**

Amman GD, Thier RW, McGregor MD, Schmitz RF. 1989. Efficacy of verbenone in reducing lodgepole pine infestation by mountain pine beetles in Idaho. *Canadian Journal of Forest Research* **19**: 60–4

Amman GD, Thier RW, Weatherby JC, Rasmussen LA, Munson AS. 1991. Optimum dosage of verbenone to reduce infestation of mountain pine beetle in lodgepole pine stands in central Idaho. *United States Department of Agriculture Forest Service Research Paper* **INT-446**

Bentz B, Lister CK, Schmid JM, Mata SA, Rasmussen LA, Haneman D. 1989. Does verbenone reduce mountain pine beetle attacks in susceptible stands of ponderosa pine? *United States Department of Agriculture Forest Service Research Note* **RM-495**

Billings RF. 1985. Southern pine bark beetles and associated insects: effects of rapidly-released host volatiles on response to aggregation pheromones. *Zeitschrift für Angewandte Entomologie* **99**: 483–91

Billings RF, Berisford CW, Salom SM, Payne TL. 1995. Applications of semiochemicals in the management of southern pine beetle infestations: current status of research. pp. 30–8 *in* SM Salom, KR Hobson (Tech Eds), Application of Semiochemicals for Management of Bark Beetle Infestations, Proceedings of an Informal Conference. *United States Department of Agriculture Forest Service General Technical Report* **INT-GTR-318**

Birch MC. 1984. Aggregation in bark beetles. pp. 331–53 *in* WJ Bell, RT Cardé (Eds), *Chemical ecology of insects*. Sunderland: Sinauer Associates

Birch MC, Svihra P. 1979. Novel approaches to forest insect control: exploiting olfactory interactions between species of Scolytidae. pp. 135–8 *in* WE Walters (Ed), *Current topics in forest entomology*. New York: John Wiley and Sons, Inc.

Birch MC, Light DM, Wood DL, Browne LE, Silverstein RM, Bergot BJ, Ohloff G, West JR, Young JC. 1980*a*. Pheromonal attraction and allomonal interruption of *Ips pini* in California by the two enantiomers of ipsdienol. *Journal of Chemical Ecology* **6**: 703–17

Birch MC, Svihra P, Paine TD, Miller JC. 1980*b*. Influence of chemically mediated behavior on host tree colonization by four cohabiting species of bark beetles. *Journal of Chemical Ecology* **6**: 395–414

Borden JH. 1982. Aggregation pheromones. pp. 74–139 *in* JB Mitton, KB Sturgeon (Eds), *Bark beetles in North American conifers: a system for the study of evolutionary biology*. Austin: University of Texas Press

——— 1989. Semiochemicals and bark beetle populations: exploitation of natural phenomena by pest management strategists. *Holarctic Ecology* **12**: 501–10

——— 1994. Future of semiochemicals for the management of bark beetle populations. pp. 5–10 *in* PJ Shea (Technical Coordinator), Proceedings of the Symposium on Management of Western Bark Beetles with Pheromones: Research and Development. *United States Department of Agriculture Forest Service General Technical Report* **PSW-GTR-150**

——— 1995. From identifying semiochemicals to developing a suppression tactic: a historical review. pp. 3–10 *in* SM Salom, KR Hobson (Tech Eds), Application of Semiochemicals for Management of Bark Beetle Infestations, Proceedings of an Informal Conference. *United States Department of Agriculture Forest Service General Technical Report* **INT-GTR-318**

Borden JH, Devlin DR, Miller DR. 1992. Synomones of two sympatric species deter attack by the pine engraver, *Ips pini* (Coleoptera: Scolytidae). *Canadian Journal of Forest Research* **22**: 381–7

Byers JA. 1989. Chemical ecology of bark beetles. *Experientia* **45**: 271–83

Coster JE, Payne TL, Hart ER, Edson LJ. 1977. Aggregation of the southern pine beetle in response to attractive host trees. *Environmental Entomology* **6**: 725–31

Dixon WN, Payne TL. 1979. Sequence of arrival and spatial distribution of entomophagous and associate insects of southern pine beetle-infested trees. *Texas Agriculture Experiment Station Miscellaneous Publication* **1432**

Furniss RL, Carolin VM. 1980. Western forest insects. *United States Department of Agriculture Forest Service Miscellaneous Publication* **1339**

Gibson KE, Schmitz RF, Amman GD, Oakes RD. 1991. Mountain pine beetle response to different dosages in pine stands of western Montana. *United States Department of Agriculture Forest Service Research Paper* **INT-444**

Hayes JL, Strom BL, Roton LM, Ingram LL Jr. 1994. Repellent properties of the host compound 4-allylanisole to the southern pine beetle. *Journal of Chemical Ecology* **20**: 1595–1615

Hedden R, Vité JP, Mori K. 1976. Synergistic effect of a pheromone and a kairomone on host selection. *Nature (London)* **261**: 696–7

Holsten EH, Werner RA. 1987. Use of MCH bubblecaps in preventing spruce beetle attacks in Alaska. *United States Department of Agriculture Forest Service Forest Pest Management Technical Report* **R10-12**

Kohnle U, Vité JP, Meyer H, Francke W. 1994. Response of four American engraver bark beetles, *Ips* spp. (Col., Scolytidae), to synthetic racemates of chiral pheromones. *Journal of Applied Entomology* **117**: 451–6

Lanier GN, Cameron EA. 1969. Secondary sexual characters in the North American species of the genus *Ips* (Coleoptera: Scolytidae). *The Canadian Entomologist* **101**: 862–70

Lanier GN, Classon A, Stewart T, Piston JJ, Silverstein RM. 1980. *Ips pini*: the basis for interpopulational differences in pheromone biology. *Journal of Chemical Ecology* **6**: 677–87

Lindgren BS. 1983. A multiple-funnel trap for scolytid beetles. *The Canadian Entomologist* **115**: 299–302

Lindgren BS, Borden JH, Cushon GH, Chong LJ, Higgins CJ. 1989. Reduction of mountain pine beetle (Coleoptera: Scolytidae) attacks by verbenone in lodgepole pine stands in British Columbia. *Canadian Journal of Forest Research* **19**: 65–8

Lister CK, Schmid JM, Mata SA, Haneman D, O'Neil C, Pasek J, Sower L. 1990. Verbenone bubble caps ineffective as a preventative strategy against mountain pine beetle attacks in ponderosa pine. *United States Department of Agriculture Forest Service Research Note* **RM-501**

Miller DR. 1991. Reproductive and ecological isolation: community structure in the use of semiochemicals by pine bark beetles (Coleoptera: Scolytidae). Ph.D. thesis, Simon Fraser University, Burnaby, British Columbia

Miller DR, Borden JH. 1985. Life history and biology of *Ips latidens* (LeConte) (Coleoptera: Scolytidae). *The Canadian Entomologist* **117**: 859–71

——— 1992. (*S*)-(+)-Ipsdienol: interspecific inhibition of *Ips latidens* (LeConte) by *Ips pini* (Say) (Coleoptera: Scolytidae). *Journal of Chemical Ecology* **18**: 1577–82

Miller DR, Borden JH, King GGS, Slessor KN. 1991. Ipsenol: an aggregation pheromone for *Ips latidens* (LeConte) (Coleoptera: Scolytidae). *Journal of Chemical Ecology* **17**: 1517–27

Miller DR, Borden JH, Slessor KN. 1996. Enantiospecific pheromone production and response profiles for populations of pine engraver, *Ips pini* (Say) (Coleoptera: Scolytidae), in British Columbia. *Journal of Chemical Ecology* **22**: 2157–72

Paine TD, Bertram SL. 1995. Management potential of semiochemicals for protection of trees from western pine beetle. pp. 11–13 *in* SM Salom, KR Hobson (Tech Eds), Application of Semiochemicals for Management of Bark Beetle Infestations, Proceedings of an Informal Conference. *United States Department of Agriculture Forest Service General Technical Report* **INT-GTR-318**

Paine TD, Birch MC, Svihra P. 1981. Niche breadth and resource partitioning by four sympatric species of bark beetles (Coleoptera: Scolytidae). *Oecologia* **48**: 1–6

Payne TL, Coster JE, Richerson JV, Edson LJ, Hart ER. 1978. Field response of the southern pine beetle to behavioral chemicals. *Environmental Entomology* **7**: 578–82

Payne TL, Richerson JV, Dickens JC, West JR, Mori K, Berisford CW, Hedden RL, Vité JP, Blum MS. 1982. Southern pine beetle: olfactory receptor and behavior discrimination of enantiomers of the attractant pheromone frontalin. *Journal of Chemical Ecology* **8**: 873–81

Poland TM, Borden JH, Stock AJ, Chong LJ. 1998. Green leaf volatiles disrupt responses by the spruce beetle, *Dendroctonus rufipennis*, and the western pine beetle, *Dendroctonus brevicomis* (Coleoptera: Scolytidae) to attractant-baited traps. *Journal of the Entomological Society of British Columbia* **95**: 17–24

Renwick JAA, Vité JP. 1972. Pheromones and host volatiles that govern aggregation of the six-spined engraver beetle, *Ips calligraphus*. *Journal of Insect Physiology* **18**: 1215–19

Richerson JV, Payne TL. 1979. Effects of bark beetle inhibitors on landing and attack behavior of the southern pine beetle and beetle associates. *Environmental Entomology* **8**: 360–4

Ross DW, Gibson KE, Thier RW, Munson AS. 1996. Optimal dose of an antiaggregation pheromone (3-methylcyclohex-2-en-1-one) for protecting live Douglas-firs from attack by *Dendroctonus pseudotsugae* (Coleoptera: Scolytidae). *Journal of Economic Entomology* **89**: 1204–7

Shea PJ, Neustein M. 1995. Protection of a rare stand of Torrey pine from *Ips paraconfus*. pp. 39–43 *in* SM Salom, KR Hobson (Tech Eds), Application of Semiochemicals for Management of Bark Beetle Infestations, Proceedings of an Informal Conference. *United States Department of Agriculture Forest Service General Technical Report* **INT-GTR-31**

Shea PJ, McGregor MD, Daterman GE. 1992. Aerial application of verbenone reduces attack of lodgepole pine by mountain pine beetles. *Canadian Journal of Forest Research* **22**: 436–41

Shore TL, Safranyik L, Lindgren BS. 1992. The response of mountain pine beetle (*Dendroctonus ponderosae*) to lodgepole pine trees baited with verbenone and *exo*-brevicomin. *Journal of Chemical Ecology* **18**: 533–41

Smith MT, Payne TL, Birch MC. 1990. Olfactory-based behavioral interactions among five species in the southern pine bark beetle group. *Journal of Chemical Ecology* **16**: 3317–31

Smith MT, Salom SM, Payne TL. 1993. The southern pine bark beetle guild: an historical review of the research on the semiochemical-based communication system of the five principal species. *Virginia Agricultural Experiment Station Bulletin* **93-4**

Stewart TE. 1975. Volatiles isolated from *Ips pini*: isolation, identification, enantiomeric composition, biological activity, and the enantiomeric composition of other insect pheromone alcohols and bicyclic ketals. M.Sc. thesis, College of Environmental Science and Forestry, State University of New York, Syracuse, New York

Svihra P, Paine TD, Birch MC. 1980. Interspecific olfactory communications in southern pine beetles. *Naturwissenschaften* **67**: 518–9

Vité JP. 1971. Pest management systems using synthetic pheromones. *Contributions of the Boyce Thompson Institute* **24**: 343–50

Vité JP, Francke W. 1992. Research needs and technology transfer. *Journal of Applied Entomology* **114**: 212–5

Vité JP, Hedden R, Mori K. 1976. *Ips grandicollis*: field responses to the optically pure pheromone. *Naturwissenschaften* **63**: 582–3

Vité JP, Ohloff G, Billings RF. 1978. Pheromonal chirality and integrity of aggregation response in southern species of the bark beetle *Ips* sp. *Nature (London)* **272**: 817–8

(Date received: 10 January 2000; date accepted: 29 May 2000)