



United States
Department of
Agriculture
Forest Service

FOREST PEST MANAGEMENT

Technology Update

Southern Region, USDA Forest Service, 1720 Peachtree Rd., N.W. Atlanta Ga. 30367

Forestry Pesticide Fact Sheet No. 7

EFFECTIVENESS OF 2,4-D AND PICLORAM AS FORESTRY HERBICIDES

by

Daniel G. Neary, Parshall B. Bush, Jerry L. Michael, and John W. Taylor, Jr.

Foresters use 2,4-D, alone or in combination with picloram, as often as any herbicide in the South. An active analog of the plant growth hormone indole acetic acid, 2,4-D is used as a selective herbicide against broadleaved weeds. In most forest soils in the South where organic matter, moisture, and temperature are adequate, 2,4-D degrades rapidly. Amine and salt formulations most commonly used in forestry do not change readily. While 2,4-D will leach under some soil conditions, concentrations found in forest streams are infrequently detected at levels approaching the 100 ppb water quality standard.

The herbicide 2,4-D is only slightly toxic to most aquatic life and very low in toxicity to animals. It does not bioaccumulate and is readily metabolized by plants and soil microorganisms. A variety of formulations are available, including low volatile esters, high volatile esters, water-soluble amines, oil-soluble amines, the parent acid, and inorganic salts.

Picloram is a non-selective, wide-spectrum herbicide used for control of woody plants and most broadleaved weeds. This herbicide is used mainly in site preparation, for removal of undesirable hardwoods and most frequently in combination with 2,4-D. Picloram is low in toxicity to soil microorganisms, fish, birds, and mammals. Readily absorbed by plant roots, it is translocated through plants where it can accumulate and slowly degrade. This herbicide, like 2,4-D does not bioaccumulate and is readily excreted by animals.

Picloram is moderately to highly persistent in forest soils with half life ranging from one month in warm, humid regions to several years in arid or cold climates. Picloram is soluble in water and has a high potential to move off-site

under the right conditions. Some nontarget plants can be affected by low concentrations in water.

The effectiveness of 2,4-D and picloram in controlling important forest weeds has been documented by a number of studies. The more important ones are listed in this fact sheet. Contact the authors of this fact sheet for more complete listing or for any other information regarding the chemicals.

REFERENCES

Bovey, Rodney W. Response of selected woody plants in the United States to herbicides. Agriculture Handbook 493. Washington, D.C., U.S. Department of Agriculture; 1978. 101 p.

Byrd, B.C.; Foster, J. Pine growth twelve years following site preparation with picloram plus 2,4-D. Proceedings of the Southern Weed Science Society 35:200-209. 1982.

Cantrell, R.L.; Flinchum, D.M.; Neat-y, D.G. In press. Individual Tree Treatments Using Herbicides for Control of Turkey Oak (*Quercus laevis* Walt.) Southern Journal of Applied Forestry.

Coble, H.D.; Upchurch, R.P.; Keaton, J.A. Influence of time and method of application on turkey oak response to picloram plus 2,4-D. Weed Science 17:87-91. 1969.

Fitzgerald., C.H.; Reed, T.M.; Fortson, J.C. Foliar applied herbicides for site preparation in Georgia Piedmont types. Proceedings of the Southern Weed Science Society 35:161-174. 1982.

Hamel, Dennis R. Forest management chemicals. Agriculture Handbook 585. Washington, D.C.; Department of Agriculture; 1983. 645 p.

Haymond, Jacqueline L. Herbicides for forestry. Clemson University, Cooperative **Extension Service**. 1984 (rev.) 36 p.

Holt, H.A.; Voeller, JE. Hardwood control trials with the Hypod-Hatchet. Proceedings of the **Southern Weed Science Society** **25:270-276**. 1972.

Miller, **J.H.** Hardwood control using **pelleted** herbicides and burning. **Proceedings** of the **Southern Weed Science Society**. **35:210-215**. 1982.

Neary, D.G.; Douglass, JE.; **Ruehle**, J.L.; Fox, W. Converting rhododendron-laurel thickets to white pine using a picloram herbicide and **mycorrhize**-inoculated seedlings. Southern Journal of Applied Forestry **8:163-168**. 1984.

Peevy, F.A. Wide-spaced injections and herbicidal mixtures for controlling weed trees. **Proceedings** of the Southern Weed **Science Society** **24:263-267**. 1971.

Peevy, F.A. **Bromacil** and picloram under southern (U.S.) upland hardwoods. Weed Science **21:54-56**. 1973.

Peevy, EA. Injection treatment for killing bottomland hardwoods. Weed Science **20:566-568**. 1972.

Shipman, RD. Individual white oaks treated with **stem**-injected and soil-applied herbicides--cost and effectiveness. Down to Earth **28(4):19-25**. 1973.

Shipman, RD. Selective herbicide treatment of striped maple. **Proceedings** of the Northeastern Weed Science Society 36217220.1982.

Sterrett, J.P. Injection of red maple and hickory with picloram, 2,4-D and 2,4,5-T. Down to Earth **25(2)**: 18-21. 1969.

This fact sheet was prepared by Daniel G. Neary, USDA Forest Service, Southeastern Forest Experiment Station, **(904-392-1951)**; **Parshall** B. Bush, University of Georgia, (404-542-7690); Jerry L. Michael, USDA Forest Service, Southern Forest Experiment Station, **(205-826-8700)**; and John W. Taylor, Jr., USDA Forest Service, Southern Region, (404-347-2718). For further information, contact any of the authors at the listed telephone numbers.

The use of trade, **firm** or corporation names in this publication is for the information and convenience of the reader. Such use does not constitute an official endorsement or approval by the U.S. Department of Agriculture of any product to the exclusion of others that may be suitable.

CAUTION: Use pesticides selectively and **carefully**. Follow **recommended** practices for the disposal of surplus pesticides and pesticide containers. Some States have restrictions on the use of certain pesticides. Check with your **local** and State regulations. **Also**, because registration of pesticides are under constant review by the Environmental Protection Agency. Consult your State forestry agency, county agent or State extension agent to be sure the intended use is still registered.