

United States  
Department of  
Agriculture

Forest Service



Southern  
Research Station

Resource Bulletin  
SRS-22

# Forest Resources of Puerto Rico, 1990

Peter A. Franco, Peter L. Weaver, and  
Susan Eggen-McIntosh



---

**The Authors:**

---

**Peter A. Franco** is a Research Forester with the Forest Inventory and Analysis Research Work Unit, Southern Research Station, U.S. Department of Agriculture, Forest Service, Starkville, MS 39759;

**Peter L. Weaver** is a Research Forester with the International Institute of Tropical Forestry, Southern Research Station, U.S. Department of Agriculture, Forest Service, Río Piedras, PR 00928;

and **Susan Eggen-McIntosh** is a former Research Forester with the Forest Inventory and Analysis Research Work Unit, Southern Research Station, U.S. Department of Agriculture, Forest Service, Starkville, MS 39759.

October 1997

Southern Research Station  
P.O. Box 2680  
Asheville, North Carolina 28802

## ERRATA

An error occurred in the preparation of camera copy for Resource Bulletin SRS-22, *Forest Resources of Puerto Rico, 1990* by Peter A. Franco, Peter L. Weaver, and Susan Eggen-McIntosh. On the inside front cover, the address for Peter L. Weaver should have read as follows: "Peter L. Weaver is a Research Forester with the International Institute of Tropical Forestry, U.S. Department of Agriculture, Forest Service, Río Piedras, PR 00928."

—SRS, October 1997

---

## ERRATA

An error occurred in the preparation of camera copy for Resource Bulletin SRS-22, *Forest Resources of Puerto Rico, 1990* by Peter A. Franco, Peter L. Weaver, and Susan Eggen-McIntosh. On the inside front cover, the address for Peter L. Weaver should have read as follows: "Peter L. Weaver is a Research Forester with the International Institute of Tropical Forestry, U.S. Department of Agriculture, Forest Service, Río Piedras, PR 00928."

—SRS, October 1997



# Contents

	Page
<b>Introduction</b> .....	1
<b>Previous Forest Area Trends</b> .....	2
<b>Highlights</b> .....	3
<b>The 1990 Forest Survey</b> .....	4
Area .....	4
Stand Size .....	6
Species Composition .....	6
Basal Area Distribution .....	7
Timber Volume .....	8
Sawtimber Volume .....	10
Trends in Secondary Forests .....	11
Trends in Coffee Shade Forests .....	12
Abandoned coffee shade forest .....	12
Active coffee shade forest .....	13
<b>Conclusions</b> .....	14
<b>Literature Cited</b> .....	14
<b>Appendix</b> .....	15
<b>Survey Methods</b> .....	15
1980 Survey .....	15
1985 Update .....	15
1990 Resurvey .....	15
<b>Reliability of the Data</b> .....	16
<b>Definitions</b> .....	16
<b>Conversion Factors</b> .....	19
<b>List of Tree Species Included in Standard Tables</b> .....	20
<b>Index of Tables</b> .....	22
Tables 1-26 .....	23



# Forest Resources of Puerto Rico, 1990

by Peter A. Franco, Peter L. Weaver, and Susan Eggen-McIntosh

## Introduction

In 1980, the Southern Forest Experiment Station, Forest Inventory and Analysis Research Work Unit (SOFIA), in cooperation with the International Institute of Tropical Forestry, established a forest survey of the Commonwealth of Puerto Rico. The primary objective was to assess the timber-production potential of the island's forest resources. The main island, occupying about 8900 square kilometers, was partitioned into commercial and noncommercial survey regions (fig. 1). Field sampling was concentrated in the commercial survey region, which included all areas having

the potential to produce commercial timber. The commercial region was delineated to exclude urban areas, agricultural lands, critical watersheds, areas with adverse site conditions, such as excessive rainfall and excessive slope, and other areas not likely, either for physiographic or economic reasons, to sustain commercial timber production.

Because complete photographic coverage was unavailable for the 1980 survey, aerial photography was acquired for a 1985 update of forest area estimates of both the commercial and noncommercial survey regions. The 1985 update provided more current estimates of forest area by detailed




Figure 1—Exclusions from the commercial survey region of Puerto Rico (adapted from Birdsey and Weaver 1982).

cover classes for the commercial region, refined forest area estimates for the noncommercial region, and estimated rates of change in the timberland area (Birdsey and Weaver 1987). In 1990, another forest survey was conducted in the commercial region of Puerto Rico. This publication describes the results of that survey and changes that occurred within the region between 1980 and 1990.


The forests of Puerto Rico were previously classified according to the Life Zone system (Ewel and Whitmore 1973, Holdridge 1967). The primary advantage of this classification system is that it provides a means of comparison among tropical forests based on climatic data (Birdsey and Weaver 1982). The commercial region is located in only two of the island's six Life Zones, the Subtropical Moist Forest and the Subtropical Wet Forest. The Subtropical Moist Forest occupies about 58 percent of Puerto Rico; the Subtropical Wet Forest covers about 23 percent.

Broad soil associations, which reflect geologic origin, formed the basis for stratifying the data within both Life

Zones (fig. 2) because they yielded the most important prior knowledge of the commercial survey region (Birdsey and Weaver 1982). Four soil groups were recognized in the surveys. Soils of volcanic origin were divided into deep and shallow clays. Together, these soils make up about 70 percent of the commercial region. Granitic soils and limestone soils account for about 10 and 20 percent of the commercial region, respectively. Forest area statistics for both the 1980 and 1990 surveys were compiled using this stratification rather than the county-based system used in the United States.

**Previous Forest Area Trends**

In the early 16th century, Puerto Rico's 890 000 hectares of land were covered with mature forests containing a wide variety of species. These forest formations were typical of the Caribbean region at the time of discovery (Durland 1929, Little and others 1974, Little and Wadsworth 1964). In the years that followed, European settlers cleared most of the



**Subtropical Life Zones:**



<b>Wet</b>	<b>Moist</b>	
		Deep volcanic soils
		Shallow volcanic soils
		Granitic soils
		Limestone soils

Figure 2—Life Zones and soil groups in the commercial survey region of Puerto Rico (adapted from Birdsey and Weaver 1982).

land and coastal forests for pasture and cropland. Although the clearing of forests for timber, fuelwood, and charcoal had modified the land, clearing for agriculture was far the most destructive in terms of the area affected (Birdsey and Weaver 1982). By 1828, pasture and cultivated land represented one-third of Puerto Rico's total land area (Wadsworth 1950). To accommodate the rapidly growing population, most of the remaining mature forests were cleared in the 19th century (fig. 3). By 1899, forest cover was reduced to 182 000 hectares, with pasture alone accounting for 55 percent of the land area. Cultivating coffee beneath shade trees (coffee shade) had recently begun; this practice used an additional 77 000 hectares (Wadsworth 1950). Tree cover continued to decline slowly over the next 40 years, greatly diminishing the capability of the forests to provide products other than fuelwood and charcoal.




Figure 3—Area with tree cover in Puerto Rico for selected years 1828-1990 (adapted from Birdsey and Weaver 1987).

After 1916, another period of heavy cutting caused forest area to decline sharply, depleting tree cover to 81 000 hectares by 1931 (Gill 1931). By the late 1940's, Puerto Rico's forests had dwindled to only 6 percent of the total land area, and coffee shade occupied about the same area (Koenig 1953). The 1950's marked the beginning of a period of recovery for the forests of Puerto Rico. Abandoned cropland and pasture on eroded hillsides began reverting to secondary forest during this time. Widespread reversion from agricultural land to secondary forest was the primary reason Puerto Rico's total forest area increased to over 30 percent by 1980. These reverted areas, mainly abandoned hillside farms in the mountainous region of the main island (Birdsey and Weaver 1982, 1987), comprise the forest resource addressed in this study.

## Highlights

The 10 years between the 1980 and 1990 surveys was a period of major change in Puerto Rico's timberland, one that reveals a positive outlook for the future quality and utility of the island's timber resources. The most noticeable areal trend was the substantial increase in secondary forests from less than one-half to more than three-fourths of the timberland area. Reverted cropland and pasture accounted for about two-thirds of the increase in secondary forest area. Reclassification of former abandoned coffee shade forest stands also contributed to this increase.

Accompanying the expansion of secondary forests was a reduction in the areas of both abandoned and active coffee shade forests. Most of the abandoned coffee shade lands not reclassified as secondary forest were converted to other agricultural uses. Some formerly active coffee shade areas were abandoned since 1980, but most of the decline in coffee shade forests resulted from the replacement of coffee with other agricultural crops.

Some of the most dramatic areal trends resulted from the reclassification of some forest types (coffee shade and secondary forest). This reclassification helps illustrate the frequently changing land use patterns found in Puerto Rico, where people recurrently clear their land for agriculture or pasture. The island's high population density (about 1,050 people per square mile) and the high proportion of privately owned land (95 percent) equate to many small landholdings, numerous property sales, and an ever changing mosaic of land uses. Previously abandoned lands, once sold, may be placed in agricultural use, such as fruit trees or vegetable crops, used for pasture, or developed for a homesite or commercial venture.

In terms of timberland coverage, sawtimber-size stands gained a slight dominance over poletimber stands, whereas sapling-seedling stands declined. In secondary forests, sapling-seedling stands gained 4600 hectares by 1990, though their proportion of secondary forest area fell from 43 to 26 percent. Poletimber stands covered only an additional 2700 hectares in 1990, the same area by which sapling-seedling stands decreased. These trends reflect the emergence of many of the former poletimber-size stands (the dominant stand size in 1980) into sawtimber-size stands and the ascendance of the former sapling-seedling stands into the poletimber stand-size class.

Trends in basal area and stocking showed an overall improvement in Puerto Rico's timber resource, both in terms of quantity and quality. The combined basal area of poletimber and sawtimber trees was substantially higher, increasing from 1.0 to 1.5 million square meters. Average basal area per hectare of sawtimber-size trees rose most

sharply in secondary forests, whereas average basal area of poletimber had the greatest gains in abandoned coffee shade areas. The average basal area of sapling-size trees across timberland remained the same since 1980 but diminished slightly in secondary forests. The average level of stocking for both sawtimber-size and poletimber-size stands was higher in 1990. Sawtimber and poletimber stands with basal areas of 25 or more square meters per hectare covered 14 500 hectares, compared to 8500 hectares in 1980. Timberland was not only better stocked, but stocked with better quality trees, as supported by a greater proportion of basal area in growing-stock trees. More than 70 percent of timberland basal area was comprised of growing-stock quality trees, compared to 50 percent in 1980.

Increases in volume from 1980 to 1990 were remarkable and offered further evidence of a higher quality timber resource. These notable gains in timber volume resulted from both the growth of the timber surveyed and the lack of commercial timber extraction. The average timber volume per hectare in timberland doubled, with growing-stock trees contributing the majority of this accrual. Furthermore, two-thirds of the timber volume in all forest classes was growing-stock volume. In secondary forests, growing-stock volume was about evenly divided between sawtimber and poletimber trees. Sawtimber volume in timberland approximately doubled, both in terms of total volume (2.6 million cubic meters) and volume per hectare (18 cubic meters). The volume of sawtimber on secondary forests nearly tripled to 1.8 million cubic meters, representing an increase from 11 to 16 cubic meters per hectare. On a per hectare basis, abandoned coffee shade forest had the greatest increase in timber volume (53 to 102 cubic meters) and growing-stock volume (28 to 66 cubic meters), whereas active coffee shade forest had the greatest increase in sawtimber volume (7 to 26 cubic meters).

Much of the notable increase in sawtimber volume may be attributed to the ascendance of formerly poletimber-size trees into the sawtimber-size class. Likewise, many of the previously submerchantable trees have since crossed the threshold of merchantability into either the poletimber or sawtimber-size class. The lack of commercial timber harvesting on the island has particularly favored the accumulation of volume in the sawtimber component.

## The 1990 Forest Survey

### Area

Puerto Rico's total forest area increased from 279 000 hectares in 1980 to 287 000 hectares in 1990. Nonforest land uses amounted to 68 percent of the total land area, slightly less than in 1980. Forest area in the commercial region increased by more than 17 000 hectares, whereas that in the noncommercial region decreased by about 9 000 hectares. The positive net change in the commercial region's forested area resulted because reversions of cropland and pasture exceeded the clearing of forested areas to nonforest uses. Forest land classified as timberland increased to almost 144 000 hectares. Since 1980, about 35 000 hectares reverted from cropland, pasture, and other nonforest land uses to timberland. At the same time, about 21 000 hectares of timberland were cleared for nonforest uses. Therefore, the net increase in timberland area was about 14 000 hectares.

Nearly four-fifths of the commercial region's forests were classified as secondary forest compared to less than one-half in 1980 (fig. 4). The remaining 30 000 hectares of timberland represented either former or current coffee producing areas (fig. 5). Over two-thirds of that timberland




Figure 4—Timberland area by forest class, Puerto Rico, 1980 and 1990.




Figure 5—Distribution of components of timber volume, Puerto Rico timberland, 1990.

Area was secondary forest land resulting specifically from the abandonment of coffee production under shade trees (abandoned coffee shade). Only about 7000 hectares in 1990 supported forest stands, which were being used as coffee shade (active coffee shade). Reversions from nonforest land uses accounted for almost one-third of the 30 000 hectares of secondary forest land in the commercial region. Most of the other 67 percent of secondary forest land was classified as such in 1980. Other minor additions to secondary forest land came from the classification of active and abandoned coffee shade and other previously cutover areas to secondary forest. Unlike the 1980 survey, no areas were classified as nonstocked in 1990.

Additions to secondary forest land amounted to about 10 000 hectares in the Subtropical Moist Forest and about 10 000 hectares in the Subtropical Wet Forest. Secondary forests accounted for about two-thirds of the Subtropical Moist Forest's timberland area in 1980 but represented more than 90 percent of its timberland in 1990. The largest accretion in the Subtropical Moist Forest, about 15 000 hectares, occurred on shallow volcanic soils. In the Subtropical Wet Forest, the proportion of secondary forests in timberland increased from about one-third to two-thirds. Deep volcanic soils and granitic soils of the Subtropical Wet Forest each amassed about 10 000 hectares.

The areas of both abandoned and active coffee shade forests decreased by about one-half. In 1980, both forest classes combined covered about 59 000 hectares. By 1990, the estimated total area was 30 000 hectares, of which 7 000 hectares were in active coffee shade. These active coffee shade forests fell from 16 percent to 5 percent of all

timberland. The Life Zone soil group stratum with the greatest loss of active coffee shade forest area, more than 10 000 hectares, was the deep volcanic classification within the Subtropical Wet Forest. About one-half of the total reduction in active coffee shade area resulted from the conversion of these lands to alternative agricultural uses; the remainder was simply abandoned.

Many of the older, abandoned coffee shade areas are very difficult to distinguish from reverted pasture and cropland (Birdsey and Weaver 1987). The estimated area of forests classified as abandoned coffee shade decreased from 38 000 to 23 000 hectares between 1980 and 1990. About one-half of this decrease resulted from reclassifying these areas to secondary forest. About one-third of the abandoned coffee shade areas were converted to agricultural uses other than coffee production. The remaining 15 percent were either developed for nonforest land uses such as roads and rights-of-way or returned to active coffee production.

Deep volcanic soils of the Subtropical Wet Forest are the stratum encompassing the main coffee-producing region of the island. Consequently, this stratum had the most extensive coverage of abandoned and active coffee shade areas in both 1980 and 1990. By 1990, however, there were about 5500 fewer hectares of abandoned coffee shade and about 10 500 fewer hectares of active coffee shade than in 1980. Active coffee shade forests were also present, albeit to a lesser degree, on limestone and granitic soils in 1980. In 1990, the 7000 hectares of active coffee shade were confined only to deep soils of volcanic origin, with about 80 percent in the Subtropical Wet Forest and the remainder on the Subtropical Moist Forest.

**Stand Size**

In 1980, timberland was about evenly divided among sawtimber, poletimber, and sapling-seedling stands (fig. 6). By 1990, sawtimber stands occupied nearly 66 000 hectares and dominated Puerto Rico's timberland. These stands covered about 1.4 times the area of poletimber stands and 2.2 times the area of sapling-seedling stands.

The area of sapling-seedling stands on all timberland fell from 33 200 hectares to about 30 544 hectares, despite a 20-percent increase in the area of sapling-seedling stands in secondary forests. This decrease resulted from a large loss of sapling-seedling stands in abandoned and active coffee shade forests. Sapling-seedling stands in both coffee shade forests occupied more than 8 000 hectares in 1980 but only 700 hectares in 1990, all in abandoned coffee shade forest. Moreover, poletimber stands in coffee shade forests decreased from 19 000 to 7 000 hectares. By 1990, most of the poletimber stands were found in abandoned coffee shade forest. In contrast, sawtimber stands in abandoned and active coffee shade forests remained fairly stable, covering about 23 000 hectares in 1980 and 1990. The majority of these sawtimber stands were in abandoned coffee shade forest.

In secondary forests, the area of sawtimber stands almost tripled, and the area of poletimber stands more than doubled between 1980 and 1990. Poletimber and sawtimber stands combined occupied over 80 000 hectares, almost three-fourths of all secondary forest land. This total area was almost evenly divided between these two stand-size classes.

The increased coverage of poletimber and sawtimber stands is supported by comparison of the 1980 and 1990 diameter

distributions (fig. 7). Numbers of live trees in the poletimber diameter classes (15, 20, and 25 centimeters) increased by an average of 37 percent. Numbers of live trees in the sawtimber diameter classes (30 centimeters and larger) increased by an average of 50 percent.

**Species Composition**

Ten species accounted for about one-half of all live basal area both in 1980 and in 1990. Nine of these species were the same in both surveys, while *Erythrina poeppigiana*




Figure 7—Number of live trees by diameter class, Puerto Rico timberland, 1980 and 1990.





Figure 6—Timberland area by stand-size class, Puerto Rico, 1980 and 1990.

placed *Citrus sinensis* (fig. 8). However, their rankings by basal area plurality changed. Two species, *Inga vera* and *Anthodea campanulata*, showed significant changes in their relative contributions to live basal area.

*Anthodea campanulata*, formerly ranked seventh in basal area plurality, replaced *I. vera* as the species with the most basal area. This species tripled its relative proportion of live basal area from 3 percent to 10 percent. The proportion of live basal area represented by *I. vera* decreased by about one-fifth to 5 percent. Furthermore, *I. vera* and *I. fagifolia* were only two species for which basal area decreased.

In 1980, coffee shade forests were less complex than secondary forests, containing fewer than half the number of species found in the latter (Birdsey and Weaver 1982). This same trend in species diversity between coffee shade and secondary forests was apparent in 1990. The 10 most common species in secondary forests accounted for 47 percent of total basal area. In contrast, the 10 most common species in abandoned and active coffee shade forests accounted for 68 and 83 percent, respectively, of their total basal areas.

### Basal Area Distribution

The Subtropical Wet Forest continued to carry a larger proportion of timberland basal area than the Subtropical Moist Forest; however, timberland basal area did become more evenly distributed between the two Life Zones. In 1980, timberland basal area in the Subtropical Wet Forest was almost twice that found in the Subtropical Moist

Forest. By 1990, timberland basal area in the Subtropical Wet Forest was only about 1.5 times greater than in the Subtropical Moist Forest. Total timberland basal area in the Subtropical Moist Forest increased 78 percent to over 650 000 square meters. Growing-stock quality trees accounted for 90 percent of that increase. In addition, average basal area increased from 6 to 9 square meters per hectare in the Subtropical Moist Forest and from 10 to 13 square meters per hectare in the Subtropical Wet Forest.

Total basal area changes in the Subtropical Wet Forest were most noticeable for sawtimber-size trees, which increased by 33 percent, whereas the total basal area of poletimber trees increased by 25 percent. Within the sawtimber tree-size class, basal area of the growing-stock component more than doubled, whereas that of the rough-and-rotten component decreased to almost one-half the amount present in 1980.

In 1990, about 60 percent of the sawtimber-size stands in timberland had basal areas >15 square meters per hectare. This reflects an improvement in the stocking of these stands since 1980 when over 50 percent of these sawtimber stands had basal areas <15 square meters per hectare. Stocking levels in poletimber stands also showed improvement. The area of poletimber stands having basal areas >20 square meters per hectare increased from 10 000 to over 16 000 hectares since 1980.

The total area of sapling-seedling stands decreased by <10 percent, but no sapling-seedling stands had basal areas of >15 square meters per hectare by 1990. In addition, more




Figure 8—Relative species importance by basal area plurality, Puerto Rico timberland, 1990.

than one-half of the 30 000 hectares of sapling-seedling stands had basal areas averaging <5 square meters per hectare.

The average basal area of all live trees in timberland increased from 13.2 to 15.2 square meters per hectare during the decade, with the poletimber and sawtimber-size classes each representing an increase of 1 square meter per hectare (fig. 9). The average basal area of sapling trees remained about the same at 4.3 square meters per hectare.

Of the three forest classes, abandoned coffee shade had the greatest increase in average basal area of all live trees, almost 3 square meters per hectare (fig. 10). An almost twofold increase in the basal area of poletimber trees effected this change (fig. 11). In active coffee shade forests, average basal area of poletimber trees increased by over 2 square meters per hectare. Average basal area of sawtimber-size trees decreased in abandoned coffee shade forests but remained the same in active coffee shade forests. In contrast, sawtimber-size tree basal area increased in secondary forests, whereas poletimber-size tree basal area remained stable.

In 1980, the ratio of growing-stock tree basal area to rough-and-rotten tree basal area for poletimber-size trees was about 1.5 to 1.0; for sawtimber-size trees, about 1 to 1. Since 1980, the basal area of poletimber trees of growing-stock quality more than doubled, and the basal area of sawtimber-size, growing-stock trees almost tripled. As a result, the ratio of growing-stock basal area to rough-and-rotten basal area reached 2 to 1 for poletimber-size trees and 3 to 1 for sawtimber-size trees by 1990.




Figure 9—Average basal area by tree-size class, Puerto Rico timberland, 1980 and 1990.




Figure 10—Average basal area of all live trees by forest class, Puerto Rico timberland, 1980 and 1990.




Figure 11—Average basal area by forest class and tree-size class, Puerto Rico timberland, 1980 and 1990.

### Timber Volume

Average timber volume on all timberland increased from 42 to 80 cubic meters per hectare between 1980 and 1990. Ten species accounted for about 55 percent of all timber volume in both surveys (fig. 12).




Figure 12—Relative species importance by timber volume, Puerto Rico timberland, 1990.

By 1990, the Subtropical Moist and Subtropical Wet forests supported average timber volumes of 59 and 103 cubic meters per hectare, respectively, compared to 31 and 51 cubic meters per hectare in 1980 (fig. 13). Total timber volume became more evenly distributed between the two Life Zones. The timber volume ratio between the Subtropical Moist Forest and the Subtropical Wet Forest was 1.0 to 1.9 in 1980 and 1.0 to 1.6 in 1990.

The Subtropical Moist Forest, with 39 percent of the total timber volume by 1990, had more than doubled to 4.5 million cubic meters during the decade. Growing-stock quality trees, responsible for over 80 percent of this increase, almost tripled in timber volume. Within the growing-stock component, the volume of timber in sawtimber-size trees more than tripled to almost 2 million cubic meters.

The Subtropical Wet Forest, accounting for 61 percent of total timber volume, supported over 7 million cubic meters in 1990, almost two times the volume in 1980. Growing-stock trees, whose timber volume more than doubled to 5.4 million cubic meters, accounted for 90 percent of this increase. Timber volume of growing-stock trees in the Subtropical Wet Forest continued to be almost equally divided between poletimber and sawtimber-size trees.

The total volume of timber in sawtimber-size, growing-stock trees across all timberland increased from 1.8 million to almost 4.9 million cubic meters (fig. 14). Volume accretion in the saw-log portion of these trees contributed to almost one-half the total increase in timber volume. Branch and fork sections accounted for over one-fourth of the volume increase.




Figure 13—Timber volume by Life Zone and tree class, Puerto Rico timberland, 1980 and 1990.

In poletimber trees, volume increment in the bole sections accounted for about one-half the increase in timber volume (fig. 15); branch and fork sections and the sound cull component made up the remainder. Pronounced gains in the volumes of branches and forks (figs. 14, 15) are characteristic of stands in the late poletimber-early sawtimber stages of development.




Figure 14—Distribution of components of timber volume in sawtimber-size, growing-stock trees, Puerto Rico timberland, 1980 and 1990.




Figure 15—Distribution of components of timber volume in poletimber-size, growing-stock trees, Puerto Rico timberland, 1980 and 1990.

Among the Life Zone-soil group strata, both the highest and lowest timber volumes were found in the Subtropical Wet Forest. The highest volume, 4.8 million cubic meters, occurred on deep volcanic soils, and the lowest volume, 40000 cubic meters, occurred on limestone soils. About three-fourths of all timber volume was on slopes >25 percent. More than one-half of this portion occurred on slopes >45 percent.

The proportions of timber volume represented by sawtimber trees, poletimber trees, and rough-and-rotten trees showed similar patterns of change among the three forest classes (fig. 16). In all forest classes, the proportion of timber volume in sawtimber trees increased, whereas the proportion of timber volume in rough-and-rotten trees decreased. The proportion of timber volume in poletimber trees changed little or not at all.

### Sawtimber Volume

Sawtimber volume, the volume in the saw-log portion of sawtimber-size trees of growing-stock quality, doubled from 1.3 to 2.6 million cubic meters across all timberland. Averaging 18 cubic meters per hectare, sawtimber volume accounted for about one-third of growing-stock volume and about one-fourth of timber volume (fig. 17). These sawtimber volume proportions remained relatively unchanged since 1980. Active coffee shade was the only forest class in which average sawtimber volume and average timber volume, increasing from 36 to 48 percent and 20 to 32 percent, respectively (fig. 18). Ten species represented slightly more than two-thirds of the total sawtimber volume in all timberland (fig. 19). Sawtimber volume in secondary forests almost tripled to 1.8 million cubic meters but stayed relatively unchanged in the remaining forest classes (fig. 20). Almost 80 percent of the total sawtimber volume, over 2 million cubic meters, was in




Figure 16—Timber volume by class of timber and forest class, Puerto Rico timberland, 1980 and 1990.




Figure 17—Average sawtimber, growing-stock, and timber volume, Puerto Rico timberland, 1980 and 1990.




Figure 18—Average growing-stock, sawtimber, and timber volume by forest class, Puerto Rico timberland 1980 and 1990.

sawtimber-size stands. Over one-half the total sawtimber volume was present in trees belonging to the 30-, 35-, and 40-centimeter diameter classes. Forty-two percent of the sawtimber volume was present in trees with a butt log grade of 1 or 2. Over 80 percent of that volume was in log grade 2 trees. Only 7 percent of the total sawtimber volume occurred in trees having a butt log grade of 1.

## Trends in Secondary Forests

The area of secondary forests nearly doubled during the decade to over 113 000 hectares. The net change in the basal area of all live trees was minor, increasing from 13.8 to 15.1 square meters per hectare. Average basal area of growing-stock trees increased from 5.0 to 7.6 square meters per hectare, primarily due to the growth of sawtimber trees. Basal area of the rough-and-rotten component fell from 3.4 to 2.8 square meters per hectare. The average combined basal area of all poletimber and sawtimber trees in secondary forests therefore rose from 8.3 to 10.4 square meters per hectare. The other component of all live basal area, sapling trees, decreased from 5.5 to 4.7 square meters per hectare.

With a total timber volume of 8.6 million cubic meters by 1990, secondary forests accounted for about three-fourths of the total timber volume of Puerto Rico's commercial region. Over three-fourths of that timber volume was found in growing-stock trees (fig. 21). Less than one-fourth of the timber volume in secondary forests was in rough trees, whereas rotten trees comprised only 1 percent. Growing-stock volume was 5.7 million cubic meters, about two-thirds of the total timber volume in secondary forests. That proportion was nearly the same in 1980; however, growing-stock volume was more evenly distributed between the poletimber and sawtimber-size classes in 1990. The remaining one-tenth of total timber volume found in growing-stock trees was sound cull volume. Sound cull amounted to 14 percent of the total timber volume present in growing-stock trees only. The total sawtimber volume of 2.6 million cubic meters reached by 1990 was comparable to the 1980 volume of all sound wood (timber volume) of sawtimber trees in all timberland.

Average timber volume increased from 44 to 75 cubic meters per hectare, and average growing-stock volume increased from 29 to 50 cubic meters per hectare in secondary forests. Sawtimber volume on the average hectare in secondary forests rose from 11 to 16 cubic meters, but its relative proportions of growing-stock volume and timber volume fell slightly to 32 percent and 22 percent, respectively. These changes in volume component proportions described for secondary forests closely typify the trends observed in timberland as a whole.

Just over one-third of the secondary forest stands supported 50 or more cubic meters of timber volume per hectare, up slightly from 30 percent in 1980. Stands with between 50 and 75 cubic meters per hectare covered over 15 000 hectares, four times the area covered in 1980. The

### Species name




Figure 19—Relative species importance by sawtimber volume,<sup>1</sup> Puerto Rico timberland, 1990.




Figure 20—Total growing-stock, sawtimber, and timber volume by forest class, Puerto Rico timberland, 1980 and 1990.

proportion of all secondary forest stands represented by this volume class increased from 6 to 13 percent. The coverage of stands having 100 or more cubic meters of timber volume per hectare increased from 8000 to over 13000 hectares, although their proportion of all secondary forest stands dropped slightly from 14 to 12 percent. Additions to these two stand-volume classes increased the average timber volume of secondary forests from 44 to 75 cubic meters per hectare.

The character of secondary forests improved during the decade as shown by the greater coverage of stands composed of higher quality trees. Poletimber and sawtimber stands of secondary forests rose from 57 to 74 percent of the total area by 1990. Moreover, the poletimber and sawtimber trees of growing-stock quality in these stands accounted for almost three-fourths of the total basal area and just over three-fourths of the total volume compared to 60 and 68 percent, respectively, in 1980. Such changes indicate that stands in secondary forest stands are experiencing good growth and development, particularly those containing the higher quality trees.

### Trends in Coffee Shade Forests

**Abandoned coffee shade forest**—Stands formerly managed as coffee shade covered an estimated 15 000 fewer hectares in 1990 than in 1980 and supported stands with basal areas ranging from 5 to 30 square meters per hectare. Average basal area of all live trees in these forested areas increased from 14 to 17 square meters per hectare. The only observed decrease in basal area was for sawtimber-size trees, which fell from 6.0 to 4.7 square meters per hectare. The basal area of poletimber and sapling trees increased from 4.9 to 8.7 and 3.0 to 3.5 square meters per hectare, respectively.

Fifty-five percent of the abandoned coffee shade area was composed of stands with an average volume of at least 50 cubic meters per hectare. Only about 700 hectares, however, supported stands with average volumes exceeding 100 cubic meters per hectare. Sawtimber stands




Figure 21—Distribution of components of timber volume (thousand cubic meters) in secondary forests, Puerto Rico timberland, 1990.

carrying 25 to 100 cubic meters per hectare occupied 3 percent of the abandoned coffee shade area. Poletimber stands, covering 90 percent of the remaining abandoned coffee shade area, generally supported <25 cubic meters per hectare.

In both surveys, abandoned coffee shade forests had the highest average volume of timber. The average volumes of growing stock and sawtimber in 1980 were very nearly the same as those found in secondary forests; but by 1990, these too had climbed well above those of secondary forests. Average sawtimber volume in abandoned coffee shade forests reached 26 cubic meters per hectare, 10 cubic meters per hectare more than in secondary forests. Abandoned coffee shade forest in 1990 only had about 350 000 cubic meters more timber volume than in 1980. However, because of the lower area estimate in 1990, average timber volume of abandoned coffee shade forests almost doubled from 53 to 102 cubic meters per hectare. These changes effected a shift in the ratio of growing-stock volume to timber volume from one-half to two-thirds. The ratio of sawtimber to timber volume remained stable at about one-fourth in this forest class.

**Active coffee shade forest**—With an estimated 7000 hectares, active coffee shade forest covered about one-third of the area that it occupied in 1980 and contained the lowest total basal areas of any forest classification. Average basal area of all live trees in these forested areas increased from 10 to 12 square meters per hectare. Total sawtimber basal area decreased to about one-third of the 1980 estimate, closely paralleling the reduction in area of active coffee shade forest. The average basal area per hectare of sawtimber trees, however, remained unchanged from 1980 at 3.5 square meters per hectare. Excluding

sapling trees, the total basal area of active coffee shade forest in 1990 averaged 10.2 square meters per hectare, with poletimber and sawtimber trees at 6.7 and 3.5 square meters per hectare, respectively. The average basal area for all size classes in 1980 was also 10.2 square meters per hectare, but this average included saplings. No stands in active coffee shade areas had basal area <5 square meters per hectare or >20 square meters per hectare in 1990.

Active coffee shade forest, as the smallest forest classification sampled, had the lowest total volume of live trees. Sixty percent of the 7000 hectares of active coffee shade forest supported stands having 25 to 50 cubic meters per hectare. Stands comprising the remaining 2800 hectares of active coffee shade forest had volumes between 50 and 100 cubic meters per hectare.

Although active coffee shade supported the lowest total volume, the average timber and growing-stock volumes per hectare in this forest class were very comparable to those found in secondary forests (fig. 18). The total timber volume found in active coffee shade in 1990 was about 574 000 cubic meters. The volume decrease of about 200 000 cubic meters since 1980 did not proportionately parallel the more significant areal decrease in active coffee shade forest. Consequently, the average timber volume of active coffee shade forests rose substantially from 36 to 82 cubic meters per hectare. Growing-stock volume comprised about two-thirds of that timber volume. Sawtimber volume per hectare more than tripled to 26 cubic meters and accounted for about one-half the average hectare's volume of growing stock. Sawtimber volume, formerly one-fifth of timber volume, represented about one-third of total timber volume on the average hectare in 1990.

## Conclusions

Puerto Rico's forest resources are recovering from the land use practices prevalent in the 19th and first half of the 20th centuries. The 1990 survey revealed an increase in forest land, better stocking with larger and higher quality timber trees, and a maturing secondary forest. Area and volume trends, such as the decreased area of sapling-seedling stands and the substantial increase in sawtimber volume, are primarily the result of the absence of management activity and commercial timber harvesting on the island.

The results of the 1990 survey portray Puerto Rico as an example of how once heavily degraded tropical forests can recover to a level of productivity and usefulness.

Knowledge of these trends can facilitate the development of management techniques to enhance the value of the timber resource and provide other forest-related benefits for the residents of Puerto Rico.

## Literature Cited

- Birdsey, Richard A.; Weaver, Peter L.** 1982. The forest resources of Puerto Rico. Resour. Bull. SO-85. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 59 p.
- Birdsey, Richard A.; Weaver, Peter L.** 1987. Forest area trends in Puerto Rico. Res. Note SO-331. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 5 p.
- Durland, W.D.** 1929. Forest regeneration in Puerto Rico. *Economic Geography*. 5: 369–381.
- Ewel, J.J.; Whitmore, J.L.** 1973. The ecological life zones of Puerto Rico and the U.S. Virgin Islands. Res. Pap. ITF-18. Río Piedras, PR: Institute of Tropical Forestry. 72 p.
- Gill, Tom.** 1931. Tropical forests of the Caribbean. Baltimore: Tropical Plant Research Foundation. 318 p.
- Holdridge, L.R.** 1967. Life zone ecology. Revised. San José, Costa Rica: Tropical Science Center. 206 p.
- Husch, Bertram; Miller, Charles I.; Beers, Thomas W.** 1972. Forest mensuration. New York: Ronald Press Co. 410 p.
- Koenig, N.** 1953. A comprehensive agricultural program for Puerto Rico. Washington, DC: U.S. Department of Agriculture and Commonwealth of Puerto Rico. 290 p.
- Little, Elbert L., Jr.; Wadsworth, Frank H.** 1964. Common trees of Puerto Rico and the Virgin Islands. Agric. Handb. 249. Washington, DC: U.S. Department of Agriculture, Forest Service. 548 p.
- Little, Elbert L., Jr.; Woodbury, R.O.; Wadsworth, Frank H.** 1974. Trees of Puerto Rico and the Virgin Islands. Agric. Handb. 449. Washington, DC: U.S. Department of Agriculture, Forest Service. 1024 p. Vol. 2.
- Wadsworth, F.H.** 1950. Notes on the climax forests of Puerto Rico and their destruction and conservation prior to 1900. *Caribbean Forester*. 11(1): 38–47.
- Zambrana, J.A.** 1978. Asociaciones de suelos de Puerto Rico y su potencial para la siembra de árboles. Río Piedras, PR: Dept. Agric. de EEUU, Servicio Forestal, Area Sureste, División de Dasonomía. 6 p.

## Appendix

### Survey Methods

#### 1980 Survey

The 1980 survey of Puerto Rico's forest resources was designed to evaluate the production potential of the island's secondary timber resources and to describe their ecological characteristics (Birdsey and Weaver 1982). The 1980 survey also provided baseline information from which changes in forest areas, species compositions, and growth could be assessed in subsequent inventories.

The purpose of the sample design was to partition the island into two major areas: (1) the area with potential for commercial production of timber where an intensive inventory would be conducted, and (2) noncommercial areas where limiting physiographic or economic factors would preclude the commercial production of timber. The noncommercial area encompassed lands designated as critical watersheds, areas with adverse site conditions such as steep slopes or high rainfalls, and areas with higher priority uses such as urban expansion or agriculture. Noncommercial areas were delineated on a 1:120,000 scale base map but excluded from the ground sample. The remainder of the island, the commercial timber production area, is concentrated in the central highlands. This area was stratified by Life Zone (Subtropical Moist and Subtropical Wet Forests) and soils (steep volcanics, shallow volcanics, granitic, and limestone).

Forest resource statistics were obtained using a systematic sampling method involving forest-nonforest classification of temporary points on aerial photographs and on-the-ground measurements of trees on permanent plots. The sample locations were at the intersections of a 3- by 5-kilometer grid design. There were 978 of these permanent plots on the entire island and 437 in the commercial region. Ground sample locations were first selected on U.S. Geological Survey maps, then transferred to aerial photographs. Initial forest area estimates were derived from interpretation of 10,925 photopoints on black-and-white aerial photographs from the mid-1970's. The photo-interpretation estimates were adjusted based on ground verification at the 437 permanent ground plots. Aerial photographs were also obtained for most of the noncommercial region. Forest cover for the remaining land was estimated using reference material.

Each ground sample location consisted of a cluster of three variable radius subplots spaced 25 meters apart and running along a north-south line. The center of each sample location was the center of the middle subplot in the cluster. This point

was established on the ground by running a computed azimuth and distance from a selected starting point. Each sample tree on the variable radius subplot represented 2.5 square meters of basal area per hectare, or 0.83 square meters of basal area per hectare when the subplots were clustered.

Each sample location was classified according to several site characteristics, including the slope and aspect at each of the three subplots. Sample trees smaller than 12.5 centimeters in diameter at breast height (d.b.h.) were tallied on fixed radius plots of approximately 40 square meters around each of the three subplot centers. Sample trees 12.5 centimeters and larger were measured to estimate volume and assessed to determine wood quality.

Volumes were derived from detailed measurements of trees at all sample locations. These measurements included d.b.h., total height, bole length, log length, and two upper stem diameters. Volume of the bole and saw-log portions were computed using the International 1/4-inch rule. Regression equations were developed for five species groups and used to estimate branch and fork volumes in the sections above the bole.

#### 1985 Update

The primary objective of the 1985 "midcycle" update (Birdsey and Weaver 1987) was to refine the procedures developed for the 1980 survey in preparation for the 1990 resurvey. Complete color infrared aerial photographic transparencies at a scale of 1:64,000 were acquired for the Commonwealth in 1984 to compensate for the lack of complete coverage in 1980. The increased spectral resolution of the aerial photography allowed for the development of a more detailed forest cover type assessment.

A dot-count method was used to identify 12 forest and 3 nonforest land cover classes on the film transparencies. A 5- by 5-dot grid was centered over each of the 978 permanent sample plot locations. Land-cover class was estimated for each plot location and for each dot, resulting in a total of 22,418 photographic interpretations. Most forested locations were identified by 8 of the 12 possible classes. The 3- by 3-kilometer grid sample did not include montane (tabonuco) forest types or plantations also present in Puerto Rico.

#### 1990 Resurvey

All plots in the commercial forest region were remeasured in the spring of 1990. The expanded forest classes used in the 1985 midcycle were used. Only four of these classes were found within the commercial region (table 1) because that area by definition excluded specific forest types such as mangroves and cloud forests found on steep, wet slopes.

Black-and-white 1988-89 aerial photographs at a scale of 1:20,000 were purchased, and a 5- by 5-dot grid was interpreted for each photographic location. There was a total of 10,545 photointerpretations and 430 field-verified sample plot interpretations.

All trees were remeasured using the same methodology established in 1980. Three subplots were measured using variable radius sampling for timber evaluation and fixed radius for regeneration. The basal area factor for the variable radius was 2.5 square meters or 0.83 square meters of basal area per hectare when the subplots were clustered. The minimum d.b.h. for trees on the variable radius plot was 12.5 centimeters.

Regeneration was assessed by tallying all trees between 2.5 and 12.4 centimeters within a 3.6-meter fixed radius plot (about 40 square meters per subplot). Tree variables included form and damage factors that would affect merchantability of timber. Trees were assigned log grades and estimates for both total cull and log cull. Volume computation procedures duplicated those used in 1980.

## Reliability of the Data

Two types of errors affect the reliability of the estimates. The first source of error—estimating error—derives from mistakes in measurement, judgment, recording or compiling, and from limitations of the equipment. The SOFIA minimizes estimating error through comprehensive training, supervision, quality-control programs, and emphasis on precision. The second type of error—sampling error—arises from the use of a sample to estimate population parameters and from variability of the items being sampled. This deviation is susceptible to a mathematical evaluation of the probability of error.

The second survey of Puerto Rico again required a special team approach to meet fieldwork requirements for accuracy. In addition to the logistical problems associated with adapting field sampling techniques to a tropical environment, expertise in Puerto Rican dendrology was needed. The approach teamed the most experienced timber cruisers from the SOFIA with dendrologists and botanists from Puerto Rico.

Statistical analysis of the data indicates a sampling error of plus or minus 1.7 percent for the estimate of total timberland and 8.0 percent for the timber-volume estimate. In regular State surveys, sampling errors are estimated for both area and volume using the random sampling formula. The sampling error therefore normally increases as the area

or volume under consideration is subdivided according to various strata. Sampling errors for estimates of volume in the Puerto Rico survey are consistent with this approach. A different methodology was used to derive sampling errors for area estimates. Area sampling for the Puerto Rico survey involved pre-stratification of the commercial region into various forest classes, and the proportion represented by each forest class was estimated based on a binomial classification of the photoplots. Because of the different type of area stratification and the assumption of a binomial distribution, sampling errors for areas were estimated by a different method and, thus, do not follow the usual pattern. These errors were estimated using the procedure described by Husch and others (1972) for calculating the variance of adjusted proportions of stratified stand classes.

The following tabulation shows the sampling errors to which the area and volume estimates for secondary forest, abandoned and active coffee shade, and all timberland are liable:

Stratum	Area	Sampling error <sup>1</sup>	Volume	Sampling error <sup>2,3</sup>
	<i>Hectares</i>	<i>Percent</i>	<i>Thousand cubic meters</i>	<i>Percent</i>
Secondary forest	113 558	4.0	8 570.1	10.5
Abandoned coffee shade forest	23 365	3.5	2 380.2	7.8
Active coffee shade forest	7 010	2.4	574.2	21.1
All forest land	143 933	1.7	11 524.6	8.0

## Definitions

### Forest Land Classes

**Abandoned coffee shade.** Secondary forest land resulting from the abandonment of coffee production under shade trees.

**Coffee shade.** A multistory, multicrop system used principally for the production of coffee. An upper story of shade trees is characteristic.

<sup>1</sup>By variance formula for adjusted proportions of a binomially distributed random variable.

<sup>2</sup>By random sampling formula.

<sup>3</sup>The sampling errors for volume are equal to one standard deviation for the sample data.

**Commercial survey region.** Area designated as having potential for commercial timber production and where field sampling was carried out. This area remained after the delineation and exclusion of areas with higher priority uses (urban areas, better agricultural land, reserved areas), dry coastal areas, areas with adverse site conditions (extreme slope, excessive rainfall, etc.), and critical watershed areas.

**Forest land.** Land at least 10 percent stocked<sup>4</sup> by forest trees of any size, or formerly having had such tree cover and not currently developed for nonforest use. The minimum area for classification of forest land is one-half hectare, and the minimum width for forest strips is 35 meters. Unimproved roads and trails, streams, and clearings in forest areas are classed as forest if less than 35 meters in width.

**Noncommercial forest land.** Forest land incapable of yielding crops of industrial wood because of adverse site conditions, forest land withdrawn from timber utilization through statute or administrative regulation, or forest land with higher priority use (except coffee shade).

**Nonstocked land.** Commercial forest land less than 10 percent stocked with growing-stock trees. This includes areas covered by inhibiting vegetation (brush, vines, ferns, etc.) classed as forest land.

**Secondary forest land.** Forest land resulting from the abandonment of cropland or pasture, forest resulting from the regeneration of previously cutover or disturbed forest land, and reclassification of both types of coffee shade.

**Timberland.** Forest land that is producing or is capable of producing crops of industrial wood and not withdrawn from timber utilization. Forest lands with higher priority uses, but not specifically withdrawn from timber utilization, are excluded from this class of forest land. These excluded lands were delineated on a base map. Coffee shade is included in this category.

#### Noncommercial Forest Strata

**Alluvial regions.** River floodplains with agricultural land use.

**Critical watersheds.** Upland areas with an average slope >60 percent or rainfall >2500 millimeters per year. These areas require a continuous protective forest cover. Some timber removal would be feasible; however, much of the area is not highly productive.

**Mangrove and swamp.** Coastal wetlands with unique characteristics and values requiring special management considerations.

**Metropolitan regions.** Regions with primarily urban or residential use.

**Subtropical Dry Forest.** An ecological Life Zone delineated by a mean annual rainfall of between 600 millimeters and 1000 or 1100 millimeters, and a mean annual biotemperature between about 18 and 24 °C (Holdridge 1967).

**Unproductive soils.** Soils incapable of yielding crops of industrial wood.

#### Stand-size Classes

**Poletimber stands.** Stands with  $\geq 5$  square meters per hectare of basal area in sawtimber or poletimber-size trees, and with poletimber basal area exceeding that of sawtimber basal area.

**Sapling-seedling stands.** Stands with  $\geq 5$  square meters per hectare of basal area, with more than one-half of this basal area in saplings or seedlings, or stands with  $< 5$  square meters per hectare of basal area, regardless of the size of existing trees.

**Sawtimber stands.** Stands with  $\geq 5$  square meters per hectare of basal area in sawtimber or poletimber-size trees, and with sawtimber basal area at least equal to poletimber basal area.

#### Timber Classes

**Acceptable trees.** Trees meeting the specifications for growing stock but not qualifying as desirable trees.

**Desirable trees.** Growing-stock trees that have no serious defects to limit present or prospective use, are of relatively high vigor, and contain no pathogens that may result in death or serious deterioration before rotation age. These trees would be favored in silvicultural operations.

**Growing-stock trees.** Sawtimber trees, poletimber trees, saplings and seedlings; that is, all live trees except rough-and-rotten trees, regardless of species.

**Poletimber trees.** Growing-stock trees 12.5 to 22.5 centimeters in d.b.h. for softwoods and 12.5 to 27.5 centimeters for hardwoods and of good form and vigor.

<sup>4</sup>Ten percent of "normal" stocking as defined for the Midsouth Forest Survey.

**Rough-and-rotten trees.** Live trees that are unmerchantable for saw logs now or prospectively because of defect or rot.

**Salvable dead trees.** Standing or down dead trees that are currently or potentially merchantable.

**Saplings.** Growing-stock trees 2.5 to 12.5 centimeters in d.b.h. and of good form and vigor.

**Sawtimber trees.** Growing-stock trees 22.5 centimeters and larger in d.b.h. for softwoods and 27.5 centimeters and larger for hardwoods, and containing at least one 3.5-meter saw log.

### Timberland Strata

**Deep volcanic soils.** Deep soils of the wet volcanic uplands with clayey texture, typically red and acid (Zambrana 1978). The soil depth is generally >50 centimeters.

**Granitic soils.** Shallow soils of the wet uplands with granitic origin and with a typically sand-clay loam texture (Zambrana 1978).

**Limestone soils.** Shallow soils over limestone found in the moist coastal hills (Zambrana 1978).

**Shallow volcanic soils.** Shallow soils of the wet volcanic uplands with clayey texture (Zambrana 1978). The soil depth is generally <50 centimeters.

**Subtropical Moist Forest.** Forest occurring in an ecological Life Zone delineated by a mean annual rainfall of about 1000 to 2000 millimeters and a mean annual biotemperature between about 18 and 24 °C (Holdridge 1967).

**Subtropical Wet Forest.** Forest occurring in an ecological Life Zone delineated by a mean annual rainfall of about 2000 to 4000 millimeters and a mean annual biotemperature between about 18 and 24 °C (Holdridge 1967).

### Volumes

**Volume of growing stock.** Volume of sound wood (less cull volume) in the bole and branches of sawtimber and poletimber trees from stump to a minimum 10-centimeter diameter outside bark or to the point past which a 1-meter section meeting minimum qualifications can no longer be measured because of limbs or other cull.

**Volume of sawtimber.** Net volume of the saw log portion of sawtimber trees in cubic meters, calculated according to the International 1/4-inch rule.

**Volume of timber.** Volume of all sound wood (including sound cull) in the bole and branches of growing stock, rough, rotten, and salvable dead trees 12.5 centimeters and larger in d.b.h. from stump to a minimum 10-centimeter diameter outside bark. The minimum length of any section included is 1 meter.

### Miscellaneous Definitions

**Basal area.** The area in square meters of the cross section at breast height of a single tree or of all the trees in a stand, expressed as square meters per hectare.

**D.b.h.** Tree diameter in centimeters, outside bark, measured at 1.3 meters aboveground.

**Diameter class.** Each 5-centimeter diameter class extends from 2.5 centimeters below to 2.4 centimeters above the stated midpoint. For example, the 30-centimeter diameter class includes trees 27.5 centimeters through 32.4 centimeters in d.b.h.

**Log grades.** A classification of logs based on external characteristics as indicators of quality or value. Logs are graded according to standards used for hardwoods in the Southern United States. Grade 1 logs have very little sweep, crook, cull, or other defects, are the largest, and are suitable for standard lumber. Grade 4 logs need only be 20 centimeters in diameter outside bark at the small end, and are used only for ties and timbers. Grades 2 and 3 are between the two extremes.

## CONVERSION FACTORS

### Metric to English and English to Metric conversions

centimeter = 0.3937 inch  
meter = 3.281 feet  
kilometer = 0.6214 mile  
square meter = 10.7639 square feet  
square kilometer = 0.3861 square mile  
hectare = 2.471 acres  
cubic meter = 35.3145 cubic feet  
square meter per hectare = 4.356 square feet per acre  
cubic meter per hectare = 14.29 cubic feet per acre

1 inch = 2.54 centimeters  
1 foot = 0.3048 meter  
1 mile = 1.6093 kilometers  
1 square foot = 0.0929 square meter  
1 square mile = 2.590 square kilometers  
1 acre = 0.4047 hectare  
1 cubic foot = 0.0283 cubic meter  
1 square foot per acre = 0.2296 square meter per hectare  
1 cubic foot per acre = 0.07 cubic meter per hectare

### Metric inventory standards used in the 1990 survey

#### Item

Trism size  
Grid spacing  
Cluster point spacing  
Fixed plot size  
Crest height  
Stump height  
Diameter classes

#### Metric standard

BAF 2.5  
3 kilometers  
25 meters  
40 square meters ( $r = 3.6$  meters)  
15 square meters ( $r = 2.2$  meters)  
1.3 meters  
30 centimeters  
5-centimeter = 2.5 to 7.5 centimeters d.b.h.,  
10-centimeter = 7.5 to 12.5 centimeters d.b.h.,  
15-centimeter = 12.5 to 17.5 centimeters d.b.h.,  
20-centimeter = 17.5 to 22.5 centimeters d.b.h., etc.

#### Tree-size classes

Sapling  
Poletimber (hardwood)  
Sawtimber (hardwood)  
Sawtimber (softwood)

2.5 to 12.5 centimeters d.b.h.  
12.5 to 27.5 centimeters d.b.h.  
27.5+ centimeters d.b.h.  
22.5+ centimeters d.b.h.

#### Minimum top diameter, outside bark

Cubic volume  
Hardwood saw log  
Softwood saw log  
Sapling

10 centimeters  
22.5 centimeters  
17.5 centimeters  
2.5 centimeters

#### Minimum saw-log diameter, inside bark

Hardwood  
Softwood

20 centimeters  
15 centimeters

#### Minimum length

Cubic section  
Saw log  
Sawtimber tree

1 meter  
2.5 meters  
3.5-meter saw log

List of Tree Species Included in Standard Tables

Code number	Species name	Common name
1	<i>Cyathea arborea</i> (L.) J. E. Smith	Helecho gigante, tree-fern
7	<i>Euterpe globosa</i> Gaertn. <sup>a</sup>	Palma de sierra, sierra palm
9	<i>Roystonea borinquena</i> O. F. Cook	Palma real, royal palm, Puerto Rico
17	<i>Artocarpus altilis</i> (Parkinson) Fosberg	Panapén, pana de pepitas, breadfruit
19	<i>Castilla elastica</i> Cervantes	Caucho, Central American rubber, castilla
20	<i>Cecropia peltata</i> L.	Yagrumo hembra, trumpet-tree
22	<i>Ficus laevigata</i> Vahl <sup>b</sup>	Jagüey blanco, shortleaf fig
33	<i>Torrubia fragrans</i> (Dum.-Cours.) Standley	Corcho, black mampoo
34	<i>Magnolia portoricensis</i> Bello	Jagüilla
48	<i>Ocotea leucoxydon</i> (Sw.) Mez	Laurel geo
51	<i>Persea americana</i> Mill.	Aguacate, avocado
51	<i>Albizia procera</i> (Roxb.) Benth.	Albizia, tall albizia
52	<i>Inga fagifolia</i> (L.)	Guamá, "sweetpea"
53	<i>Inga quaternata</i> Poepp. & Endl.	Guamá venezolano
54	<i>Inga vera</i> Willd.	Guaba
54	<i>Cassia siamea</i> Lam.	Casia de Siam, Siamese cassia
56	<i>Hymenaea courbaril</i> L.	Algarrobo, West-Indian-locust, courbaril
51	<i>Andira inermis</i> (W. Wright) H.B.K. <sup>c</sup>	Moca, cabbage angelin
54	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	Bucayo gigante, mountain immortelle
56	<i>Lonchocarpus latifolius</i> (Willd.) H.B.K.	Retama
57	<i>Ormosia krugii</i> Urban	Palo de matos
59	<i>Citrus paradisi</i> Macfadyen	Toronja, grapefruit
60	<i>Citrus sinensis</i> Osbeck	China, sweet orange
62	<i>Zanthoxylum martinicense</i> (Lam.) DC.	Espino rubial, white-prickle, Martinique
65	<i>Bursera simaruba</i> (L.) Sarg.	Almácigo, turpentine-tree, gumbo-limbo
66	<i>Dacryodes excelsea</i> Vahl	Tabonuco
67	<i>Tetragastris balsamifera</i> (Sw.) Kuntze	Masa
68	<i>Cedrela odorata</i> L.	Cedro hembra, Spanish-cedar
69	<i>Guarea guidonia</i> (L.) Sleumer	Guaraguao, American muskwood
74	<i>Trichilia pallida</i> (Sw.)	Gaeta
75	<i>Byrsonima coriacea</i> (Sw.) DC.	Maricao
78	<i>Alchornea latifolia</i> Sw.	Achiotillo
75	<i>Hura crepitans</i> L.	Molinillo, sandbox, hura
79	<i>Sapium laurocerasus</i> Desf.	Tabaiba
81	<i>Mangifera indica</i> L.	Mango
83	<i>Spondias dulcis</i> Parkinson	Jobo de la India, ambarella
84	<i>Spondias mombin</i> L.	Jobo, hogplum, yellow mombin
87	<i>Turpinia paniculata</i> Vent.	Sauco cimarrón
88	<i>Cupania americana</i> L.	Guara
93	<i>Thouinia striata</i> Radlk.	Ceboruquillo
94	<i>Meliosma herbertii</i> Rolfe	Aguacatillo
99	<i>Sloanea berteriana</i> Choisy	Motillo
101	<i>Montezuma speciosissima</i> Sessé & Moc.	Maga
104	<i>Ochroma pyramidale</i> (Cav.) Urban	Guano, balsa
105	<i>Quararibæa turbinata</i> (Sw.) Poir	Garrocho
106	<i>Guazuma ulmifolia</i> Lam.	Guácima, jacocalalu
101	<i>Calophyllum brasiliense</i> Camb.	María, santa-maria
103	<i>Clusia rosea</i> Jacq.	Cupey, wild-mammeey, copey clusia
104	<i>Mammea americana</i> L.	Mamey, mammeey-apple

List of Tree Species Included in Standard Tables (continued)

Code number	Species name	Common name
69	<i>Casearia arborea</i> (L. C. Rich.) Urban	Rabo ratón
70	<i>Casearia decandra</i> Jacq.	Tostado, wild honey-tree
71	<i>Casearia guianensis</i> (Aubl.) Urban	Palo blanco, wild-coffee
72	<i>Casearia sylvestris</i> Sw.	Cafeillo
80	<i>Buchenavia capitata</i> (Vahl) Eichl.	Granadillo
81	<i>Bucida buceras</i> L.	Ucar, gregre, oxhorn bucida
84	<i>Terminalia catappa</i> L.	Almendra, Indian-almond
87	<i>Eugenia aeruginea</i> DC.	Guasábara
88	<i>Eugenia jambos</i> L.	Pomarrosa, Rose-apple
92	<i>Myrcia deflexa</i> (Poir.) DC.	Cieneguillo
93	<i>Myrcia splendens</i> (Sw.) DC.	Hoja menuda
95	<i>Psidium guajava</i> L.	Guayaba, common guava
98	<i>Miconia prasina</i> (Sw.) DC.	Camasey
99	<i>Tetrazygia elaeagnoides</i> (Sw.) DC.	Verdiseco
00	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	Pollo
01	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	Yagrumo macho, matchwood
03	<i>Rapanea ferruginea</i> (Ruiz & Pav.) Mez	Mantequero
08	<i>Dipholis salicifolia</i> (L.) A. DC.	Sanguinaria, wild mespel, willow bustic
09	<i>Manilkara bidentata</i> (A. DC.) Chev.	Ausubo, balata
11	<i>Micropholis chrysophylloides</i> Pierre	Caimitillo
12	<i>Micropholis garciniaefolia</i> Pierre	Caimitillo verde
13	<i>Pouteria multiflora</i> (A. DC.) Eyma	Jácana
21	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	Capá prieto, capa
24	<i>Cordia sulcata</i> DC.	Moral, white manjack
27	<i>Citharexylum fruticosum</i> L.	Péndula, pasture fiddlewood, Florida
28	<i>Petitia domingensis</i> Jacq.	Capá blanco
30	<i>Vitex divaricata</i> Sw.	Higüerillo, white fiddlewood
34	<i>Spathodea campanulata</i> Beauv.	Tulipán africano, African tuliptree
36	<i>Tabebuia heterophylla</i> (DC.) Britton	Roble blanco, "white-cedar"
40	<i>Coffea arabica</i> L.	Café, coffee
49	<i>Terebraria resinosa</i> (Vahl) Sprague	Aquilón
60	<i>Pinus caribaea</i> Morelet	Pino hodureño, Caribbean pine
79	<i>Piper amalago</i> L.	Higuillo de limón
94	<i>Ficus stahlii</i> Warb. (E)	Jagüey
30	<i>Nectandra antillana</i> Meisn.	Aguacatillo
34	<i>Nectandra sintenisii</i> Mez	Laurel amarillo
40	<i>Phoebe elongata</i> (Vahl) Nees	Laurel avispollo
54	<i>Calyptanthus pallens</i> Griseb.	Tapón blanco, pale lidflower
92	<i>Clidemia umbrosa</i> (Sw.) Cogn.	Camasey
12	<i>Miconia subcorymbosa</i> Britton	Camasey
21	<i>Coffea dewevrei</i> Wildem. & T. Dur.	Café excelsa, dewevre coffee

Referred to as *Prestoea montana* (R. Grah.) Nichols. In 1980 survey.

Referred to as *Fiscus citrifolia* Mill. In 1980 survey.

Referred to as *Andira inermis* (W. Wright) DC. In 1980 survey.

## **Index of Tables**

### **Forest Area**

1. Land area by detailed land cover class and survey region, Puerto Rico, 1990
2. Land area by survey region and broad land cover class, Puerto Rico, 1980-1990
3. Area of timberland by Life Zone, soil group, and forest class, Puerto Rico, 1990
4. Area of timberland by stand-volume class and forest class, Puerto Rico, 1990
5. Area of timberland by stand-volume class and soil group, Puerto Rico, 1990
6. Area of timberland by stand-size class and forest class, Puerto Rico, 1990
7. Area of timberland by stand-size class and soil group, Puerto Rico, 1990
8. Area of timberland by stand-age class and forest class, Puerto Rico, 1990
9. Area of timberland by stand-age class and soil group, Puerto Rico, 1990
10. Area of timberland by basal area class and forest class, Puerto Rico, 1990
11. Area of timberland by basal area class and stand-size class, Puerto Rico, 1990
12. Area of stocked timberland by Life Zone, soil group, and slope class, Puerto Rico, 1990

### **Species Composition**

13. Number of live trees by species and diameter class, ranked by basal area plurality, Puerto Rico, 1990
14. Number of growing-stock trees by species and diameter class, ranked by basal area plurality, Puerto Rico, 1990

15. Number of live trees on timberland by diameter class and forest class, Puerto Rico, 1990
- 16a. Basal area and number of live trees per hectare of timberland for selected species, secondary forests, Puerto Rico, 1990
- 16b. Basal area and number of live trees per hectare of timberland for selected species, abandoned coffee shade, Puerto Rico, 1990
- 16c. Basal area and number of live trees per hectare of timberland for selected species, active coffee shade, Puerto Rico, 1990
17. Basal area of poletimber and sawtimber trees by tree class and forest class, Puerto Rico, 1990
18. Basal area of poletimber and sawtimber trees by tree class, Life Zone, and soil group, Puerto Rico, 1990

### **Timber Volume**

19. Volume of timber on timberland by class of timber and forest class, Puerto Rico, 1990
20. Volume of timber on timberland by Life Zone, soil group, and tree class, Puerto Rico, 1990
21. Volume of timber on timberland by species and diameter class, Puerto Rico, 1990
22. Volume of growing stock on timberland by species and diameter class, Puerto Rico, 1990
23. Volume of sawtimber on timberland by species and diameter class, Puerto Rico, 1990
24. Volume of sawtimber on timberland by species and butt log grade, Puerto Rico, 1990
25. Volume of timber, growing stock, and sawtimber on timberland by stand-size class, Puerto Rico, 1990
26. Volume of timber on timberland by Life Zone, soil group, and slope class, Puerto Rico, 1990

**Table 1—Land area by detailed land cover class and survey region, Puerto Rico, 1990**

Land cover class	Commercial survey region	Noncommercial survey region <sup>a</sup>	All regions
	----- <i>Hectares</i> -----		
<b>Forest</b>			
Secondary	113 600	83 900	197 500
Abandoned coffee shade	23 400	16 300	9 700
Active coffee shade	7 000	6 800	13 800
Upper mountain	--	2 300	2 300
Palm	--	1 800	1 800
Dwarf	--	1 800	1 800
Xeric scrub <sup>b</sup>	4 200	21 700	25 900
Mangrove	--	4 700	4 700
<b>Total</b>	<b>148 100</b>	<b>139 300</b>	<b>287 400</b>
<b>Nonforest</b>			
Cropland	29 400	60 300	85 000
Pasture	174 100	146 100	312 300
Other	85 000	108 000	205 500
<b>Total</b>	<b>288 500</b>	<b>314 400</b>	<b>602 900</b>
<b>Total land</b>	<b>436 600</b>	<b>453 700</b>	<b>890 300</b>

-- = Dashes indicate the corresponding cell was not sampled.

<sup>a</sup> 1985 estimate.

<sup>b</sup> Not included in "timberland" in subsequent tables.

**Table 2—Land area by survey region and broad land cover class, Puerto Rico, 1980–1990**

Survey region and land cover class	Survey year	
	1980	1990
----- <i>Hectares</i> -----		
<b>Commercial survey region</b>		
Forest	130 500	148 100
Nonforest	306 100	288 500
<b>Total</b>	<b>436 600</b>	<b>436 600</b>
<b>Noncommercial survey region<sup>a</sup></b>		
Forest	148 200	139 399
Nonforest	305 500	314 400
<b>Total</b>	<b>453 700</b>	<b>453 700</b>
<b>All regions</b>		
Forest	278 700	287 400
Nonforest	611 600	602 900
<b>Total</b>	<b>890 300</b>	<b>890 300</b>

<sup>a</sup> The noncommercial region was not surveyed; estimates reported are from the 1985 forest area assessment.

**Table 3—Area of timberland by Life Zone, soil group, and forest class, Puerto Rico, 1990**

Life Zone and soil group	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
-----Hectares-----						
<b>Subtropical Moist Forest</b>						
Deep volcanic soils	12 928	5 841	4 268	10 110	1 416	1402
Shallow volcanic soils	28 938	11 683	15 839	27 522	1 416	0
Granitic soils	12 994	0	12 994	12 994	0	0
Limestone soils	21 061	10 709	8 228	18 937	2 124	0
Total	75 921	28 233	41 329	69 563	4 956	1402
<b>Subtropical Wet Forest</b>						
Deep volcanic soils	43 862	1 947	22 147	24 094	14 161	5608
Shallow volcanic soils	5 628	974	1 114	2 088	3 540	0
Granitic soils	16 574	1 947	13 919	15 866	708	0
Limestone soils	1 947	1 947	0	1 947	0	0
Total	68 011	6 815	37 180	43 995	18 409	5608
<b>Total, all areas</b>	143 932	35 048	78 509	113 558	23 365	7010

**Table 4—Area of timberland by stand-volume class and forest class, Puerto Rico, 1990**

Timber volume (m <sup>3</sup> per hectare)	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
-----Hectares-----						
< 25	61 665	29 207	26 793	56 000	5 664	0
25 to 50	32 666	3 894	19 609	23 504	4 956	4206
50 to 75	25 807	974	14 227	15 201	9 205	1402
75 to 100	9 617	0	5 382	5 382	2 832	1402
> 100	14 178	974	12 496	13 470	708	0
Total	143 933	35 049	78 507	113 557	23 365	7010

**Table 5—Area of timberland by stand-volume class and soil group, Puerto Rico, 1990**

Timber volume (m <sup>3</sup> per hectare)	Soil group				
	All groups	Deep volcanic	Shallow volcanic	Granitic	Limestone
----- <i>Hectares</i> -----					
< 25	61 665	14 049	21 257	10 981	15 377
25 to 50	32 666	9 441	8 789	8 228	6 209
50 to 75	25 807	17 019	2 390	6 399	0
75 to 100	9 617	7 486	2 131	0	0
> 100	14 178	8 795	0	3 960	1 423
Total	143 933	56 790	34 567	29 568	23 009

**Table 6—Area of timberland by stand-size class and forest class, Puerto Rico, 1990**

Stand-size class	Forest class					
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	Active coffee shade
----- <i>Hectares</i> -----						
Sawtimber	65 930	5 841	37 488	43 329	16 993	5608
Poletimber	47 458	7 789	32 603	40 392	5 664	1402
Sapling and seedling	30 544	21 419	8 417	29 836	708	0
Total	143 932	35 049	78 508	113 557	23 365	7010

**Table 7—Area of timberland by stand-size class and soil group, Puerto Rico, 1990**

Stand-size class	Soil group				
	All groups	Deep volcanic	Shallow volcanic	Granitic	Limestone
----- <i>Hectares</i> -----					
Sawtimber	65 930	30 980	14 872	15 741	4 337
Poletimber	47 458	20 801	8 968	8 893	8 795
Sapling and seedling	30 544	5 009	10 725	4 933	9 876
Total	143 932	56 790	34 565	29 567	23 008

**Table 8—Area of timberland by stand-age class and forest class, Puerto Rico, 1990**

Stand age (years)	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
----- <i>Hectares</i> -----						
< 30	78 588	35 049	31 559	66 608	6 372	5608
> 30	27 679	0	17 072	17 072	9 205	1402
Mixed ages	5 684	0	4 268	4 268	1 416	0
Mature without past age <sup>a</sup>	31 981	0	25 609	25 609	6 372	0
Total	143 932	35 049	78 508	113 557	23 365	7010

<sup>a</sup>Stand age could not be reliably estimated for these stands because past age data were unavailable.

**Table 9—Area of timberland by stand-age class and soil group, Puerto Rico, 1990**

Stand age (years)	Soil group				
	All groups	Deep volcanic	Shallow volcanic	Granitic	Limestone
----- <i>Hectares</i> -----					
< 30	78 588	25 564	24 620	13 210	15 193
> 30	27 679	14 180	6 392	2 845	4 262
Mixed ages	5 684	2 131	0	1 423	2 131
Mature without past age <sup>a</sup>	31 981	14 915	3 553	12 090	1 423
Total	143 932	56 790	34 565	29 568	23 009

<sup>a</sup>Stand age could not be reliably estimated for these stands because past age data were unavailable.

**Table 10—Area of timberland by basal area class and forest class, Puerto Rico, 1990**

Basal area (m <sup>2</sup> per hectare)	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
----- <i>Hectares</i> -----						
< 5	17 000	15 577	1 423	17 000	0	0
5 to 10	23 551	8 762	11 263	20 025	2 124	1402
10 to 15	35 238	4 868	18 376	23 244	7 789	4206
15 to 20	30 570	3 894	19 609	23 504	5 664	1402
20 to 25	23 094	974	16 456	17 429	5 664	0
25 to 30	4 970	0	2 845	2 845	2 124	0
30 to 35	5 691	0	5 691	5 691	0	0
> 35	3 819	974	2 845	3 819	0	0
Total	143 933	35 049	78 508	113 557	23 365	7010

**Table 11—Area of timberland by basal area class and stand-size class, Puerto Rico, 1990**

Basal area (m <sup>2</sup> per hectare)	Stand-size class			
	All classes	Sawtimber	Poletimber	Sapling and seedling
----- <i>Hectares</i> -----				
< 5	17 000	0	0	17 000
5 to 10	23 551	9 719	4 463	9 368
10 to 15	35 238	15 568	15 495	4 176
15 to 20	30 570	19 385	11 185	0
20 to 25	23 094	14 866	8 228	0
25 to 30	4 970	3 547	1 423	0
30 to 35	5 691	1 423	4 268	0
> 35	3 819	1 423	2 396	0
<b>Total</b>	<b>143 933</b>	<b>65 931</b>	<b>47 458</b>	<b>30 544</b>

**Table 12—Area of stocked timberland by Life Zone, soil group, and slope class, Puerto Rico, 1990**

Life Zone and soil group	Percent slope class				
	All classes	0 to 10	11 to 25	26 to 45	> 45
----- <i>Hectares</i> -----					
<b>Subtropical Moist Forest</b>					
Deep volcanic soils	12 928	2 287	1 123	4 164	5 353
Shallow volcanic soils	28 938	3 107	6 325	10 239	9 267
Granitic soils	12 994	474	2 166	3 011	7 343
Limestone soils	21 061	2 614	2 571	4 841	11 036
<b>Total</b>	<b>75 921</b>	<b>8 482</b>	<b>12 185</b>	<b>22 255</b>	<b>32 999</b>
<b>Subtropical Wet Forest</b>					
Deep volcanic soils	43 862	4 007	6 609	12 609	20 637
Shallow volcanic soils	5 628	0	944	1 303	3 381
Granitic soils	16 575	0	1 659	4 380	10 536
Limestone soils	1 947	0	649	1 298	0
<b>Total</b>	<b>68 012</b>	<b>4 007</b>	<b>9 861</b>	<b>19 590</b>	<b>34 554</b>
<b>Total, all areas</b>	<b>143 933</b>	<b>12 489</b>	<b>22 046</b>	<b>41 845</b>	<b>67 553</b>

Table 13—Number of live trees by species and diameter class, ranked by basal area plurality, Puerto Rico, 1990\*

Species code	Species name	Total basal area (m <sup>2</sup> )	Diameter class (centimeters)											
			All classes	5	10	15	20	25	30	35	40	45	50	55+
----- Thousand trees -----														
234	<i>Spathodea campanulata</i> Beauv.	229 543	21 436	12 014	5 529	1 550	929	717	289	195	61	41	54	56
109	<i>Guarea guidonia</i> (L.) Sleumer	229 502	17 569	11 346	2 614	1 463	740	494	297	252	99	94	67	101
64	<i>Inga vera</i> Willd.	116 731	6 761	3 164	1 397	898	415	370	205	122	99	44	32	14
20	<i>Cecropia peltata</i> L.	104 570	3 752	1 232	902	362	214	257	214	170	61	64	41	31
81	<i>Anandra inermis</i> (W. Wright) H.B.K.	92 232	10 415	5 667	2 820	1 025	389	311	120	38	14	19	9	3
236	<i>Tabebuia heterophylla</i> (DC.) Britton	91 324	11 711	7 649	1 680	1 487	470	248	64	75	20	8	6	4
188	<i>Eugenia jambos</i> L.	73 453	14 370	11 491	1 531	677	348	167	81	47	7	15	0	5
62	<i>Inga fagifolia</i> (L.)	61 134	3 367	1 371	969	266	315	143	186	34	54	15	6	9
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	51 428	996	312	254	0	108	101	47	64	14	19	18	56
131	<i>Mangifera indica</i> L.	44 451	393	117	0	52	31	20	43	19	10	23	9	64
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	43 031	2 831	1 042	778	472	222	163	103	19	10	12	3	8
7	<i>Euterpe globosa</i> Gaertn.	40 102	1 762	0	117	929	575	130	11	0	0	0	0	0
48	<i>Ocotea leucoxylon</i> (Sw.) Mez	37 708	7 050	5 636	739	203	213	198	24	25	0	12	0	0
9	<i>Roustea borinquena</i> O. F. Cook	35 532	674	117	80	44	91	77	89	81	73	21	0	2
171	<i>Casearia guianensis</i> (Aubl.) Urban	33 753	23 171	552	0	30	0	0	0	0	0	0	0	0
100	<i>Citrus sinensis</i> Osbeck	28 041	3 576	1 850	867	708	79	72	0	0	0	0	0	0
200	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	24 763	2 693	2 147	58	190	142	78	41	20	0	11	3	3
1	<i>Cyathea arborea</i> (L.) J. E. Smith	23 641	2 237	230	1 405	524	79	0	0	0	0	0	0	0
172	<i>Casearia sylvestris</i> Sw.	23 024	15 461	14 941	441	79	0	0	0	0	0	0	0	0
138	<i>Cupania americana</i> L.	22 714	2 876	1 344	1 159	220	60	61	18	0	14	0	0	0
592	<i>Clidemia umbrosa</i> (Sw.) Cogn.	22 550	9 310	8 664	458	57	36	49	0	36	9	0	0	0
118	<i>Alchornea latifolia</i> Sw.	22 513	1 942	1 262	117	232	224	63	33	0	0	0	3	8
161	<i>Calophyllum brasiliense</i> Camb.	21 980	2 609	2 380	0	53	0	53	50	11	26	14	12	10
224	<i>Cordia sulcata</i> DC.	21 457	2 331	1 355	440	244	163	79	24	26	0	0	0	0
163	<i>Clusia rosea</i> Jacq.	20 620	1 826	668	779	39	201	126	0	12	0	0	0	0
134	<i>Spondias mombin</i> L.	20 134	798	456	0	228	0	48	25	0	9	9	0	22
193	<i>Myrcia splendens</i> (Sw.) DC.	18 016	7 336	6 364	873	64	35	0	0	0	0	0	0	0
334	<i>Nectandra sintenisii</i> Mez (E)	17 920	2 458	1 810	350	197	23	24	27	6	5	0	0	17
198	<i>Miconia prasina</i> (Sw.) DC.	17 694	7 994	7 200	641	153	0	0	0	0	0	0	0	0
227	<i>Citharexylum fruticosum</i> L.	17 456	2 119	1 224	598	170	120	90	17	0	0	0	0	0
17	<i>Artocarpus altitilis</i> (Parkinson) Fosberg	17 318	817	478	80	109	56	0	12	31	28	6	7	10
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	17 304	2 016	1 391	367	108	36	51	34	13	6	7	0	2
22	<i>Ficus laevigata</i> Vahl	15 583	390	183	0	78	59	0	8	18	10	22	0	12
249	<i>Terebraria resinosa</i> (Vahl) Sprague	15 219	2 728	1 380	1 109	205	33	0	0	0	0	0	0	0
228	<i>Pettitia domingensis</i> Jacq.	15 147	609	239	0	236	15	0	92	21	0	0	0	0
76	<i>Hymenaea courbaril</i> L.	14 132	449	0	208	74	0	81	35	26	20	0	6	0
221	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	14 033	1 759	1 412	116	32	93	56	28	6	11	0	0	5
211	<i>Micropholis chrysophylloides</i> Pierre	12 914	1 331	699	350	112	117	0	54	0	0	0	0	0
180	<i>Buchenavia capitata</i> (Vahl) Eichl.	12 800	261	0	117	0	56	22	18	13	20	11	0	4
721	<i>Coffea dewevrei</i> Wildem. & T. Dur.	12 601	2 812	2 014	622	120	36	21	0	0	0	0	0	0
195	<i>Psidium guajava</i> L.	12 201	9 686	9 436	251	0	0	0	0	0	0	0	0	0
184	<i>Terminalia catappa</i> L.	11 991	454	319	0	0	44	20	0	12	36	23	0	0
170	<i>Casearia decandra</i> Jacq.	11 989	6 385	6 231	0	95	39	17	0	0	0	0	0	3

Species code	Species name	Diameter class (centimeters)															
		All classes	5	10	15	20	25	30	35	40	45	50	55+				
199	<i>Tetragyia elaeagnoides</i> (Sw.) DC.	11 006	3 274	429	125	0	29	0	0	0	0	0	0	0	0	0	0
240	<i>Coffea arabica</i> L.	10 901	9 710	116	0	0	0	0	0	0	0	0	0	0	0	0	0
87	<i>Ormosia krugii</i> Urban	10 756	0	117	201	126	49	0	0	0	0	0	0	0	0	0	0
106	<i>Dacryodes excensae</i> Vahl	10 666	0	0	0	62	0	51	12	20	0	6	0	0	0	0	0
999	Unidentified species	10 562	2 401	466	47	38	0	18	0	0	4	0	0	0	0	0	0
169	<i>Casearia arborea</i> (L. C. Rich.) Urban	10 545	3 641	175	50	77	0	0	0	0	0	0	0	0	0	0	0
125	<i>Hura crepitans</i> L.	10 543	0	0	0	55	31	10	8	6	5	0	16	0	0	0	0
115	<i>Byrsonima coriacea</i> (Sw.) DC.	10 228	318	175	0	31	47	35	0	19	12	0	0	0	0	0	0
151	<i>Montezuma speciosissima</i> Sessé & Moc.	10 017	247	0	117	0	85	8	12	18	8	0	0	0	0	0	0
192	<i>Myrcia deflexa</i> (Poir.) DC.	9 650	2 076	466	73	0	0	15	0	0	0	13	0	0	0	0	0
156	<i>Guazuma ulmifolia</i> Lam.	9 350	404	91	136	40	0	0	0	10	7	0	4	0	0	0	0
149	<i>Sloanea berteriana</i> Choisy	9 260	611	233	117	133	71	30	18	0	9	0	0	0	0	0	0
208	<i>Dipholis salicifolia</i> (L.) A. DC.	8 813	599	80	233	117	169	0	0	0	0	0	0	0	0	0	0
105	<i>Bursera inaruba</i> (L.) Sarg.	8 781	1 013	600	291	0	27	63	18	12	0	0	0	0	0	0	0
33	<i>Torrubia fragrans</i> (Dum.-Cours.) Standley	8 578	1 620	1 194	74	0	0	0	0	0	0	0	0	0	0	0	0
213	<i>Pouteria multiflora</i> (A. DC.) Eyma	8 274	55	0	0	0	0	18	12	14	4	0	5	0	0	0	0
61	<i>Albizia procera</i> (Roxb.) Benth.	7 930	503	159	45	106	0	34	0	0	0	0	0	0	0	0	0
340	<i>Phoebe elongata</i> (Vahl) Nees	7 891	790	464	175	62	42	25	10	11	0	0	3	0	0	0	0
108	<i>Cedrela odorata</i> L.	7 747	329	115	115	0	66	0	0	19	8	0	7	0	0	0	0
114	<i>Trichilia pallida</i> Sw.	7 701	4 895	313	0	0	0	0	0	0	0	0	0	0	0	0	0
133	<i>Spondias dulcis</i> Parkinson	7 453	134	80	0	0	0	0	0	27	0	16	6	5	0	0	0
143	<i>Thouinia striata</i> Radlk.	7 117	2 138	1 907	159	0	72	0	0	0	0	0	0	0	0	0	0
	Other species (134)	200 205	49 571	42 005	5 358	958	402	338	193	179	62	39	23	22			
	Total	2 194 205	308 780	232 253	40 615	15 676	8425	5104	2717	1665	893	593	329	511			

-----  
Thousand trees  
-----

\*Columns may not add up to totals as a result of rounding numbers.

Table 14—Number of growing-stock trees by species and diameter class, ranked by basal area plurality, Puerto Rico, 1990<sup>a</sup>

Species code	Species name	Total basal area (m <sup>2</sup> )	Diameter class (centimeters)											
			All classes	5	10	15	20	25	30	35	40	45	50	55+
----- Thousand trees -----														
234	<i>Spathodea campanulata</i> Beauv.	190 169	14 049	6 239	4 522	1 229	807	641	255	168	61	31	45	49
109	<i>Guarea guidonia</i> (L.) Sleumer	123 615	5 909	2 994	662	872	485	375	188	158	33	51	42	48
20	<i>Cecropia peltata</i> L.	96 080	2 782	602	669	316	402	257	188	170	61	56	31	31
64	<i>Inga vera</i> Willd.	75 438	3 570	1 335	582	765	380	211	111	73	68	16	16	12
236	<i>Tabebuia heterophylla</i> (DC.) Britton	72 186	5 786	2 367	1 404	1 235	397	226	51	75	20	0	6	4
81	<i>Andira inermis</i> (W. Wright) H.B.K.	59 359	4 498	1 547	1 532	789	212	286	92	21	0	19	0	0
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	44 232	470	116	58	0	31	76	16	64	14	19	18	56
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	40 435	2 535	926	661	427	222	163	83	19	10	12	3	8
7	<i>Euterpe globosa</i> Gaertn.	38 360	1 555	0	0	838	575	130	11	0	0	0	0	0
62	<i>Inga jagifolia</i> (L.)	36 474	1 493	382	472	142	180	105	138	27	36	7	0	3
9	<i>Roystonia borinquena</i> O. F. Cook	35 532	674	117	80	44	91	77	89	81	73	21	0	2
188	<i>Eugenia jambos</i> L.	25 962	6 089	4 629	1 092	128	184	33	16	0	0	8	0	0
40	<i>Ocotea leucoxylon</i> (Sw.) Mez	22 172	4 721	3 840	408	203	145	125	0	0	0	0	0	0
1	<i>Cyathea arborea</i> (L.) J. E. Smith	20 620	1 840	115	1 199	448	79	0	0	0	0	0	0	0
161	<i>Calophyllum brasiliense</i> Camb.	20 534	2 384	2 183	0	53	0	25	50	11	26	14	12	10
118	<i>Alchornea latifolia</i> Sw.	20 194	987	323	117	232	224	50	33	0	0	0	0	8
138	<i>Cupania americana</i> L.	14 746	1 520	641	598	168	30	61	18	0	5	0	0	0
163	<i>Clusia rosea</i> Jacq.	13 937	1 287	472	583	0	121	100	0	12	0	0	0	0
17	<i>Artocarpus altitilis</i> (Parkinson) Fosberg	13 854	634	319	80	109	56	0	12	24	19	0	7	8
76	<i>Hymenaea courbaril</i> L.	13 617	333	0	91	74	0	81	35	26	20	0	6	0
334	<i>Nectandra sintenisii</i> (E)	13 177	977	581	117	197	23	24	8	6	5	0	0	17
224	<i>Cordia sulcata</i> DC.	12 147	987	567	116	128	70	79	15	12	0	0	0	0
134	<i>Spondias mombin</i> L.	12 125	284	183	0	0	0	48	25	0	9	4	0	15
592	<i>Clidemia umbrosa</i> (Sw.) Cogn.	11 243	2 785	2 553	58	57	36	49	0	23	9	0	0	0
228	<i>Peltita domingensis</i> Jacq.	11 043	556	239	0	236	0	0	72	8	0	0	0	0
249	<i>Terebraria resinosa</i> (Vahl) Sprague	10 951	1 606	515	950	141	0	0	0	0	0	0	0	0
106	<i>Dacryodes excelsea</i> Vahl	10 666	151	0	0	0	62	0	51	12	20	0	6	0
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	10 649	1 029	779	80	53	36	20	34	13	6	7	0	0
184	<i>Terminalia catappa</i> L.	10 592	206	80	0	0	44	20	0	12	27	23	0	0
221	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	10 228	903	590	116	32	93	39	16	6	11	0	0	0
193	<i>Myrcia splendens</i> (Sw.) DC.	9 805	3 300	2 654	582	64	0	0	0	0	0	0	0	0
200	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	9 515	714	429	58	104	65	29	8	14	0	4	3	1
211	<i>Micropholis chrysophylloides</i> Pierre	9 202	1 039	699	117	112	73	0	38	0	0	0	0	0
87	<i>Ormosia knugii</i> Urban	8 385	375	0	117	118	92	49	0	0	0	0	0	0
131	<i>Mangifera indica</i> L.	8 268	93	0	0	52	0	20	0	5	0	0	3	13
180	<i>Buchenavia capitata</i> (Vahl) Eichl.	7 700	103	0	0	0	32	22	18	0	20	11	0	0
22	<i>Ficus laevigata</i> Vahl	7 666	82	0	0	0	43	0	8	0	10	15	0	6
171	<i>Casearia guianensis</i> (Aubl.) Urban	7 477	6 007	5 977	0	0	30	0	0	0	0	0	0	0
208	<i>Dipholis salicifolia</i> (L.) A. DC.	7 347	461	0	233	59	169	0	0	0	0	0	0	0
999	Unidentified species	7 247	1 431	1 012	350	47	0	0	18	0	0	4	0	0
192	<i>Myrcia deflexa</i> (Poir.) DC.	6 837	1 482	1 105	350	0	0	0	15	0	0	0	13	0
74	<i>Cassia siamea</i> Lam.	6 756	1 644	1 116	478	40	0	0	0	9	0	0	0	0
115	<i>Byrsonima coriacea</i> (Sw.) DC.	6 690	258	175	0	0	31	22	0	0	19	12	0	0



**Table 15—Number of life trees on timberland by diameter class and forest class, Puerto Rico, 1990**

Diameter class (centimeters)	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
	----- <i>Hectares</i> -----					
5	1614	1536	1758	1690	1556	573
10	282	209	348	305	196	197
15	109	58	138	114	91	93
20	59	26	81	64	46	12
25	35	20	43	36	35	31
30	19	7	23	18	24	12
35	12	4	14	11	15	7
40	6	2	7	6	9	4
45	4	1	5	4	7	3
50	2	0	3	2	4	2
55+	4	2	3	3	7	11
<b>Total</b>	<b>2146</b>	<b>1865</b>	<b>2423</b>	<b>2253</b>	<b>1990</b>	<b>945</b>

**Table 16a—Basal area and number of live trees per hectare of timberland for selected species, secondary forests, Puerto Rico, 1990<sup>a</sup>**

Species code	Species name	Basal area		Number of trees
		<i>m<sup>2</sup>/ha</i>	<i>Percent</i>	<i>Hectares</i>
234	<i>Spathodea campanulata</i> Beauv.	1.86	12.4	177
109	<i>Guarea guidonia</i> (L.) Sleumer	1.19	7.9	106
236	<i>Tabebuia heterophylla</i> (DC.) Britton	.78	5.2	103
81	<i>Andira inermis</i> (W. Wright) H.B.K.	.68	4.5	82
20	<i>Cecropia peltata</i> L.	.59	3.9	21
188	<i>Eugenia jambos</i> L.	.54	3.6	86
64	<i>Inga vera</i> Willd.	.46	3.1	26
7	<i>Euterpe globosa</i> Gaertn.	.35	2.3	16
62	<i>Inga fagifolia</i> (L.)	.34	2.3	18
9	<i>Roystonea borinquena</i> O. F. Cook	.28	1.8	5
171	<i>Casearia guianensis</i> (Aubl.) Urban	.28	1.8	196
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	.28	1.9	16
48	<i>Ocotea leucoxylon</i> (Sw.) Mez	.28	1.9	42
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	.26	1.7	6
131	<i>Mangifera indica</i> L.	.19	1.3	3
161	<i>Calophyllum brasiliense</i> Camb.	.19	1.3	23
138	<i>Cupania americana</i> L.	.19	1.3	25
163	<i>Clusia rosea</i> Jacq.	.18	1.2	16
172	<i>Casearia sylvestris</i> Sw.	.18	1.2	122
118	<i>Alchornea latifolia</i> Sw.	.17	1.1	13
592	<i>Clidemia umbrosa</i> (Sw.) Cogn.	.17	1.1	66
1	<i>Cyathea arborea</i> (L.) J. E. Smith	.15	1.0	12
193	<i>Myrcia splendens</i> (Sw.) DC.	.15	1.0	61
227	<i>Citharexylum fruticosum</i> L.	.15	1.0	19
134	<i>Spondias mombin</i> L.	.15	1.0	7
17	<i>Artocarpus altilis</i> (Parkinson) Fosberg	.14	.9	7
198	<i>Miconia prasina</i> (Sw.) DC.	.13	.9	57
249	<i>Terebraria resinosa</i> (Vahl) Sprague	.13	.9	24
76	<i>Hymenaea courbaril</i> L.	.12	.8	4
100	<i>Citrus sinensis</i> Osbeck	.12	.8	13
	Other species (157)	4.36	29.0	876
	Total	15.07	100.0	2248

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

**Table 16b—Basal area and number of live trees per hectare of timberland for selected species, abandoned coffee shade, Puerto Rico, 1990<sup>a</sup>**

Species code	Species name	Basal area		Number of trees
		<i>m<sup>2</sup>/ha</i>	<i>Percent</i>	<i>Hectares</i>
109	<i>Guarea guidonia</i> (L.) Sleumer	3.51	20.7	225
64	<i>Inga vera</i> Willd.	2.15	12.7	135
20	<i>Cecropia peltata</i> L.	1.34	7.9	57
131	<i>Mangifera indica</i> L.	.78	4.6	3
62	<i>Inga fagifolia</i> (L.)	.77	4.6	49
200	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	.71	4.2	59
84	<i>Erythrina poeppigiana</i> (Walp.) O.F. Cook	.63	3.7	13
234	<i>Spathodea campanulata</i> Beauv.	.60	3.5	52
81	<i>Andira inermis</i> (W. Wright) H.B.K.	.53	3.1	44
188	<i>Eugenia jambos</i> L.	.53	3.1	198
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	.46	2.7	44
224	<i>Cordia sulcata</i> DC.	.37	2.2	45
240	<i>Coffea arabica</i> L.	.28	1.6	236
221	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	.26	1.5	29
48	<i>Ocotea leucoxydon</i> (Sw.) Mez	.24	1.4	90
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	.23	1.4	6
100	<i>Citrus sinensis</i> Osbeck	.21	1.2	21
334	<i>Nectandra sintenisii</i> Mez (E)	.21	1.3	25
213	<i>Pouteria multiflora</i> (A.DC.) Eyma	.20	1.2	2
721	<i>Coffea dewevrei</i> Wildem. & T. Dur.	.19	1.1	52
9	<i>Roystonea borinquena</i> O.F. Cook	.18	1.0	5
134	<i>Spondias mombin</i> L.	.15	.9	1
230	<i>Vitex divaricata</i> Sw.	.15	.9	6
340	<i>Phoebe elongata</i> (Vahl) Nees	.14	.8	23
198	<i>Miconia prasina</i> (Sw.) DC.	.13	.8	63
	Other species (46)	2.00	11.8	501
	Total	16.94	100.0	1694

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

**Table 16c—Basal area and number of live trees per hectare of timberland for selected species, active coffee shade, Puerto Rico, 1990<sup>a</sup>**

Species code	Species name	Basal area		Number of trees
		<i>m<sup>2</sup>/ha</i>	<i>Percent</i>	<i>Hectares</i>
64	<i>Inga vera</i> Willd.	2.04	16.3	87
109	<i>Guarea guidonia</i> (L.) Sleumer	1.70	13.6	43
100	<i>Citrus sinensis</i> Osbeck	1.40	11.2	227
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	1.00	8.0	2
1	<i>Cyathea arborea</i> (L.) J. E. Smith	.99	8.0	125
20	<i>Cecropia peltata</i> L.	.83	6.7	11
131	<i>Mangifera indica</i> L.	.67	5.3	2
62	<i>Inga fagifolia</i> (L.)	.63	5.1	23
234	<i>Spathodea campanulata</i> Beauv.	.58	4.6	23
22	<i>Ficus laevigata</i> Vahl	.50	4.0	2
81	<i>Andira inermis</i> (W. Wright) H.B.K.	.33	2.7	4
721	<i>Coffea dewevrei</i> Wildem. & T. Dur.	.23	1.8	19
240	<i>Coffea arabica</i> L.	.20	1.6	229
51	<i>Persea americana</i> Mill.	.17	1.3	1
115	<i>Byrsonima coriacea</i> (Sw.) DC.	.17	1.3	2
99	<i>Citrus paradisi</i> Macfadyen	.17	1.3	3
118	<i>Alchornea latifolia</i> Sw.	.17	1.3	5
236	<i>Tabebuia heterophylla</i> (DC.) Britton	.17	1.3	2
294	<i>Ficus stahlii</i> Warb. (E)	.17	1.3	1
221	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	.17	1.3	1
108	<i>Cedrela odorata</i> L.	.09	.7	33
592	<i>Clidemia umbrosa</i> (Sw.) Cogn.	.05	.4	49
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	.04	.3	16
48	<i>Ocotea leucoxylon</i> (Sw.) Mez	.01	.1	16
279	<i>Piper amalago</i> L.	.01	.1	16
	Total	12.49	99.6	942

<sup>a</sup>Columns may not add up to totals as a result of rounding numbers.

**Table 17—Basal area of poletimber and sawtimber trees by tree class and forest class, Puerto Rico, 1990**

Tree class	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
----- <i>Square meters of basal area</i> -----						
<b>Poletimber-size trees</b>						
Growing stock	544 257	39 738	47 926	387 665	127 398	29 195
Rough-and-rotten	254 642	29 195	132 435	161 630	75 495	17 517
Total	798 899	68 933	480 361	549 295	202 893	46 712
<b>Sawtimber-size trees</b>						
Growing stock	572 224	68 123	406 371	474 493	80 214	17 517
Rough-and-rotten	197 469	27 573	132 809	160 382	30 080	7 007
Total	769 693	95 696	539 180	634 875	110 294	24 524
Total, growing stock	1 116 481	107 861	754 297	862 158	207 612	46 712
Total, rough-and-rotten	452 111	56 768	265 244	322 012	105 575	24 524
<b>Total, poletimber and sawtimber trees</b>	1 568 592	164 629	1 019 541	1 184 170	313 187	71 236

**Table 18—Basal area of poletimber and sawtimber trees by tree class, Life Zone, and soil group, Puerto Rico, 1990**

Tree class and Life Zone	Soil group				
	groups	All volcanic	Deep volcanic	Shallow Granitic	Limestone
----- <i>Square meters of basal area</i> -----					
<b>Subtropical Moist Forest</b>					
Poletimber-size trees					
Growing stock	222 129	88 539	58 201	43 236	32 153
Rough-and-rotten	102 157	29 112	34 224	24 117	14 704
Total	<u>324 286</u>	<u>117 651</u>	<u>92 425</u>	<u>67 353</u>	<u>46 857</u>
Sawtimber-size trees					
Growing stock	220 117	48 594	54 555	51 847	65 121
Rough-and-rotten	108 312	22 355	35 282	28 501	22 175
Total	<u>328 429</u>	<u>70 949</u>	<u>89 837</u>	<u>80 348</u>	<u>87 296</u>
Total, growing stock	442 246	137 133	112 756	95 084	97 274
Total, rough-and-rotten	210 470	51 466	69 506	52 618	36 879
Total, Subtropical Moist Forest	<u>652 716</u>	<u>188 599</u>	<u>182 262</u>	<u>147 702</u>	<u>134 153</u>
<b>Subtropical Wet Forest</b>					
Poletimber-size trees					
Growing stock	322 128	211 796	27 131	79 146	4 055
Rough-and-rotten	152 485	91 873	20 864	33 259	6 488
Total	<u>474 613</u>	<u>303 669</u>	<u>47 995</u>	<u>112 405</u>	<u>10 543</u>
Sawtimber-size trees					
Growing stock	352 107	246 818	20 848	78 764	5 677
Rough-and-rotten	89 157	58 910	12 664	15 961	1 622
Total	<u>441 264</u>	<u>305 728</u>	<u>33 512</u>	<u>94 725</u>	<u>7 299</u>
Total, growing stock	674 235	458 614	47 979	157 910	9 732
Total, rough-and-rotten	241 642	150 783	33 528	49 220	8 110
Total, Subtropical Wet Forest	<u>915 877</u>	<u>609 397</u>	<u>81 507</u>	<u>207 130</u>	<u>17 842</u>
<b>Total, Life Zones</b>	<u>1 568 593</u>	<u>797 996</u>	<u>263 769</u>	<u>354 832</u>	<u>151 995</u>

**Table 19—Volume of timber on timberland by class of timber and forest class, Puerto Rico, 1990<sup>a</sup>**

Class of timber	Forest class					Active coffee shade
	All classes	Reversion	Other secondary forest	All secondary forest	Abandoned coffee shade	
----- <i>Thousand cubic meters</i> -----						
<b>Growing-stock trees</b>						
Sawtimber trees						
Saw-log portion	2 647.3	154.9	1695.7	1850.6	613.1	183.6
Upper stem	648.3	38.9	419.4	458.3	155.0	35.1
Branches and forks	895.4	51.2	586.4	637.6	225.7	32.1
Sound cull	686.5	38.1	486.2	524.2	142.5	19.7
Total, timber volume	4 877.5	283.0	3187.8	3470.8	1136.3	270.5
Poletimber trees						
Bole volume	2 583.7	321.4	1778.5	2100.0	385.4	98.3
Branches and forks	862.1	75.3	586.1	661.4	170.1	30.6
Sound cull	416.9	14.1	369.2	383.3	29.9	3.7
Total, timber volume	3 862.8	410.8	2733.9	3144.7	585.4	132.7
Total, all growing-stock trees	8 740.2	693.8	5921.7	6615.5	1721.7	403.1
<b>Rough-and-rotten trees</b>						
Rough trees						
Bole volume	1 217.2	136.1	705.1	841.3	296.6	79.3
Branches and forks	946.4	144.8	558.4	703.2	184.4	58.7
Sound cull	485.0	53.0	262.8	315.8	145.0	24.1
Total, timber volume	2 648.5	334.0	1526.3	1860.3	626.0	162.2
Rotten trees						
Bole volume	56.0	7.1	34.3	39.3	9.4	7.3
Branches and forks	36.8	2.9	17.1	20.9	15.2	.7
Sound cull	43.0	3.1	29.9	34.1	8.0	.9
Total, timber volume	135.8	13.1	81.3	94.4	32.6	8.9
Total, all rough-and-rotten trees	2 784.3	347.0	1607.6	1954.7	658.6	171.1
<b>Total, all timber</b>	11 524.6	1040.8	7529.3	8570.1	2380.2	574.2

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

**Table 20—Volume of timber on timberland by Life Zone, soil group, and tree class, Puerto Rico, 1990<sup>a</sup>**

Life zone and tree class	Soil group				Limestone
	groups	All volcanic	Deep volcanic	Shallow Granitic	
----- <i>Thousand cubic meters</i> -----					
<b>Subtropical Moist Forest</b>					
Sawtimber trees	1 940.5	781.3	487.2	372.2	299.7
Poletimber trees	1 390.2	298.4	355.7	317.1	418.9
Total, all growing-stock trees	3 330.6	1079.7	842.9	689.4	718.6
Rough trees	1 123.1	276.6	319.0	295.6	231.8
Rotten trees	52.1	21.5	16.0	12.0	2.6
Total, all rough-and-rotten trees	1 175.2	298.1	335.0	307.6	234.5
Total, all timber	4 505.8	1377.8	1178.0	997.0	953.1
<b>Subtropical Wet Forest</b>					
Sawtimber trees	2 937.0	1992.6	221.1	690.7	32.6
Poletimber trees	2 472.6	1778.1	162.1	493.7	38.7
Total, all growing-stock trees	5 409.6	3770.7	383.2	1184.4	71.3
Rough trees	1 525.4	944.5	207.8	303.0	70.2
Rotten trees	83.7	72.6	5.0	6.1	0
Total, all rough-and-rotten trees	1 609.1	1017.1	212.7	309.1	70.2
Total, all timber	7 018.7	4787.8	596.0	1493.5	141.5
<b>Total, all Life Zones</b>	11 524.6	6165.6	1773.9	2490.5	1094.5

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

Table 21—Volume of timber on timberland by species and diameter class, Puerto Rico, 1990<sup>a</sup>

Species code	Species name	Diameter class (centimeters)									
		All classes	15	20	25	30	35	40	45	50	55+
----- Thousand cubic meters -----											
109	<i>Guarea guidonia</i> (L.) Sleumer	1 282.4	123.4	155.7	142.5	161.6	178.9	85.8	109.1	94.2	214.8
234	<i>Spathodea campanulata</i> Beauv.	1 109.4	104.5	167.0	195.5	146.7	140.2	51.2	53.5	79.7	158.4
64	<i>Inga vera</i> Willd.	820.6	102.5	109.4	176.6	121.2	104.1	90.2	44.4	42.8	29.4
20	<i>Cecropia peltata</i> L.	812.0	28.9	96.5	129.3	131.5	135.1	77.3	85.8	60.5	67.1
62	<i>Inga fagifolia</i> (L.)	398.9	55.4	67.1	42.2	119.6	27.1	55.3	15.5	6.0	10.5
236	<i>Tabebuia heterophylla</i> (DC.) Britton	395.0	120.3	65.0	89.3	29.8	51.7	14.7	9.6	7.1	7.7
81	<i>Andira inermis</i> (W. Wright) H.B.K.	381.5	84.2	75.3	94.6	54.7	20.1	5.7	30.4	11.5	4.9
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	364.4	0	12.5	28.9	18.0	53.2	14.1	23.6	23.4	175.7
9	<i>Roystonea borinquena</i> O. F. Cook	310.7	1.1	7.8	18.4	64.8	79.6	96.8	35.5	0	6.6
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	295.7	28.0	39.6	77.1	67.7	18.8	16.0	17.2	8.0	23.2
188	<i>Eugenia jambos</i> L.	291.0	47.0	78.2	69.8	34.7	28.7	5.2	19.4	0	7.9
131	<i>Mangifera indica</i> L.	234.8	5.6	2.9	4.7	26.4	16.5	2.7	15.8	21.3	121.4
161	<i>Calophyllum brasiliense</i> Camb.	201.3	5.9	0	28.2	45.6	11.6	36.9	25.1	24.3	23.6
48	<i>Ocotea leucoxylo</i> n (Sw.) Mez	200.7	16.0	37.9	93.4	14.5	26.4	0	12.5	0	0
7	<i>Euterpe globosa</i> Gaertn.	189.3	64.5	89.9	33.4	1.4	0	0	0	0	0
118	<i>Alchornea latifolia</i> Sw.	146.6	21.8	48.4	27.8	21.6	0	0	0	4.3	22.8
22	<i>Ficus laevigata</i> Vahl	144.3	23.6	22.0	0	6.3	16.4	11.8	29.5	0	34.7
76	<i>Hymenaea courbaril</i> L.	141.0	7.2	24.6	20.2	30.8	22.2	19.5	11.9	4.5	0
200	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	133.4	12.1	30.3	24.2	30.0	12.6	0	12.4	3.6	8.2
134	<i>Spondias mombin</i> L.	120.4	6.3	0	10.0	16.8	0	12.6	10.9	0	63.9
224	<i>Cordia sulcata</i> DC.	110.9	13.5	24.3	31.0	18.3	23.8	0	0	0	0
106	<i>Dacryodes excelsa</i> Vahl	110.2	0	19.4	0	37.7	12.4	29.7	0	11.0	0
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	109.8	11.1	6.8	14.0	36.0	13.9	8.7	12.8	0	6.4
163	<i>Clusia rosea</i> Jacq.	109.0	1.8	46.4	50.6	0	10.3	0	0	0	0
334	<i>Nectandra sintenisii</i> Mez (E)	106.4	12.6	7.0	11.6	15.5	6.4	5.8	0	0	47.4
228	<i>Petitia domingensis</i> Jacq.	103.8	21.3	2.4	0	60.6	14.0	0	0	5.5	0
17	<i>Artocarpus altitilis</i> (Parkinson) Fosberg	102.7	8.3	10.0	0	4.4	22.3	19.9	7.2	6.6	24.0
115	<i>Byrsonima coriacea</i> (Sw.) DC.	101.0	0	7.1	19.8	32.3	0	24.0	17.8	0	0
100	<i>Citrus sinensis</i> Osbeck	100.1	57.5	22.0	20.6	0	0	0	0	0	0
138	<i>Cupania americana</i> L.	98.0	17.7	21.4	18.7	10.8	0	29.4	0	0	0
180	<i>Buchenavia capitata</i> (Vahl) Eichl.	97.9	0	20.6	5.7	10.4	6.5	27.4	17.2	0	7.3
184	<i>Terminalia catappa</i> L.	87.3	0	7.9	5.9	0	7.5	33.1	32.9	0	0
151	<i>Montezuma speciosissima</i> Sessé & Moc.	83.8	0	0	40.5	5.1	9.9	19.5	8.8	0	0
87	<i>Ormosia krugii</i> Urban	81.0	18.5	36.7	25.8	0	0	0	0	0	0
211	<i>Micropholis chrysophylloides</i> Pierre	76.2	7.1	29.2	0	29.0	10.9	0	0	0	0
149	<i>Sloanea berteriana</i> Choisy	74.5	11.1	27.9	10.4	10.8	0	14.2	0	0	0
227	<i>Citharexylum fruticosum</i> L.	71.1	5.8	10.8	43.9	10.6	0	0	0	0	0
721	<i>Coffea dewevrei</i> Wildem. & T. Dur.	52.5	28.1	13.6	10.8	0	0	0	0	0	0
125	<i>Hura crepitans</i> L.	48.5	0	3.7	5.1	2.8	3.5	5.0	4.4	0	24.1
156	<i>Guazuma ulmifolia</i> Lam.	47.4	18.0	5.2	0	0	0	12.9	5.1	0	6.2
192	<i>Myrcia deflexa</i> (Poir.) DC.	40.7	2.5	0	0	15.5	0	0	0	22.7	0
169	<i>Casearia arborea</i> (L. C. Rich) Urban	38.9	14.6	24.3	0	0	0	0	0	0	0
1	<i>Cyathia arborea</i> (L.) J. E. Smith	37.8	21.7	16.1	0	0	0	0	0	0	0
193	<i>Myrcia splendens</i> (Sw.) DC.	37.2	5.5	31.7	0	0	0	0	0	0	0
249	<i>Terebraria resinosa</i> (Vahl) Sprague	33.8	24.2	9.6	0	0	0	0	0	0	0
170	<i>Casearia decandra</i> Jacq.	30.9	9.3	15.2	1.7	0	0	0	0	0	4.8
999	<i>Tetrazygia elaeagnoides</i> (Sw.) DC.	27.6	19.9	0	7.7	0	0	0	0	0	0
999	Unidentified species	26.8	1.7	6.1	0	12.3	0	0	6.5	0	0
	Other Species (149)	1 316.3	172.7	238.8	155.8	193.2	193.7	109.0	83.0	44.6	93.8
	Total	11 524.8	1374.9	1818.7	1831.5	1661.8	1333.9	965.8	757.8	481.6	1200.2

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

Table 22—Volume of growing stock on timberland by species and diameter class, Puerto Rico, 1990<sup>a</sup>

Species code	Species name	Diameter class (centimeters)									
		All classes	15	20	25	30	35	40	45	50	55+
----- Thousand cubic meters -----											
4	<i>Spathodea campanulata</i> Beauv.	964.6	91.2	147.2	180.0	131.5	118.4	49.0	41.6	58.2	134.9
9	<i>Guarea guidonia</i> (L.) Sleumer	762.9	88.3	115.1	98.0	113.0	114.0	31.1	59.1	57.4	86.9
	<i>Cecropia peltata</i> L.	737.3	27.4	83.7	127.1	102.2	134.3	72.2	74.2	52.7	63.5
	<i>Inga vera</i> Willd.	546.7	100.1	102.4	88.2	72.1	62.0	61.5	12.4	22.0	26.0
6	<i>Tabebuia heterophylla</i> (DC.) Britton	342.4	100.3	60.9	81.3	24.4	49.5	11.7	0	6.9	7.4
	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	328.2	0	5.2	22.2	5.9	53.2	14.0	22.9	22.8	170.4
1	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	289.1	26.9	39.6	77.1	62.8	18.8	16.0	16.8	8.0	23.2
	<i>Inga fagifolia</i> (L.)	276.3	40.9	46.5	32.6	91.5	19.9	37.2	5.6	0	2.0
	<i>Andira inermis</i> (W. Wright) H.B.K.	272.7	68.0	35.5	83.7	46.2	10.3	0	29.0	0	0
1	<i>Calophyllum brasiliense</i> Camb.	175.3	5.9	0	9.8	45.2	11.6	36.9	24.3	22.6	18.9
8	<i>Alchornea latifolia</i> Sw.	126.8	21.8	45.0	19.7	21.2	0	0	0	0	19.1
6	<i>Dacryodes excelsa</i> Vahl	108.0	0	19.4	0	37.7	12.4	27.5	0	11.0	0
8	<i>Eugenia jambos</i> L.	100.8	19.8	46.1	17.4	6.7	0	0	10.7	0	0
	<i>Hymenaea courbaril</i> L.	97.8	6.8	0	20.2	30.3	21.6	15.2	0	3.6	0
	<i>Ocotea leucoxydon</i> (Sw.) Mez	97.4	16.0	26.7	54.6	0	0	0	0	0	0
4	<i>Nectandra sintenisii</i> Mez (E)	89.8	12.6	7.0	11.6	5.2	6.4	5.4	0	0	41.5
4	<i>Cordia sulcata</i> DC.	84.1	9.6	16.5	30.8	14.0	13.2	0	0	0	0
4	<i>Spondias mombin</i> L.	83.3	0	0	9.0	16.8	0	11.9	5.9	0	39.7
2	<i>Clidemia umbrosa</i> (Sw.) Cogn.	81.2	8.6	6.3	21.5	0	26.3	18.5	0	0	0
2	<i>Zanthoxylum martinicense</i> (Lam.) DC.	78.9	2.8	6.4	3.1	34.6	13.5	7.3	11.0	0	0
4	<i>Terminalia catappa</i> L.	77.3	0	7.9	5.9	0	6.8	26.1	30.8	0	0
	<i>Artocarpus altilis</i> (Parkinson) Fosberg	75.0	8.3	10.0	0	4.2	17.2	14.7	0	6.6	14.0
	<i>Ficus laevigata</i> Vahl	74.4	0	20.1	0	6.2	0	11.8	18.9	0	17.4
8	<i>Petitia domingensis</i> Jacq.	72.7	20.8	0	0	48.9	3.1	0	0	0	0
3	<i>Clusia rosea</i> Jacq.	69.6	0	18.8	41.7	0	9.0	0	0	0	0
0	<i>Buchenavia capitata</i> (Vahl) Eichl.	67.3	0	8.5	5.1	10.4	0	26.9	16.3	0	0
1	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	65.4	3.5	18.1	15.7	10.2	5.4	12.5	0	0	0
8	<i>Cupania americana</i> L.	63.7	16.0	15.1	18.5	9.6	0	4.5	0	0	0
8	<i>Dipholis salicifolia</i> (L.) A. DC.	63.4	23.9	39.5	0	0	0	0	0	0	0
	<i>Ormosia krugii</i> Urban	61.9	14.7	21.6	25.5	0	0	0	0	0	0
9	<i>Sloanea berteriana</i> Choisy	58.8	7.6	27.9	0	10.1	0	13.3	0	0	0
3	<i>Pouteria multiflora</i> (A.DC.) Eyma	58.4	0	0	0	4.7	0	17.9	5.2	0	16.7
1	<i>Montezuma speciosissima</i> Sessé & Moc.	58.0	0	0	40.0	0	9.1	0	8.8	0	0
5	<i>Byrsonima coriacea</i> (Sw.) DC.	55.0	0	7.1	9.3	0	0	22.4	16.2	0	0
	<i>Lonchocarpus latifolius</i> (Willd.) H.B.K.	49.7	0	0	7.2	6.9	26.4	9.1	0	0	0
	<i>Albizia procera</i> (Roxb.) Benth.	46.9	3.7	22.6	0	20.6	0	0	0	0	0
8	<i>Cedrela odorata</i> L.	46.3	0	18.9	0	0	0	12.5	4.9	0	10.1
0	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	43.8	4.9	7.8	5.1	6.6	9.0	0	4.2	3.6	2.6
1	<i>Micropholis chrysophylloides</i> Pierre	40.4	6.8	18.6	0	14.9	0	0	0	0	0
1	<i>Mangifera indica</i> L.	37.1	5.0	0	4.7	0	4.6	0	0	4.1	18.6
2	<i>Myrcia deflexa</i> (Poir.) DC.	35.9	0	0	0	14.8	0	0	0	21.1	0
2	<i>Micropholis garciniaefolia</i> Pierre	32.5	0	0	6.2	0	15.7	10.6	0	0	0
	<i>Castilla elastica</i> Cervantes	31.0	0	4.2	13.9	12.9	0	0	0	0	0
3	<i>Spondias dulcis</i> Parkinson	28.9	0	0	0	0	19.2	0	0	9.7	0
0	<i>Casearia decandra</i> Jacq.	28.3	9.3	15.2	0	0	0	0	0	0	3.8
9	<i>Casearia arborea</i> (L. C. Rich) Urban	26.6	14.1	12.5	0	0	0	0	0	0	0
1	<i>Coffea dewevrei</i> Wildem. & T. Dur.	26.4	16.0	0	10.4	0	0	0	0	0	0
0	<i>Citrus sinensis</i> Osbeck	25.7	12.5	5.1	8.1	0	0	0	0	0	0
	Other species	504.3	83.6	88.7	100.0	99.4	39.7	21.9	21.1	10.3	39.8
	Total	7636.9	939.5	1200.7	1311.9	1141.7	862.9	619.6	439.9	320.6	761.3

Columns may not add up to totals as a result of rounding numbers.

Table 23—Volume of sawtimber on timberland by species and diameter class, Puerto Rico, 1990<sup>a</sup>

Species code	Species name	Diameter class (centimeters)						
		All classes	30	35	40	45	50	55+
----- Thousand cubic meters -----								
234	<i>Spathodea campanulata</i> Beauv.	399.5	87.5	82.7	35.3	31.6	43.6	106.7
20	<i>Cecropia peltata</i> L.	349.4	60.6	91.9	48.7	59.5	34.0	54.7
109	<i>Guarea guidonia</i> (L.) Sleumer	272.7	54.4	58.7	17.6	38.6	37.5	65.9
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	243.3	3.2	33.7	11.2	17.5	19.1	148.2
64	<i>Inga vera</i> Willd.	140.5	33.0	33.8	30.0	7.3	14.2	22.2
62	<i>Inga fagifolia</i> (L.)	93.6	49.8	13.4	24.6	4.5	0	1.3
201	<i>Didymopanax morototoni</i> (Aubl.) Decne & Planch.	89.7	39.3	9.1	9.2	8.4	5.8	18.0
161	<i>Calophyllum brasiliense</i> Camb.	83.7	17.2	3.3	21.0	15.2	14.7	12.4
236	<i>Tabebuia heterophylla</i> (DC.) Britton	62.5	15.2	29.6	8.5	0	4.3	4.9
106	<i>Dacryodes excelsea</i> Vahl	56.2	20.2	9.6	20.0	0	6.4	0
81	<i>Andira inermis</i> (W. Wright) H.B.K.	54.5	28.8	4.9	0	20.8	0	0
184	<i>Terminalia catappa</i> L.	51.9	0	5.8	17.7	28.3	0	0
134	<i>Spondias mombin</i> L.	46.1	6.4	0	6.6	3.7	0	29.4
334	<i>Nectandra sintenisii</i> Mez (E)	43.8	3.0	4.8	2.1	0	0	33.8
76	<i>Hymenaea courbaril</i> L.	43.7	16.7	14.0	9.3	0	3.6	0
213	<i>Pouteria multiflora</i> (A. DC.) Eyma	41.5	2.3	0	9.9	2.8	0	14.6
17	<i>Artocarpus altiss</i> (Parkinson) Fosberg	39.4	1.7	10.8	10.3	0	5.5	11.1
180	<i>Buchenavia capitata</i> (Vahl) Eichl.	31.8	9.0	0	14.1	8.7	0	0
22	<i>Ficus laevigata</i> Vahl	30.4	1.9	0	4.6	11.9	0	12.1
592	<i>Clitellium umbrosa</i> (Sw.) Cogn.	27.6	0	15.4	12.2	0	0	0
115	<i>Byrsonima coriacea</i> (Sw.) DC.	27.1	0	0	15.8	11.3	0	0
102	<i>Zanthoxylum martinicense</i> (Lam.) DC.	27.0	14.2	6.1	2.0	4.7	0	0
228	<i>Pettitia domingensis</i> Jacq.	22.3	20.6	1.7	0	0	0	0
86	<i>Lonchocarpus latifolius</i> (Willd.) H.B.K.	20.0	5.2	11.0	3.8	0	0	0
118	<i>Alchornea latifolia</i> Sw.	19.9	9.0	0	0	0	0	10.9
131	<i>Mangifera indica</i> L.	18.1	0	1.3	0	0	0	14.6
221	<i>Cordia alliodora</i> (Ruiz & Pav.) Oken	17.8	5.6	3.1	9.1	0	2.3	0
108	<i>Cedrela odorata</i> L.	17.4	0	0	7.2	3.6	0	6.6
192	<i>Myrcia deflexa</i> (Poir.) DC.	17.0	6.1	0	0	0	10.9	0
63	<i>Inga quaternata</i> Poepp. & Endl.	15.6	0	0	0	0	0	15.6
224	<i>Cordia sulcata</i> DC.	15.1	6.7	8.4	0	0	0	0
260	<i>Pinus caribaea</i> Morelet	13.1	0	0	7.5	0	0	0
133	<i>Spondias dulcis</i> Parkinson	12.9	0	7.6	0	0	5.3	0
212	<i>Micropholis garciniaefolia</i> Pierre	12.6	0	8.1	4.6	0	0	0
19	<i>Castilla elastica</i> Cervantes	12.5	0	0	0	0	0	0
149	<i>Sloanea berteriana</i> Choisy	11.7	5.0	0	6.7	0	0	0
188	<i>Eugenia jambos</i> L.	11.5	4.9	0	0	6.5	0	0
209	<i>Munikara bidentata</i> (A. DC.) Chev.	11.4	7.8	3.5	0	0	0	0
200	<i>Dendropanax arboreus</i> (L.) Decne. & Planch.	11.3	1.9	3.4	0	2.1	2.0	2.0
340	<i>Phoebe elongata</i> (Vahl) Nees	10.4	0	7.8	0	0	0	2.6
144	<i>Meliosma herbertii</i> Rolfe	10.3	10.3	0	0	0	0	0
151	<i>Montezuma speciosissima</i> Sessé & Moc.	10.2	0	2.1	0	8.2	0	0
164	<i>Mammea americana</i> L.	9.5	0	0	3.4	0	0	6.2

Species code	Species name	Diameter class (centimeters)						
		All classes	30	35	40	45	50	55+
----- Thousand cubic meters -----								
330	<i>Nectandra antillana</i> Meisn.	9.2	3.2	0	0	0	6.0	0
211	<i>Micropholis chrysophylloides</i> Pierre	7.9	7.9	0	0	0	0	0
999	Unidentified species	7.2	4.1	0	0	3.1	0	0
203	<i>Rapanea ferruginea</i> (Ruiz & Pav.) Mez	6.9	0	6.9	0	0	0	0
34	<i>Magnolia portoricensis</i> Bello	6.8	0	6.8	0	0	0	0
61	<i>Albizia procera</i> (Roxb.) Benth.	5.5	5.5	0	0	0	0	0
155	<i>Quararibaea turbinata</i> (Sw.) Poir.	5.2	0	0	0	5.2	0	0
163	<i>Clusia rosea</i> Jacq.	5.1	0	5.1	0	0	0	0
138	<i>Cupania americana</i> L.	4.4	2.5	0	2.0	0	0	0
230	<i>Vitex divaricata</i> Sw.	4.0	0	2.5	1.5	0	0	0
51	<i>Persea americana</i> Mill.	3.9	3.9	0	0	0	0	0
107	<i>Tetragastris balsamifera</i> (Sw.) Kuntze	3.9	0	0	0	0	0	3.9
227	<i>Citharexylum fruticosum</i> L.	3.5	3.5	0	0	0	0	0
125	<i>Hura crepitans</i> L.	3.4	1.9	0	0	0	0	1.5
105	<i>Bursera imaruba</i> (L.) Sarg.	3.2	3.2	0	0	0	0	0
187	<i>Eugenia aeruginosa</i> DC.	2.7	0	0	0	2.7	0	0
154	<i>Ochroma pyramidale</i> (Cav.) Urban	2.3	2.3	0	0	0	0	0
170	<i>Casearia decandra</i> Jacq.	1.6	0	0	0	0	0	1.6
137	<i>Turpinia paniculata</i> Vent.	1.5	0	0	1.5	0	0	0
612	<i>Miconia subcorymbosa</i> Britton	1.2	0	0	0	0	0	1.2
74	<i>Cassia stamea</i> Lam.	.9	0	.9	0	0	0	0
554	<i>Calyptrothrix pallens</i> Griseb.	.4	0	0	0	0	0	.4
Total		2647.2	598.0	507.8	378.0	306.2	215.2	602.4

<sup>a</sup> Columns may not add up to totals as a result of rounding numbers.

**Table 24—Volume of sawtimber on timberland by species and butt log grade, Puerto Rico, 1990<sup>a</sup>**

Species code	Species name	All classes	Butt log grade <sup>b</sup>			Volume in ungraded portion
			1	2	3	
----- Thousand cubic meters -----						
234	<i>Spathodea campanulata</i> Beauv.	399.5	9.3	154.6	191.0	44.2
20	<i>Cecropia peltata</i> L.	349.4	50.3	165.1	114.1	19.9
109	<i>Guarea guidonia</i> (L.) Sleumer	272.7	6.8	52.7	96.5	116.7
84	<i>Erythrina poeppigiana</i> (Walp.) O. F. Cook	243.3	73.7	90.4	58.8	20.4
64	<i>Inga vera</i> Willd.	140.5	7.8	32.7	77.2	22.8
62	<i>Inga fagifolia</i> (L.) Willd.	93.6	5.2	33.1	50.8	4.5
201	<i>Didymopanax morototoni</i> (Aubl.) Decne. & Planch.	89.7	16.4	25.0	48.3	0
161	<i>Calophyllum brasiliense</i> Camb.	83.7	0	16.7	62.1	4.9
236	<i>Tabebuia heterophylla</i> (DC.) Britton	62.5	0	5.9	36.3	20.3
106	<i>Dacryodes excelsa</i> Vahl	56.2	0	28.2	28.0	0
81	<i>Andira inermis</i> (W. Wright) H.B.K.	54.5	0	18.4	33.8	2.3
184	<i>Terminalia catappa</i> L.	51.9	0	11.3	26.4	14.2
134	<i>Spondias mombin</i> L.	46.1	7.9	25.3	12.8	0
334	<i>Nectandra sintenisii</i> Mez (E)	43.8	0	30.5	3.0	10.3
	Other species	659.8	8.9	238.8	289.1	123.0
	Total	2647.2	186.3	928.7	1128.2	404.0

<sup>a</sup>Columns may not add up to totals as a result of rounding numbers.

<sup>b</sup>Graded by hardwood log grade standards used in the Southern United States.

**Table 25—Volume of timber, growing stock, and sawtimber on timberland by stand-size class, Puerto Rico, 1990**

Stand-size class	Volume of timber	Volume of growing stock	Volume of sawtimber
----- Thousand cubic meters -----			
Sawtimber	7 031.2	4615.1	2060.9
Poletimber	4 244.5	2928.0	572.2
Sapling-seedling	248.9	93.8	14.3
Total	11 524.6	7636.9	2647.4

**Table 26—Volume of timber on timberland by Life Zone, soil group, and slope class, Puerto Rico, 1990**

Life Zone and soil group	Percent slope class				
	groups	All 0 to 10	11 to 25	26 to 45	> 45
----- <i>Thousand cubic meters</i> -----					
<b>Subtropical Moist Forest</b>					
Deep volcanic soils	1 377.8	287.1	174.4	487.8	428.5
Shallow volcanic soils	1 178.0	134.5	315.0	361.6	366.8
Granitic soils	977.0	0	144.3	210.4	642.2
Limestone soils	953.1	123.0	163.3	156.4	510.5
Total	4 505.9	544.6	797.0	1216.2	1948.0
<b>Subtropical Wet Forest</b>					
Deep volcanic soils	4 787.8	282.6	787.1	1435.0	2283.1
Shallow volcanic soils	596.0	0	88.0	141.8	366.2
Granitic soils	1 493.5	0	224.2	309.4	959.9
Limestone soils	141.5	0	42.9	98.6	0
Total	7 018.8	282.6	1142.2	1984.8	3609.2
<b>Total, all Life Zones</b>	11 524.7	827.2	1939.2	3201.0	5557.2



**Franco, Peter A.; Weaver, Peter L.; Eggen-McIntosh, Susan.** 1997. Forest resources of Puerto Rico, 1990. Resour. Bull. SRS-22. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southern Research Station. 45 p.

The principal findings of the second forest survey of Puerto Rico (1990) and changes that have occurred since the survey was established in 1980 are presented. The forest inventory estimates describe the timber resource found within the potential commercial region designated in the first survey. The timber resource addressed consists primarily of regrown areas on abandoned pastures and cropland, including coffee production areas. The status and trends of the timber resource are presented for the two Life Zones occurring in the commercial region, as well as for various forest classes, which are based on stand history and origin. Topics discussed include forest area, timberland area, basal area, species composition, timber volume, growing-stock volume, and sawtimber volume. Results of the 1990 survey are promising, showing increases in numbers of trees across all diameter classes and substantial increases in volume. These trends offer evidence that Puerto Rico's forests are continuing to recover following a dramatic decline of the late 19<sup>th</sup> and early 20<sup>th</sup> centuries.

**Keywords:** Coffee shade forests, forest area, forest inventory, secondary forests, timber volume, tropical forest management.



The Forest Service, U.S. Department of Agriculture, is dedicated to the principle of multiple use management of the Nation's forest resources for sustained yields of wood, water, forage, wildlife, and recreation. Through forestry research, cooperation with the States and private forest owners, and management of the National Forests and National Grasslands, it strives—as directed by Congress—to provide increasingly greater service to a growing Nation.

The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs). Persons with disabilities who require alternative means for communication of program information (braille, large print, audiotape, etc.) should contact the USDA's TARGET Center at 202-720-2600 (voice and TDD). To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, DC 20250, or call 1-800-245-6340 (voice) or 202-720-1127 (TDD). USDA is an equal employment opportunity employer.