



United States
Department of
Agriculture



Forest Service

Southern Forest
Experiment Station

Research Note

SO-347
June 1988

Volume Equations for Plantation Cottonwood Trees (*Populus deltoides*)

Roger M. Krinard

SUMMARY

Equations and tables for cubic-foot volume of outside bark to the top of the tree and inside bark to a 3-inch top are given for plantation-grown cottonwoods (*Populus deltoides* Bartr. ex Marsh.) from 5 to 22 inches in d.b.h.

INTRODUCTION

A previous publication of volume tables for plantation-grown cottonwood (*Populus deltoides* Bartr. ex Marsh.) trees was based on 650 3- or 5-year-old trees sampled from 3 plantations near Greenville, Mississippi (Mohn and Krinard 1971). Diameters ranged from 2.3 to 9.9 inches, and heights ranged from about 25 to 65 feet. Volume equations have now been developed from planted cottonwoods near Fidler, Mississippi, that are up to 20 years old and more than 20 inches in d.b.h. This expands the range of tree sizes used in cottonwood volume determinations.

SAMPLE TREES

The sample trees came from a spacing study in the Mississippi River batture on Crown Zellerbach land (Krinard and Johnson 1984). Soil series were mainly Commerce and Convent silt loams, which have estimated site indices at age 30 of 105 to 125 feet for natural cottonwood stands (Broadfoot 1976). Cuttings used in the plantings were run-of-the-bar stock (nonselect material). Spacings were 4 by 9, 8 by 9, 12 by 12, and 16 by 18 feet. Several thinning treatments were applied, starting after the 4th year and ending after the 15th year.

Volumes were obtained both from trees cut in thinnings, with diameters measured at 4-foot intervals start-

ing with a 1-foot stump, and from standing trees, using a dendrometer and also starting with a 1-foot stump. Whether trees were cut or standing, volumes were determined by summing 4-foot frustrums (Grosenbaugh 1954), with the top calculated as a cone. Two bark thicknesses were measured at d.b.h. Inside bark diameters (d.i.b.) were calculated from outside bark diameters (d.o.b.) by $d.i.b. = d.o.b. (1 - (1 - d.b.h.i.b./d.b.h.o.b.)(1/(2 - d.o.b./d.b.h.o.b.)))$ (Mesavage 1969).

Distribution of sample trees by age, spacing, and whether cut or standing are given in table 1. Tree distribution by 1-inch diameter class and 10-foot height class is given in table 2 with the average measured total outside bark volume.

Tables for total stem cubic-foot volume outside bark from a 1-foot stump to the top of the tree (table 3) and inside bark stem volume to a 3-inch top (table 4) were constructed from the following regression equations:

$$\begin{aligned} \text{Volume outside bark} &= 0.06 + 0.002221 D^2H \\ r^2 &= 0.987, S_e = 2.4 \text{ ft}^3, \text{ CV} = 10.5\%, \bar{V} = 23.1 \text{ ft}^3 \\ \text{Volume inside bark} &= -0.86 + 0.001904 D^2H \\ r^2 &= 0.987, S_e = 2.1 \text{ ft}^3, \text{ CV} = 10.9\%, \bar{V} = 18.9 \text{ ft}^3 \\ &\text{where } D = \text{d.b.h. and } H = \text{total height.} \end{aligned}$$

Intercept and slope coefficients based on smaller, younger trees (Mohn and Krinard 1971) were 0.21 and 0.00221 for outside bark volumes and -0.62 and 0.00204 for inside bark volumes.

A weighted regression may be used to correct for nonhomogeneity of variance and may give a more precise estimate. Use of weights, whether $1/(D^2H)$ or $1/(D^2H)^2$, provided the same or very slightly larger values of S_e , CV, and fit index $(1 - \sum(Y - \hat{Y})^2 / \sum(Y - \bar{Y})^2)$ with this data. The index of fit (Furnival 1961) was slightly better for weighted regressions, with values of 1.6 and 1.3 compared to 2.4 for outside bark volumes and 1.4 and 1.2 compared to 2.1 for inside bark volumes. Weighted regressions, with



weight $1/(D^2H)^2$, changed the coefficients slightly, as follows:

$$\text{Volume outside bark} = 0.09 + 0.002216 D^2H$$

$$\text{Volume inside bark} = -0.41 + 0.001852 D^2H$$

The volumes generated by these equations should be applicable to most commercial cottonwood plantation spacings. The equations represent trees from a mixture of four planting spacings, with 20, 26, 31, and 23 percent of sample trees coming from 4- by 9-, 8- by 9-, 12- by 12-, and 16- by 18-foot spacings, respectively. There were differences between spacings in volume equations, partly due to the number of trees measured and to differences in tree

sizes between spacings, where mean measured outside bark volumes were 19.0, 19.4, 23.6, and 30.2 cubic feet from closest to widest spacing. Spacing variables incorporated in the volume equations improved the fit statistics so very little that there was no useful benefit in doing so. Whether these equations would apply to cottonwood trees of genetically improved material—which would include most current and future plantations—is not known, but with slash (Tankersley and others 1983) and loblolly (Buford and Burkhart 1987) pine, improved and unimproved trees did not need separate equations.

Table 1 — Number of sample trees by age, spacing, and whether cut or standing

Spacing	Age of cut trees					Age of standing trees					Standing and cut tree total
	6	7	8	12	15	7	9	10	15	20	
Feet	-----										Number
4 by 9	20			11	19	7	26	12		7	102
8 by 9	30	7	5	10	15	14	23	16	3	8	131
12 by 12	32	10	20	6	10	11	23	35	3	10	160
16 by 18		10	10	1	10	4	37	38	2	7	119
Total	82	27	35	28	54	36	109	101	8	32	512

Table 2. — Number of trees and average measured cubic-foot volume outside bark from a 1-foot stump to the top of the tree by 10-foot total height class and 1-inch diameter class

D.b.h.	Height classes								Total
	40-49	50-59	60-69	70-79	80-89	90-99	100-109	110-119	
Inches	-----								Cubic feet
4	1.8(1)	2.6(6)	3.9(1)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	(8)
5	0.0(0)	3.6(16)	4.8(6)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	(22)
6	0.0(0)	5.0(9)	6.2(43)	7.4(11)	9.1(1)	0.0(0)	0.0(0)	0.0(0)	(64)
7	0.0(0)	0.0(0)	7.7(21)	9.4(24)	11.6(3)	0.0(0)	0.0(0)	0.0(0)	(48)
8	0.0(0)	0.0(0)	9.5(8)	12.0(37)	15.0(12)	16.8(5)	0.0(0)	0.0(0)	(62)
9	0.0(0)	0.0(0)	14.4(1)	15.0(26)	16.1(15)	21.7(2)	0.0(0)	0.0(0)	(44)
10	0.0(0)	0.0(0)	0.0(0)	18.4(20)	20.9(36)	24.7(12)	24.6(1)	0.0(0)	(69)
11	0.0(0)	0.0(0)	0.0(0)	21.6(12)	24.9(40)	27.4(11)	30.3(3)	0.0(0)	(66)
12	0.0(0)	0.0(0)	0.0(0)	23.5(6)	27.9(27)	32.4(14)	36.7(2)	0.0(0)	(49)
13	0.0(0)	0.0(0)	0.0(0)	27.8(3)	31.4(11)	38.5(6)	43.2(3)	0.0(0)	(23)
14	0.0(0)	0.0(0)	0.0(0)	0.0(0)	36.4(3)	40.0(2)	0.0(0)	0.0(0)	(5)
15	0.0(0)	0.0(0)	0.0(0)	0.0(0)	38.5(2)	45.2(2)	58.0(5)	65.4(2)	(11)
16	0.0(0)	0.0(0)	0.0(0)	0.0(0)	45.2(1)	48.9(2)	67.4(4)	74.7(1)	(8)
17	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	60.8(1)	73.0(11)	78.6(1)	(13)
18	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	79.4(2)	91.8(4)	(6)
19	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	90.0(1)	97.4(3)	(4)
20	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	90.8(4)	100.9(3)	(7)
21	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	118.5(1)	(1)
22	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	0.0(0)	121.9(2)	(2)
Total	(1)	(31)	(80)	(139)	(151)	(57)	(36)	(17)	(512)

† Example: the 4-inch d.b.h. class represents trees whose d.b.h.'s range from 4.0 to 4.9 inches.

‡ Numeral in parentheses represents the number of trees in this height and d.b.h. class.

Roger M. Krinard is mensurationist at the Southern Hardwoods Laboratory, maintained at Stoneville, Mississippi, by the Southern Forest Experiment Station, Forest Service—USDA, in cooperation with the Mississippi Agricultural and Forestry Experiment Station and the Southern Hardwood Forest Research Group.

The use of trade or firm names in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture of any product or service.



Table 3 — Cubic-foot volumes outside bark from a 1-foot stump to the top of the tree

D.b.h. Inches	Total tree height (ft)							
	50	60	70	80	90	100	110	120
5	2.8	3.4	3.9	4.5				
6	4.1	4.9	5.7	6.5	7.3			
7	5.5	6.6	7.7	8.8	9.9			
8	7.2	8.6	10.0	11.4	12.9	14.3		
9	9.1	10.9	12.7	14.5	16.3	18.0		
10		13.4	15.6	17.8	20.0	22.3	24.5	
11		16.2	18.9	21.6	24.2	26.9	29.6	
12		19.2	22.4	25.6	28.8	32.0	35.2	
13		22.6	26.3	30.1	33.8	37.6	41.3	
14			30.5	34.9	39.2	43.6	47.9	
15			35.0	40.0	45.0	50.0	55.0	60.0
16			39.9	45.5	51.2	56.9	62.6	68.3
17				51.4	57.8	64.2	70.7	77.1
18					64.8	72.0	79.2	86.4
19					72.2	80.2	88.3	96.3
20					80.0	88.9	97.8	106.7
21						98.0	107.8	117.6
22						107.6	118.3	129.1

† Area outlined represents range of measured trees.

Table 4 — Cubic-foot volumes inside bark from a 1-foot stump to a 3-inch top

D.b.h. Inches	Total tree height (ft)							
	50	60	70	80	90	100	110	120
5	1.5	2.0	2.5	2.9				
6	2.6	3.3	3.9	4.6	5.3			
7	3.8	4.7	5.7	6.6	7.5			
8	5.2	6.5	7.7	8.9	10.1	11.3		
9	6.9	8.4	9.9	11.5	13.0	14.6		
10		10.6	12.5	14.4	16.3	18.2	20.1	
11		13.0	15.3	17.6	19.9	22.2	24.5	
12		15.6	18.3	21.1	23.8	26.6	29.3	
13		18.4	21.7	24.9	28.1	31.3	34.5	
14			25.3	29.0	32.7	36.5	40.2	
15			29.1	33.4	37.7	42.0	46.3	50.5
16			33.3	38.1	43.0	47.9	52.8	57.6
17				43.2	48.7	54.2	59.7	65.2
18					54.7	60.8	67.0	73.2
19					61.0	67.9	74.7	81.6
20					67.7	75.3	82.9	90.5
21						83.1	91.5	99.9
22						91.3	100.5	109.7

† Area outlined represents range of measured trees.

LITERATURE CITED


Broadfoot, Walter M. 1976. Hardwood suitability for and properties of important Midsouth soils. Res. Pap. SO-127. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 84 p.

Buford, Marilyn A.; Burkhart, Harold E. 1987. Genetic improvement effects on growth and yield of loblolly pine plantations. Forest Science. 33(3): 707-724.

Furnival, George M. 1961. An index for comparing equations used in constructing volume tables. Forest Science. 7(4): 337-341.

Grosenbaugh, L. R. 1954. New tree measurement concepts: height accumulation, giant tree, taper and shape. Occas. Pap. 134. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 32 p.

Krinard, Roger M.; Johnson, Robert L. 1984. Cottonwood plantation growth through 20 years. Res. Pap. SO-212. New Orleans, LA: U.S. Department of Agriculture,



- Forest Service, Southern Forest Experiment Station. 11 p.
- Mesavage, Clement. 1969. Converting dendrometer estimates of outside-bark stem diameters to wood diameters on major southern pines. Res. Note SO-93. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 p.
- Mohn, C.A.; Krinard, R. M. 1971. Volume tables for small cottonwoods in plantations. Res. Note SO-113. New Orleans, LA: U.S. Department of Agriculture, Forest Service, Southern Forest Experiment Station. 4 p.
- Tankersley, Larry; Bongarten, Bruce; Brister, Graham; Zoerb, Marvin. 1983. Operational plantations of improved slash pine: Age 15 results. In: Proceedings, 17th Southern Forest Tree Improvement Conference; 1983 June 7-9; Athens, GA. Springfield, VA: National Technical Information Service: 271-280.

