

NTFPs from Trees: Nontimber Forest Products that Support our Society and Economy

PINYON PINE

Product: Pine nuts
Plant parts used: Seeds



Pinyon pine (*Pinus edulis* Engelm.) is a long-lived, drought-tolerant tree common throughout the Southwestern United States. The pinyon-juniper forest type is generally located on rocky plateaus and lower mountain slopes. It is typically found at elevations 4,500–6,000 feet above sea level. Forests in which pinyon pine is the major pine species cover close to 40 million acres primarily in Arizona, Colorado, New Mexico, and Utah (shown in green on the map below).



Key Points

- The principal contemporary use of pinyon pine is for food. Pine “nuts” are the seeds from trees, which can produce for hundreds of years.
- There is a multi-million-dollar global market for pine nuts; the United States is a net importer.
- In recent years, there has been a negative trend in overall change in cubic feet per acre of pinyon pine.
- Management of pinyon-juniper woodlands entails wildfire management, grazing, and control of invasive species and grassland invasion.
- Large-scale mortality of pinyon pine is expected as climate change impacts increase.

Nontimber Uses

- Native Americans depended on pinyon pine nuts for survival, especially during the winter when food was scarce. Pinyon pine is a cultural keystone species for Tribes of the Southwestern United States.
- In the mid-1800s, charcoal made from pinyon wood fueled smelters to process silver mined in the Southwest.
- Pinyon pine has been a staple Christmas tree in States with large pinyon-juniper forests.
- Pinyon pine nuts are delicacies. They contain about 20 amino acids, high levels of iron and vitamin A, and >2,800 calories per pound. Consumption can contribute to livelihood and food security.



Markets

- In the late 1980s, >1 million pounds of seed were harvested each year by commercial operators. There is little current information about annual harvest amounts.
- In 2004, the U.S. market for pine nuts was reportedly valued at about \$100 million.
- Commercial importance of pine nuts is underestimated, so pinyon woodlands are not managed for pine nuts. 80 percent of pine nuts consumed in the United States are imported.
- The average annual value of imported pine nuts for the years 2009–2013 was approximately \$25 million. Over the same period, the United States exported an estimated \$922,000 annually.
- Pine nuts are one of the most expensive nuts on the market, with prices in 2004 ranging from \$10–16 per pound.

Any medical or pesticide use described in this publication is for reader information and does not imply endorsement by the U.S. Department of Agriculture.

Acknowledgments: Literature review and tree species maps supporting this Science Update were contributed by Ben Addlestone, Thomas Metzger, Wenyu Gao, and John Munsell through a collaboration with Virginia Polytechnic Institute and State University.

Photo Credits: ABOVE: Dave Powell, USDA Forest Service (retired), Bugwood.org.
NEXT PAGE TOP TO BOTTOM: USD01 National Park Service; USDA Forest Service.



Status

- According to Forest Inventory and Analysis (FIA) data, the mean annual number of trees per acre ranged from 0.0057 (Nevada) to 58.42 (New Mexico) over the years 2005–2017.
- From 2005 to 2017, net change (i.e., difference between growth and sum of mortality and removals) trended negative in Arizona, Colorado, New Mexico, and Utah; Nevada and Wyoming had barely positive net change.
- Arizona had the greatest negative net change (a loss of >3 cubic feet per acre) and the highest mortality (~2.07 cubic feet per acre) between 2005 and 2017.
- In general, the negative net change has been decreasing and trending toward zero over the years [see chart (A)].
- There has been a positive trend in estimated mean annual growth of pinyon pine, although estimated growth is still negative [see chart (B)].

Management and Implications

- Pinyon pine trees reach cone-bearing age at around 25 years when the trees are 4–9 meters tall. Seed cone production is highly correlated with late summer temperatures at the time of initial cone formation.
- A good stand of pinyon pines can produce about 400 pounds of seed per acre, but seeds are not produced in large quantities until the tree is 75–100 years old.
- Pinyon pine is shade intolerant and does not compete well with grasses. Disturbance, especially fire, is needed for the forest type to flourish.
- Higher temperatures and lower precipitation associated with climate change could cause shifts in species composition, structure, and distribution of pinyon-juniper woodlands; greater infestations from the pinyon Ips beetle (*Ips confusus* LeConte); and large-scale mortality.
- Long-lived mast species like pinyon pine have shown decline in cone production from 1970s to the 2000s in New Mexico and northwest Oklahoma.
- Additional analytical methods may be needed to capture regional ecological variations in discrete populations.

References

- Evans, R.A. 1988. Management of pinyon-juniper woodlands. Gen. Tech. Rep. INT-249. Ogden, UT: U.S. Department of Agriculture Forest Service, Intermountain Research Station. 34 p.
- Fagan, D. 2017. Pinyon pine trees: icons of the Southwest. <https://www.desertusa.com/flora/pinyon-pine.html>. [Date accessed: January 16, 2018].
- Kyllo, R. 2016. Piñon-juniper woodlands of the Western United States: Are we on the brink of piñon oblivion? Flagstaff, AZ: Northern Arizona University. 52 p. Professional paper.
- Lanner, R.M. 1981. The piñon pine: a natural and cultural history. Reno, NV: University of Nevada Press. 208 p.
- Redmond, M.D.; Forcella, F.; Barger, N.N. 2012. Declines in pinyon cone production associated with regional warming. *Ecosphere*. 3(12): 120. <https://doi.org/10.1890/ES12-00306.1>.
- Ronco, F.P., Jr. 1990. Pinyon. In: Burns, R.M.; Honkala, B.H., tech. coords. *Silvics of North America: volume 1. Conifers*. Agric. Handb. 654. Washington, DC: U.S. Department of Agriculture Forest Service: 327–337.
- Sharashkin, L.; Gold, M. 2004. Pine nuts: species, products, markets and potential for U.S. production. In: Northern Nut Growers Association 95th annual report. Proceedings for the 95th annual meeting; Columbia, Missouri, August 16–19, 2004. <https://pinenut.com/growing-pine-nuts/ringing-cedar-nuts-oil.shtml>. [Date last accessed: October 27, 2020].
- Shaw, J.D.; Steed, B.E.; DeBlander, L.T. 2005. Forest Inventory and Analysis (FIA) annual inventory answers the question: What is happening to pinyon-juniper woodlands? *Journal of Forestry*. 10(36): 280–286.
- Wagstaff, F.J. 1987. Economics of managing pinyon-juniper lands for woodland products. In: Everett, R.L., comp. Proceedings—pinyon-juniper conference. Gen. Tech. Rep. INT-215. Ogden, UT: U.S. Department of Agriculture Forest Service, Intermountain Research Station: 168–172.

Citation: Chamberlain, J. 2020. NTFPs from trees: pinyon pine. Science Update SRS-139. Asheville, NC: U.S. Department of Agriculture Forest Service, Southern Research Station. 2 p. <https://doi.org/10.2737/SRS-SU-139>.

Learn more about nontimber forest products: Jim Chamberlain • james.l.chamberlain@usda.gov • <https://www.srs.fs.usda.gov/staff/524>

www.srs.fs.usda.gov/research/nontimber-forest-products/